

GEHRING–HAYMAN THEOREM FOR CONFORMAL DEFORMATIONS

PEKKA KOSKELA AND PÄIVI LAMMI

ABSTRACT. We study conformal deformations of a uniform space that satisfies the Ahlfors Q -regularity condition on balls of Whitney type. We verify the Gehring–Hayman Theorem by using a Whitney Covering of the space.

1. INTRODUCTION

Given $x, y \in B^2(0, 1)$, the hyperbolic geodesic $[x, y]$ is essentially the shortest curve joining x to y in $B^2(0, 1)$. More precisely

$$\ell([x, y]) \leq \frac{\pi}{2} \ell(\gamma)$$

whenever γ is a path that joins x to y in $B^2(0, 1)$. This simple fact is an instance of a theorem of Gehring and Hayman in [GH]: If $f: B^2(0, 1) \rightarrow \Omega \subset \mathbb{C}$ is a conformal mapping and γ is a path joining points x and y , then

$$(1.1) \quad \int_{[x, y]} |f'(z)| ds \leq C \int_{\gamma} |f'(z)| ds,$$

where $C \geq 1$ is an absolute constant. The density $\rho(z) = |f'(z)|$ satisfies a Harnack inequality

$$\frac{\rho(z)}{A} \leq \rho(w) \leq A\rho(z)$$

whenever $z \in B^2(0, 1)$ and $w \in B(z, (1 - |z|)/2)$. It also satisfies the area growth estimate

$$\int_{B_{\rho}(w, r)} \rho^2(z) dA \leq \pi r^2,$$

where $B_{\rho}(w, r)$ refers to the ball with centre w and radius r in the path metric

$$d_{\rho}(x, y) = \inf \int_{\gamma} \rho ds,$$

where the infimum is taken over all curves γ joining points x and y .

In [BKR] the Gehring–Hayman inequality (1.1) was extended to $B^n(0, 1)$, $n \geq 2$, for conformal deformations of the Euclidean metric. By a conformal deformation

2000 *Mathematics Subject Classification.* Primary 30C65.

Key words and phrases. Conformal deformations, uniform space, Whitney covering.

Both authors were supported by the Academy of Finland, grant no. 120972.

(a conformal density) ρ we mean a continuous function $\rho: B^n(0, 1) \rightarrow (0, \infty)$ that satisfies a Harnack inequality with a constant $A \geq 1$,

$$\frac{\rho(x)}{A} \leq \rho(w) \leq A\rho(x) \quad \text{for all } w \in B(x, (1 - |x|)/2) \text{ and all } x \in B^n(0, 1),$$

and a volume growth condition with a constant $B > 0$,

$$\int_{B_\rho(w, r)} \rho^n(z) dm_n \leq Br^n \quad \text{for all } w \in B^n(0, 1) \text{ and all } r > 0,$$

with respect to n -dimensional Lebesgue measure m_n .

Subsequently, Herron showed in [H1] that $B^n(0, 1)$ can be replaced by any uniform space (Ω, d) of bounded geometry. In this setting conformal densities are defined by conditions analogous to those given above — see Section 2 for details. Here uniformity is a substitute for the “roundness” of $B^n(0, 1)$. The assumption of bounded geometry includes two conditions. First, it requires that Ω carries a Borel regular measure μ that satisfies *the (Ahlfors) Q -regularity condition on balls of Whitney type* for some $Q > 1$. That is, there is a constant $C_1 \geq 1$ such that if $r \leq d(x, \partial\Omega)/2$, then

$$C_1^{-1}r^Q \leq \mu(B(x, r)) \leq C_1r^Q.$$

Secondly, it requires that balls $B(x, d(x, \partial\Omega)/2)$ allow for nice lower bounds for the Q -modulus (see e.g. [HK], [BHK]). In fact, the Q -regularity condition on balls of Whitney type is not explicitly stated in [H1] but it follows from the other assumptions. The precise definition of a uniform space is given in Section 2 below. This concept, introduced in [BHK], generalizes the notion of a uniform domain introduced by Jones [Jo] and Martio and Sarvas [MS], see also [GO]. The volume growth condition for ρ then refers to integrals of ρ^Q with respect to the measure μ . For predecessors of the results in [H1], see [HN], [HR]. For connections to Gromov hyperbolicity, see [Gr], [BHK] and [BB].

In this paper we show that, suprisingly, lower bounds on the Q -modulus are not needed for the Gehring–Hayman inequality.

Theorem 1.1 (Gehring–Hayman Theorem). *Let $Q > 1$ and let (Ω, d, μ) be a uniform space equipped with a measure that is Q -regular on balls of Whitney type. If $\rho: \Omega \rightarrow (0, \infty)$ is a conformal density on Ω , then*

$$\ell_\rho([x, y]) \leq C\ell_\rho(\gamma)$$

whenever $[x, y]$ is a quasihyperbolic geodesic and γ is a curve joining x to y in Ω , where $C \geq 1$.

The definition of a quasihyperbolic geodesic is given in Section 2. The Gehring–Hayman theorem was a central tool in [BHR], [BKR],[H1] and [H2]. We expect that Theorem 1.1 will allow one to remove the use of modulus bounds in [BHR], [BKR],[H1] and [H2] and thus extend large parts of those papers to a much more

general setting. A very simple example of a space that satisfies the assumptions of Theorem 1.1 but does not support lower bounds for the Q -modulus is

$$\Omega = \{(x, y) \in \mathbb{R}^2 : |y| \leq |x|, -1 < x < 1\}$$

equipped with the path metric and Lebesgue measure.

2. PRELIMINARIES

Let (Ω, d) be a metric space. A *curve* means a continuous map $\gamma: [a, b] \rightarrow \Omega$ from an interval $[a, b] \subset \mathbb{R}$ to Ω . We also denote the image set $\gamma([a, b])$ of γ by γ . The *length* $\ell_d(\gamma)$ of γ with respect to the metric d is defined as

$$\ell_d(\gamma) = \sup \sum_{i=0}^{m-1} d(\gamma(t_i), \gamma(t_{i+1})),$$

where the supremum is taken over all partitions $a = t_0 < t_1 < \dots < t_m = b$ of the interval $[a, b]$. If $\ell_d(\gamma) < \infty$, then γ is said to be a *rectifiable curve*. When the parameter interval is open or half-open, we set

$$\ell_d(\gamma) = \sup \ell_d(\gamma|_{[c, d]}),$$

where supremum is taken over all compact subintervals $[c, d]$. For a rectifiable curve γ we define the *arc length* $s: [a, b] \rightarrow [0, \infty)$ along γ by

$$s(t) = \ell_d(\gamma|_{[a, t]}).$$

Next, let $\rho: \Omega \rightarrow [0, \infty]$ be a Borel function. For each rectifiable curve $\gamma: [a, b] \rightarrow \Omega$ we define the ρ -*length* $\ell_\rho(\gamma)$ of γ by

$$\ell_\rho(\gamma) = \int_\gamma \rho ds = \int_a^b \rho(\gamma(t)) ds(t).$$

If Ω is *rectifiably connected* — that is every pair of points in Ω can be joined by a rectifiable curve — then ρ determines a distance function

$$d_\rho(x, y) = \inf \ell_\rho(\gamma),$$

where the infimum is taken over all rectifiable curves γ joining $x, y \in \Omega$. In general, the distance function d_ρ need not be a metric. However, it is a metric — called a ρ -*metric* — if ρ is positive and continuous. If $\rho \equiv 1$, then $\ell_\rho(\gamma) = \ell_d(\gamma)$ is the length of the curve γ with respect to the metric d . Furthermore if $d(x, y) = \ell_d(\gamma)$ for some curve γ joining points $x, y \in \Omega$, then γ is said to be a *geodesic arc* or just a *geodesic*. If every pair of points in Ω can be joined by a geodesic arc, then Ω is called a *geodesic space*.

Let (Ω, d) be a locally compact, rectifiably connected and noncomplete metric space and denote by $\bar{\Omega}$ its metric completion. Then the *boundary* $\partial\Omega = \bar{\Omega} \setminus \Omega$ is nonempty. We denote

$$d(z) = \text{dist}_d(z, \partial\Omega) = \inf \{d(z, x) : x \in \partial\Omega\}$$

for $z \in \Omega$. If we choose

$$\rho(z) = \frac{1}{d(z)},$$

we obtain the *quasihyperbolic metric* k in Ω . In this special case we denote the metric d_ρ by k and the quasihyperbolic length of the curve γ by $\ell_k(\gamma)$. That $\ell_k(\gamma) = \ell_\rho(\gamma)$ is shown in [BHK, Appendix]. Moreover, $[x, y]$ refers to a quasihyperbolic geodesic joining points x and y in Ω .

Given a real number $D \geq 1$, a curve $\gamma: [a, b] \rightarrow (\Omega, d)$ is called a D -uniform curve if it is *quasiconvex*:

$$(2.1) \quad \ell_d(\gamma) \leq Dd(\gamma(a), \gamma(b)),$$

and

$$(2.2) \quad \min\{\ell_d(\gamma|[a, t]), \ell_d(\gamma|[t, b])\} \leq Dd(\gamma(t))$$

for every $t \in [a, b]$. A metric space (Ω, d) is called a D -uniform space if every pair of points in it can be joined by a D -uniform curve. If (Ω, d) is a uniform space, then by [BHK, Proposition 2.8 and Theorem 2.10] the quasihyperbolic space (Ω, k) is complete, proper (closed balls are compact), and geodesic. Furthermore, each quasihyperbolic geodesic $[x, y]$ is a D' -uniform curve for every $x, y \in \Omega$, where $D' = D'(D) \geq 1$. Quasihyperbolic geodesics are also *locally D' -uniform curves* — that is, every subcurve $[u, v] \subset [x, y]$ is a D' -uniform curve — because $[u, v]$ is a quasihyperbolic geodesic as well. We also have an estimate for a quasihyperbolic distance of every pair of points x and y in the D -uniform space (Ω, d) (see [BHK, Lemma 2.13]):

$$(2.3) \quad k(x, y) \leq 4D^2 \log\left(1 + \frac{d(x, y)}{\min\{d(x), d(y)\}}\right).$$

Let us consider a continuous function $\rho: \Omega \rightarrow (0, \infty)$, called a *density*. The metric d_ρ is then well-defined. We use the subscript ρ for metric notations which refer to d_ρ , and similarly for k and d . For example, $B_\rho(a, r)$, $B_k(a, r)$ and $B_d(a, r)$ are open balls with centre a and radius r in metrics d_ρ , k and d . Furthermore we abbreviate the “Whitney ball” $B_d(z, \frac{1}{2}d(z))$ to B_z .

Let μ be a Borel regular measure on (Ω, d) with dense support. We call ρ a *conformal density* provided it satisfies both a *Harnack type inequality* HI(A) for some constant $A \geq 1$:

$$\text{HI(A)} \quad \frac{1}{A} \leq \frac{\rho(x)}{\rho(y)} \leq A \quad \text{for all } x, y \in B_z \text{ and all } z \in \Omega,$$

and a *volume growth condition* VG(B) for some constant $B > 0$:

$$\text{VG(B)} \quad \mu_\rho(B_\rho(z, r)) \leq Br^Q \quad \text{for all } z \in \Omega \text{ and } r > 0.$$

Here μ_ρ is the Borel measure on Ω defined by

$$\mu_\rho(E) = \int_E \rho^Q d\mu \quad \text{for a Borel set } E \subset \Omega,$$

and Q is a positive real number. Generally Q will be the Hausdorff dimension of our space (Ω, d) .

We defined in the introduction the concept of Q –regularity on balls of Whitney type. The immediate consequence is that the measure μ is also *doubling on balls of Whitney type*: there exists a constant $C_2 \geq 1$ such that

$$(2.4) \quad \mu(B_d(z, 2r)) \leq C_2 \mu(B_d(z, r))$$

for every $z \in \Omega$ and every $0 < r \leq \frac{1}{4} d(z)$.

3. WHITNEY COVERING

In this section we assume that (Ω, d, μ) is a locally compact, rectifiably connected, and non–complete metric measure space such that the measure μ is doubling on balls of Whitney type. Let $r(z) = d(z)/50$. From the family of balls $\{B_d(z, r(z))\}_{z \in \Omega}$ we select a maximal (countable) subfamily $\{B_d(z_i, r(z_i)/5)\}_{i \in I}$ of pairwise disjoint balls. We denote $\mathcal{B} = \{B_i\}_{i \in I}$, where $B_i = B_d(z_i, r_i)$ and $r_i = r(z_i)$. We call the family \mathcal{B} the *Whitney covering* of Ω . Let us list a few facts concerning the Whitney covering. The last property is a consequence of the doubling on balls of Whitney type property of the measure μ . For more properties of the Whitney covering, see e.g. [HKT, Lemma 7].

Lemma 3.1. *There is $N \in \mathbb{N}$ such that*

(i) *the balls $B(z_i, r_i/5)$ are pairwise disjoint,*

(ii) $\Omega = \bigcup_{i \in I} B(z_i, r_i)$,

(iii) $B(z_i, 5r_i) \subset \Omega$,

(iv) $\sum_{i=1}^{\infty} \chi_{B(z_i, 5r_i)}(x) \leq N$ for all $x \in \Omega$.

The family \mathcal{B} has same kind of properties as the usual Whitney decomposition \mathcal{W} of a domain $\Omega \subset \mathbb{R}^n$ and next we prove a couple of them. In addition to the assumptions above, we assume that for each pair of points in $B \in \mathcal{B}$ for every $B \in \mathcal{B}$ can be joined by a D –uniform curve in Ω .

Lemma 3.2. *Let $x, y \in (\Omega, d, \mu)$ and $d(x, y) \geq d(x)/2$. There is a constant $C = C(C_2, D) > 0$ such that*

$$C^{-1}N(x, y) \leq k(x, y) \leq CN(x, y),$$

where $N(x, y)$ is the number of balls $B \in \mathcal{B}$ intersecting the quasihyperbolic geodesic $[x, y]$.

Proof. Let $x, y \in \Omega$ be points such that $d(x, y) \geq d(x)/2$. Because $24 \operatorname{diam}_d(B) \leq d(z)$ for every $B \in \mathcal{B}$ and every $z \in B$, then the basic estimate (2.3) implies

$$\operatorname{diam}_k(B) \leq 4D^2 \log \left(1 + \frac{\operatorname{diam}_d(B)}{24 \operatorname{diam}_d(B)} \right) = 4D^2 \log \frac{25}{24}.$$

Thus

$$N(x, y) \geq \frac{k(x, y)}{4D^2 \log \frac{25}{24}}.$$

Lemma 3.1 (iv) says that there are only N balls $B \in \mathcal{B}$ that contain x . Fix one of them and denote it B_1 . A *neighbour* of the ball B_1 is a ball $B \in \mathcal{B}$ which intersects the ball $5B_1 = B_d(z_1, 5r_1) = B_d(z_1, d(z_1)/10)$. Because the measure μ is doubling in every ball $B_d(z, r)$ with radius $0 < r \leq d(z)/4$, the ball B_1 has a uniformly bounded number of neighbours. Let this number be $N' \in \mathbb{N}$ and let $y_1 \in [x, y]$ be the first point such that y_1 does not belong to any neighbour of B_1 . This choice is possible because $d(x, y) \geq d(x)/2$. The geodesic $[x, y_1]$ intersects at most N' balls $B \in \mathcal{B}$ and

$$\begin{aligned} (3.1) \quad k(x, y_1) &= \int_{[x, y_1]} \frac{1}{d(z)} ds \geq \int_{5B_1 \cap [x, y_1]} \frac{10}{11 d(z_1)} ds \\ &\geq \frac{10}{11 d(z_1)} \left(\frac{d(z_1)}{10} - \frac{d(z_1)}{50} \right) = \frac{4}{55}. \end{aligned}$$

Let $B_2 \in \mathcal{B}$ be a ball such that $y_1 \in B_2$ and $B_2 \cap B \neq \emptyset$ for some neighbour $B \in \mathcal{B}$ of B_1 . Again there are only N' balls $B \in \mathcal{B}$ which are neighbours of B_2 . Let $y_2 \in [x, y]$ be the first point so that y_2 does not belong to any neighbour of B_2 . Then the geodesic $[y_1, y_2]$ intersects at most N' balls $B \in \mathcal{B}$ and $k(y_1, y_2) \geq \frac{4}{55}$, by the same way than in (3.1). We continue this process until we end up with a ball B_m whose neighbours contain $[y_{m-1}, y]$. This process really ends and $m < \infty$, because $[x, y]$ is compact. We may start doing this process from every ball B that contains x . Thus we obtain the upper bound to the number of balls that intersects the quasihyperbolic geodesic $[x, y]$:

$$N(x, y) \leq \frac{55}{4} NN' k(x, y). \quad \square$$

Fix a ball B_0 from the Whitney covering \mathcal{B} and let z_0 be its centre point. For each $B_i \in \mathcal{B}$ we fix a geodesic $[z_0, z_i]$. Furthermore, for each $B_i \in \mathcal{B}$ we set $P(B_i) = \{B \in \mathcal{B} : B \cap [z_0, z_i] \neq \emptyset\}$ and define the *shadow* $S(B)$ of a ball $B \in \mathcal{B}$ by

$$S(B) = \bigcup_{\substack{B_i \in \mathcal{B} \\ B \in P(B_i)}} B_i.$$

For $n \in \mathbb{N}$ we set

$$\mathcal{B}_n = \{B_i \in \mathcal{B} : n \leq k(z_0, z_i) < n + 1\}.$$

The next two lemmas are metric space analogues of [KL, Lemma 2.1 and Lemma 2.2].

Lemma 3.3. *Let γ be a quasihyperbolic geodesic in Ω starting at the point z_0 . Then there is a constant $C = C(C_2, D) > 0$ such that, for each $n \in \mathbb{N}$,*

$$\#\{B \in \mathcal{B}_n : B \cap \gamma \neq \emptyset\} \leq C.$$

Proof. Denote

$$a_n := \#\{B \in \mathcal{B}_n : B \cap \gamma \neq \emptyset\} < \infty.$$

Let $B_1, \dots, B_{a_n} \in \mathcal{B}_n$ be the balls intersecting γ , ordered so that if $k < l$, then there exists $x_k \in B_k \cap \gamma$ such that for every $z \in B_l \cap \gamma$, we have $k(z_0, x_k) \leq k(z_0, z)$. We may assume that $d(x_1, x_{a_n}) \geq d(x_1)/2$, otherwise $x_{a_n} \in B_{x_1}$ and we get the result by doubling on balls of Whitney type. Thus by Lemma 3.2, $k(x_1, x_{a_n}) \geq \frac{a_n}{C}$. Since $k(z_i, x_i) \leq \frac{1}{49} < 1$ for all $i = 1, \dots, a_n$, we calculate

$$\begin{aligned} \frac{a_n}{C} &\leq k(x_1, x_{a_n}) = k(z_0, x_{a_n}) - k(z_0, x_1) \\ &\leq k(z_0, z_{a_n}) + k(z_{a_n}, x_{a_n}) - (k(z_0, z_1) - k(x_1, z_1)) \\ &\leq (n+1) + 1 - n + 1 = 3. \end{aligned}$$

Hence $a_n \leq 3C$. \square

Lemma 3.4. *There is a constant $C = C(C_2, D) > 0$ such that, for each $n \in \mathbb{N}$,*

$$\sum_{B \in \mathcal{B}_n} \chi_{S(B)}(x) \leq C$$

whenever $x \in \Omega$.

Proof. Let $x \in \Omega$. The number of balls $B \in \mathcal{B}$ containing x is bounded, so we may assume that there is a unique ball, denote it B_1 , in \mathcal{B} such that $x \in B_1$. Let $[z_0, z_1]$ be the fixed geodesic joining z_0 to z_1 . Then $x \in S(B)$ for $B \in \mathcal{B}_n$ if and only if $[z_0, z_1] \cap B \neq \emptyset$. By Lemma 3.3, the number of balls $B \in \mathcal{B}_n$ is bounded by a constant that is independent of n . \square

4. GEHRING–HAYMAN THEOREM

We begin with *Frostman's Lemma*. First we recall the definitions of the Hausdorff measure and the weighted Hausdorff measure. Let (X, d) be a compact metric space. Let $0 \leq s < \infty$ and $0 < \delta \leq \infty$. We set

$$\lambda_\delta^s(X) = \inf \left\{ \sum_{i=1}^{\infty} c_i \text{diam}_d(E_i)^s : \chi_X \leq \sum_i c_i \chi_{E_i}, c_i > 0, \text{diam}_d(E_i) \leq \delta \right\}.$$

The *weighted Hausdorff s -measure* of X is

$$\lambda^s(X) = \lim_{\delta \rightarrow 0} \lambda_\delta^s(X).$$

In the special case, where $c_i = 1$ for every $i = 1, 2, \dots$, we denote $\mathcal{H}_\delta^s(X) = \lambda_\delta^s(X)$ and we obtain the *Hausdorff s -measure*

$$\mathcal{H}^s(X) = \lim_{\delta \rightarrow 0} \mathcal{H}_\delta^s(X).$$

The Hausdorff s -content of X is

$$\mathcal{H}_\infty^s(X) = \inf \left\{ \sum_{i=1}^{\infty} \text{diam}_d(E_i)^s : X \subset \bigcup_{i=1}^{\infty} E_i \right\}.$$

By [Ma, Lemma 8.16] we know that $\mathcal{H}^s(X) \leq 30^s \lambda^s(X)$, but in fact from the proof of that lemma one obtains that

$$\mathcal{H}_{30\delta}^s(X) \leq 30^s \lambda_\delta^s(X) \quad \text{for every } 0 < \delta \leq \infty.$$

In particular

$$\mathcal{H}_\infty^s(X) \leq 30^s \lambda_\infty^s(X).$$

The following formulation of Frostman's Lemma (cf. [Ma, Theorem 8.17.]) is suitable for our purposes.

Theorem 4.1 (Frostman's Lemma). *For any $s \geq 0$ there is a Radon measure ω on X such that*

$$\omega(X) = \lambda_\infty^s(X)$$

and

$$\omega(E) \leq \text{diam}_d(E)^s \quad \text{for all } E \subset X.$$

In particular, when $s = 1$ and X is connected, we obtain

$$\omega(X) \geq \frac{1}{30} \mathcal{H}_\infty^1(X) \geq \frac{\text{diam}_d(X)}{60}.$$

For the rest of the paper we assume that (Ω, d, μ) is a locally compact, non-complete and D -uniform metric measure space such that the measure μ is Q -regular on balls of Whitney type for some $Q > 1$. Let ρ be a conformal density such that the number Q in the definition $\text{VG}(B)$ coincides with the previous $Q > 1$.

Proof of Theorem 1.1. Let x and y be points in $\bar{\Omega}$ and let $[x, y]$ be a quasihyperbolic geodesic in Ω joining points x and y . Because quasihyperbolic geodesics are D' -uniform curves, $[x, y]$ is rectifiable in the metric d . Let γ be another rectifiable curve in Ω joining points x and y . Let $a \in [x, y]$ be the point such that $\ell_d([x, a]) = \ell_d([a, y])$ and write $p = d(x, a)$. Moreover, for each $j \in \mathbb{N}$ write $A_j = (\bar{B}_d(x, 2^{-j}p) \setminus B_d(x, 2^{-j-1}p)) \cap \Omega$. Let $[x_{j+1}, x_j] \subset [x, y]$ be a subcurve, where x_{j+1} is the last point of $[x, y]$ in $\bar{B}(x, 2^{-j-1}p)$ and x_j is the last point of $[x, y]$ in $\bar{B}(x, 2^{-j}p)$, and set $\gamma_j = \gamma \cap A_j$. We may clearly assume that γ_j is connected. By summing and symmetry it suffices to prove that

$$(4.1) \quad \ell_\rho([x_{j+1}, x_j]) \leq C \ell_\rho(\gamma_j)$$

for every $j \in \mathbb{N}$.

Let $j \in \mathbb{N}$. From the definition of the curve γ_j it follows that

$$(4.2) \quad \ell_d(\gamma_j) \geq 2^{-j-1}p.$$

From the definition of the quasihyperbolic geodesic $[x_{j+1}, x_j]$ and from the local D' -uniformity of the curve $[x, y]$, we have that

$$(4.3) \quad \ell_d([x_{j+1}, x_j]) \leq D' d(x_{j+1}, x_j) \leq D' 2^{-j+1} p,$$

$$(4.4) \quad 2^{-j-1} p \leq \ell_d([x, z]) \leq D' d(z) \quad \text{for every } z \in [x_{j+1}, x_j]$$

and

$$(4.5) \quad \begin{aligned} k(x_{j+1}, x_j) &= \ell_k([x_{j+1}, x_j]) = \int_{[x_{j+1}, x_j]} \frac{1}{d(z)} ds \\ &\leq \frac{D'}{p} 2^{j+1} \ell_d([x_{j+1}, x_j]) \leq 4D'^2. \end{aligned}$$

We first prove that inequality (4.1) holds when the curves $[x_{j+1}, x_j]$ and γ_j are “close” to each other in the quasihyperbolic metric. Let

$$M > \max \left\{ 4D'^2 \frac{\log(4D'^2)}{\log 2} + 1, 4D'^2 \frac{\log(B(2 + A^2/6)^Q / c_7)}{\log 2} \right\},$$

where the constant $c_7 > 0$ is a sufficiently small constant depending on A, C_1, D and Q , and let us assume that $\text{dist}_k([x_{j+1}, x_j], \gamma_j) \leq M$. Let $y_j \in [x_{j+1}, x_j]$ and $\tilde{y}_j \in \gamma_j$ be points such that $k(y_j, \tilde{y}_j) \leq M$. Let us show that we may estimate the ρ -length of the quasihyperbolic geodesic $[x_{j+1}, x_j]$ from above by $2^{-j} p \rho(y_j)$ in the following way

$$(4.6) \quad \ell_\rho([x_{j+1}, x_j]) \leq A^b D' \rho(y_j) 2^{-j+1} p,$$

where $b = 4c_1 D'^2$ and $c_1 = c_1(C_1) > 0$ is the constant from Lemma 3.2.

If there exists $z \in [x_{j+1}, x_j]$ such that $[x_{j+1}, x_j] \subset B_z = B_d(z, d(z)/2)$, we obtain from HI(A) and (4.3)

$$\ell_\rho([x_{j+1}, x_j]) \leq A \rho(y_j) \ell_d([x_{j+1}, x_j]) \leq A D' \rho(y_j) 2^{-j+1} p.$$

Otherwise we may assume that $d(x_{j+1}, x_j) \geq d(x_{j+1})/2$. From Lemma 3.2 and inequality (4.5), it follows that

$$N([x_{j+1}, x_j]) \leq 4c_1 D'^2 = b,$$

where the constant $c_1 = c_1(C_1) > 0$ is the constant from Lemma 3.2. Then by HI(A) every $z \in [x_{j+1}, x_j]$ satisfies

$$\rho(z) \leq A^b \rho(y_j).$$

This with (4.3) gives us inequality (4.6)

$$\begin{aligned} \ell_\rho([x_{j+1}, x_j]) &\leq A^b \rho(y_j) \ell_d([x_{j+1}, x_j]) \\ &\leq A^b D' \rho(y_j) 2^{-j+1} p. \end{aligned}$$

Next we estimate the ρ -length of the curve γ_j from below by $2^{-j}p\rho(y_j)$. If $[x_{j+1}, x_j] \cap B_{\tilde{y}_j} \neq \emptyset$, we easily get from HI(A) an estimate for $\ell_\rho(\gamma_j)$:

$$(4.7) \quad \ell_\rho(\gamma_j) \geq \frac{1}{A^{b+1}} \rho(y_j) \ell_d(\gamma_j \cap B_{\tilde{y}_j}).$$

Furthermore, for every $z \in [x_{j+1}, x_j] \cap B_{\tilde{y}_j}$, using inequalities (4.2) and (4.4) it holds that

$$(4.8) \quad \ell_d(\gamma_j \cap B_{\tilde{y}_j}) \geq \begin{cases} 2^{-j-1}p, & \text{if } \gamma_j \subset B_{\tilde{y}_j} \\ \frac{1}{2}d(\tilde{y}_j) \geq \frac{1}{2}\left(\frac{2}{3}d(z)\right) \geq \frac{1}{3D'}2^{-j-1}p, & \text{if } \gamma_j \not\subset B_{\tilde{y}_j}. \end{cases}$$

In this case, combining (4.6), (4.7) and (4.8) we obtain the desired result (4.1)

$$\ell_\rho([x_{j+1}, x_j]) \leq 12A^{2b+1}D'^2\ell_\rho(\gamma_j).$$

Therefore we may assume that $[x_{j+1}, x_j] \cap B_{\tilde{y}_j} = \emptyset$. This implies that $d(y_j, \tilde{y}_j) \geq d(\tilde{y}_j)/2$. By Lemma 3.2 there are at most $h := Mc_1$ balls from the Whitney covering \mathcal{B} that intersect $[y_j, \tilde{y}_j]$ and hence, by HI(A),

$$(4.9) \quad \rho(y_j) \leq A^h \rho(\tilde{y}_j).$$

On the other hand by HI(A) and (4.2)

$$(4.10) \quad \ell_\rho(\gamma_j) \geq \frac{1}{A} \rho(\tilde{y}_j) \ell_d(\gamma_j \cap B_{\tilde{y}_j}) \geq \begin{cases} \frac{1}{A} \rho(\tilde{y}_j) 2^{-j-1}p, & \text{if } \gamma_j \subset B_{\tilde{y}_j} \\ \frac{1}{2A} \rho(\tilde{y}_j) d(\tilde{y}_j), & \text{if } \gamma_j \not\subset B_{\tilde{y}_j}. \end{cases}$$

If $\gamma_j \subset B_{\tilde{y}_j}$, again we obtain the desired inequality (4.1) by combining inequalities (4.6), (4.9) and (4.10). If $\gamma_j \not\subset B_{\tilde{y}_j}$, then (4.10) with (4.9) gives

$$(4.11) \quad \rho(y_j) \leq A^{h+1} \frac{2}{d(\tilde{y}_j)} \ell_\rho(\gamma_j).$$

By elementary inequalities in [GP, Lemma 2.1] and [BHK, Inequality (2.4)] we obtain

$$\log\left(1 + \frac{d(y_j, \tilde{y}_j)}{\min\{d(y_j), d(\tilde{y}_j)\}}\right) \leq k(y_j, \tilde{y}_j) \leq M$$

and further

$$(4.12) \quad \frac{1}{d(\tilde{y}_j)} \leq \frac{e^M - 1}{d(y_j, \tilde{y}_j)}.$$

Moreover, the assumption $d(y_j, \tilde{y}_j) \geq d(\tilde{y}_j)/2$ gives us

$$d(y_j) \leq d(y_j, \tilde{y}_j) + d(\tilde{y}_j) \leq 3d(y_j, \tilde{y}_j).$$

This, along with inequalities (4.11), (4.12) and (4.4), yields an estimate for the ρ –length of γ_j :

$$\begin{aligned}
 (4.13) \quad \rho(y_j) &\leq 2A^{h+1} \frac{e^M - 1}{d(y_j, \tilde{y}_j)} \ell_\rho(\gamma_j) \leq 6A^{h+1} \frac{e^M - 1}{d(y_j)} \ell_\rho(\gamma_j) \\
 &\leq 6A^{h+1} (e^M - 1) \frac{D'}{p} 2^{j+1} \ell_\rho(\gamma_j).
 \end{aligned}$$

Now combining (4.6) and (4.13) we obtain

$$\ell_\rho([x_{j+1}, x_j]) \leq 24(e^M - 1)A^{b+h+1}D'^2 \ell_\rho(\gamma_j).$$

Thus (4.1) is proven when the curves $[x_{j+1}, x_j]$ and γ_j are “close” to each other in the quasihyperbolic metric. Therefore we may assume that $\text{dist}_k([x_{j+1}, x_j], \gamma_j) > M$. Let $w_j \in [x_{j+1}, x_j]$ satisfy $d(x, w_j) = 3 \cdot 2^{-j-2}p$. Denote $\ell_\rho(\gamma_j) = r$ and let $w \in \gamma_j$. Let us consider the ρ –ball $B_\rho(w, 2r)$. If $\text{dist}_k(w_j, B_\rho(w, 2r)) < M$, there exists $u \in B_\rho(w, 2r)$ such that $k(w_j, u) \leq M$ and hence $\rho(w_j) \leq A^h \rho(u)$. We may assume that $\gamma_j \cap \frac{1}{2}B_u = \emptyset$. Otherwise $\text{dist}_k([x_{j+1}, x_j], \gamma_j) \leq M + 1$ and replacing M with $M + 1$ we obtain the result by the previous case. As we have assumed $\gamma_j \cap \frac{1}{2}B_u = \emptyset$,

$$\begin{aligned}
 2\ell_\rho(\gamma_j) = 2r > \text{dist}_\rho(u, \gamma_j) &\stackrel{\text{HI(A)}}{\geq} \frac{1}{4A} \rho(u) d(u) \\
 &\stackrel{(4.9)}{\geq} \frac{1}{4A^{h+1}} \rho(w_j) d(u) \stackrel{(*)}{\geq} \frac{1}{4A^{h+1}e^M} \rho(w_j) d(w_j) \\
 &\stackrel{(4.4)}{\geq} \frac{2^{-j-1}p}{4A^{h+1}D'e^M} \rho(w_j) \\
 &\stackrel{(4.6)}{\geq} \frac{1}{16A^{b+h+1}D'^2e^M} \ell_\rho([x_{j+1}, x_j]).
 \end{aligned}$$

The inequality (*) above follows from the elementary estimate ([GP, Lemma 2.1], [BHK, Inequality (2.3)])

$$\left| \log \frac{d(w_j)}{d(u)} \right| \leq k(w_j, u) \leq M.$$

Again we find a constant $C > 1$ such that $\ell_\rho([x_{j+1}, x_j]) \leq C\ell_\rho(\gamma_j)$. So (4.1) is satisfied.

Hence we may assume that the ρ –ball $B_\rho(w, 2r)$ is “far away” from the quasihyperbolic geodesic $[x_{j+1}, x_j]$. More precisely, we may assume that $\text{dist}_k(w_j, B_\rho(w, 2r)) \geq M$. Our plan is to prove that the volume growth condition $\text{VG}(\mathbb{B})$ does not hold for such a ρ –ball.

Let for every $z \in \gamma_j$, $[z, w_j]$ be a quasihyperbolic geodesic which joins z and w_j . Cover $[z, w_j]$ with balls $\{B_1, \dots, B_{n(z)}\} \subset \mathbb{B}$ ordered so that if $m < n$, then there exists $z_m \in B_m \cap [z, w_j]$ such that for every $\tilde{z} \in B_n \cap [z, w_j]$, we have

$k(z, z_m) \leq k(z, \tilde{z})$. Recall that $n(z) < \infty$. Denote $[z, w_z] \subset [z, w_j]$, where w_z is the first point which does not belong to $B_\rho(w, 2r)$. Thus $\ell_\rho([z, w_z]) \geq r$. Let $\{B_1, \dots, B_{n_r(z)}\} \subset \{B_1, \dots, B_{n(z)}\}$ be those balls which cover $[z, w_z]$. So by HI(A) and by the local D' -uniformity (quasiconvexity) of quasihyperbolic geodesics we obtain

$$(4.14) \quad \begin{aligned} r \leq \ell_\rho([z, w_z]) &\leq \sum_{i=1}^{n_r(z)} A\rho(z_i)\ell_d([z, w_z] \cap B_i) \\ &\leq AD' \sum_{i=1}^{n_r(z)} \rho(z_i) \operatorname{diam}_d(B_i). \end{aligned}$$

We next provide a tool that will be used to estimate the μ_ρ -measure of the ρ -ball $B_\rho(w, 2r)$. We claim that if $B \in \mathcal{B}$ intersects $B_\rho(w, 2r)$, then $B \subset B_\rho(w, (2 + \frac{A^2}{6})r)$. To show this, it suffices to prove that if $B \in \mathcal{B}$ intersects $B_\rho(w, 2r)$ then

$$(4.15) \quad \operatorname{diam}_\rho(B) \leq \frac{A^2}{6}r.$$

Consider such a ball $B \in \mathcal{B}$. It follows from HI(A) that

$$\operatorname{diam}_\rho(B) \leq A\rho(z_B) \operatorname{diam}_d(B) = \frac{A}{25}\rho(z_B) d(z_B)$$

for each $B \in \mathcal{B}$, where z_B is the centre of B . Hence it actually suffices to prove that

$$(4.16) \quad \rho(z_B) d(z_B) \leq 4Ar.$$

Let $y \in B \cap B_\rho(w, 2r)$. If $w \notin B_{z_B}$, then there exists a curve γ , which joins points w and y and

$$\begin{aligned} 2r &\geq \int_\gamma \rho(z) ds \geq \frac{1}{A}\rho(z_B)\ell_d(\gamma \cap B_{z_B}) \\ &\geq \left(\frac{1}{2} - \frac{1}{50}\right) \frac{1}{A}\rho(z_B) d(z_B) \geq \frac{24}{25A}\rho(z_B) d(z_B) \end{aligned}$$

and the inequality (4.16) is proven. Let us assume that $w \in B_{z_B}$. The elementary estimate (2.3) implies

$$M \leq k(w_j, w) \leq 4D^2 \log\left(1 + \frac{d(w_j, w)}{\min\{d(w_j), d(w)\}}\right).$$

Along with the assumption that $M > 4D^2 \frac{\log(4D'^2)}{\log 2} + 1$, we also see that

$$(4.17) \quad \min\{d(w_j), d(w)\} \leq \frac{d(w_j, w)}{e^{M/4D^2} - 1} \leq 2^{-j+1-(M-1)/4D^2} p$$

The assumption $M > 4D^2 \frac{\log(4D'^2)}{\log 2} + 1$ and (4.4) give us

$$(4.18) \quad \begin{aligned} d(w_j) &\geq \frac{p}{D'} 2^{-j-1} = 2^{-j+1-(M-1)/4D^2} p \frac{2^{(M-1)/4D^2}}{2^2 D'} \\ &\geq 2^{-j+1-(M-1)/4D^2} p. \end{aligned}$$

Thus it follows from inequality (4.17) that

$$d(w) \leq 2^{-j+1-(M-1)/4D^2} p \leq 2^{-j-1} p.$$

Hence, from the definition of the curve γ_j and inequality (4.2) we know that γ_j is can not be a subset of B_w . Then by HI(A)

$$r = \int_{\gamma_j} \rho(z) ds \geq \frac{1}{2A} \rho(z_B) d(w) \geq \frac{1}{4A} \rho(z_B) d(z_B),$$

and (4.16) is proven.

Now we know that if $B \in \mathcal{B}$ intersects $B_\rho(w, 2r)$, then $B \subset B_\rho(w, (2 + \frac{A^2}{6})r)$. Then by HI(A), Lemma 3.1 (iv) and Q -regularity on balls of Whitney type, we have

$$(4.19) \quad \begin{aligned} \mu_\rho(B_\rho(w, (2 + \frac{A^2}{6})r)) &= \int_{B_\rho(w, (2 + \frac{A^2}{6})r)} \rho^Q d\mu \geq \sum_{\substack{B \in \mathcal{B} \\ B \cap B_\rho(w, 2r) \neq \emptyset}} \frac{1}{NA^Q} \rho(z_B)^Q \mu(B) \\ &\geq \sum_{\substack{B \in \mathcal{B} \\ B \cap B_\rho(w, 2r) \neq \emptyset}} c_2 \rho(z_B)^Q \left(\frac{\text{diam}_d(B)}{2} \right)^Q, \end{aligned}$$

where $c_2 = \frac{1}{NC_1 A^Q}$.

Let us choose the basepoint z_0 to be w_j . According to Frostman's Lemma (Theorem 4.1) there is a Radon measure ω supported on γ_j such that $\omega(\gamma_j) \geq \frac{\text{diam}_d(\gamma_j)}{60}$ and $\omega(E) \leq \text{diam}_d(E)$ for every $E \subset \Omega$. Then with (4.14) we obtain (a version of Fubini's theorem)

$$(4.20) \quad \begin{aligned} \omega(\gamma_j) r &\leq AD' \int_{\gamma_j} \sum_{i=1}^{n_r(z)} \rho(z_i) \text{diam}_d(B_i) d\omega(z) \\ &\leq AD' \sum_{n=M-1}^{\infty} \sum_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \rho(z_B) \text{diam}_d(B) \omega(S(B)). \end{aligned}$$

By Hölder inequality this is less or equal to

$$AD' \left(\sum_{n=M-1}^{\infty} \sum_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \rho(z_B)^Q \text{diam}_d(B)^Q \right)^{\frac{1}{Q}} \left(\sum_{n=M-1}^{\infty} \sum_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \omega(S(B))^{\frac{Q}{Q-1}} \right)^{\frac{Q-1}{Q}}.$$

Using (4.19) and the assumption $\text{dist}_k(w_j, B_\rho(w, 2r)) \geq M$ we obtain the estimate

$$\begin{aligned} \omega(\gamma_j)r &\leq AD' \left(\frac{2^Q}{c_2} \mu_\rho(B_\rho(w, (2 + \frac{A^2}{6})r)) \right)^{\frac{1}{Q}} \left(\sum_{n=M-1}^{\infty} \sum_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \omega(S(B))^{\frac{Q}{Q-1}} \right)^{\frac{Q-1}{Q}} \\ (4.21) \quad &= c_3 \left(\mu_\rho(B_\rho(w, (2 + \frac{A^2}{6})r)) \right)^{\frac{1}{Q}} \left(\sum_{n=M-1}^{\infty} \sum_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \omega(S(B))^{\frac{Q}{Q-1}} \right)^{\frac{Q-1}{Q}}, \end{aligned}$$

where $c_3 = 2AD'c_2^{-\frac{1}{Q}} = 2(NC_1)^{\frac{1}{Q}}A^2D'$.

In order to estimate the measure of the shadow of the ball $B \in \mathcal{B}_n$, let us make a couple of preliminary estimates. For every $v \in B \cap [z, w_j]$, where $B \in \mathcal{B}$ and $z \in \gamma_j$, we have by uniformity (quasiconvexity) and inequality (4.3) that

$$d(w_j, v) \leq \ell_d([w_j, v]) \leq \ell_d([w_j, z]) \leq D'd(w_j, z) \leq 2^{-j+1}pD'.$$

In the same way as in inequalities (4.17) and (4.18), we obtain from inequality (4.4) and the assumption $n \geq M-1 \geq 4D^2 \frac{\log(4D'^2)}{\log 2}$ that for every $v \in B \cap [z, w_j]$, where $B \in \mathcal{B}_n$ and $z \in \gamma_j$, it holds that

$$(4.22) \quad d(v) \leq 2^{-j+1-n/4D^2}pD'.$$

Furthermore, for every centre point $z_B \in B \in \mathcal{B}_n$, such that $B \cap [z, w_j] \neq \emptyset$ for some $z \in \gamma_j$, it holds that

$$(4.23) \quad d(z_B) \leq \frac{50}{49} d(v) \leq 2^{-j+1-n/4D^2}p \frac{50D'}{49}.$$

Also from the uniformity of the space (Ω, d) and inequality (4.23) it follows that there exist a constant $c_4 = c_4(C_1, D) \geq 1$ such that for every $B \in \mathcal{B}_n$, so that $B \cap [z, w_j] \neq \emptyset$ for some $z \in \gamma_j$, holds

$$(4.24) \quad \text{diam}_d(S(B)) \leq c_4 \text{diam}_d(B) \leq 2^{-j+2-n/4D^2}pc_4 \frac{50D'}{49}.$$

Now we can calculate by Lemma 3.4, Frostman's Lemma and inequality (4.24) that

$$\begin{aligned}
 \sum_{n=M-1}^{\infty} \sum_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \omega(S(B))^{\frac{Q}{Q-1}} &\leq \sum_{n=M-1}^{\infty} \max_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \omega(S(B))^{\frac{1}{Q-1}} \sum_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \omega(S(B)) \\
 &\leq c_5 \omega(\gamma_j) \sum_{n=M-1}^{\infty} \max_{\substack{B \in \mathcal{B}_n \\ B \cap [z, w_j] \neq \emptyset \\ z \in \gamma_j}} \omega(S(B))^{\frac{1}{Q-1}} \\
 &\leq c_5 \left(\frac{200D'c_4}{49} \right)^{\frac{1}{Q-1}} \omega(\gamma_j) \sum_{n=M-1}^{\infty} (2^{-j-n/4D^2} p)^{\frac{1}{Q-1}} \\
 &\leq c_5 \left(\frac{200D'c_4}{49} \right)^{\frac{1}{Q-1}} \omega(\gamma_j) p^{\frac{1}{Q-1}} 2^{\frac{-j}{Q-1}} \frac{2^{\frac{-M+2}{4D^2(Q-1)}}}{2^{\frac{1}{4D^2(Q-1)}} - 1},
 \end{aligned}$$

where $c_5 = c_5(C_1)$ is from Lemma 3.4. Thus with (4.21) we have

$$\begin{aligned}
 \omega(\gamma_j)^Q r^Q &\leq (c_3)^Q \mu_\rho(B_\rho(w, (2 + \frac{A^2}{6})r)) \\
 &\quad \left(c_5 \left(\frac{200D'c_4}{49} \right)^{\frac{1}{Q-1}} \omega(\gamma_j) p^{\frac{1}{Q-1}} 2^{\frac{-j}{Q-1}} \frac{2^{\frac{-M+2}{4D^2(Q-1)}}}{2^{\frac{1}{4D^2(Q-1)}} - 1} \right)^{Q-1} \\
 &= c_6 \mu_\rho(B_\rho(w, (2 + \frac{A^2}{6})r)) \omega(\gamma_j)^{Q-1} 2^{-j - \frac{M-2}{4D^2}} p,
 \end{aligned}$$

where $c_6 = \frac{200}{49} c_4 N C_1 (2A^2)^Q D^{Q+1} (c_5)^{Q-1} (2^{\frac{1}{4D^2(Q-1)}} - 1)^{1-Q}$. Furthermore $\omega(\gamma_j) \geq \frac{\text{diam}_d(\gamma_j)}{60}$ and this gives us

$$\begin{aligned}
 \mu_\rho(B_\rho(w, (2 + \frac{A^2}{6})r)) &\geq \omega(\gamma_j) \frac{1}{c_6} \frac{2^{j + \frac{M-2}{4D^2}}}{p} r^Q \\
 &\geq \frac{2^{-j-1} p}{60} \frac{1}{c_6} \frac{2^{j + \frac{M-2}{4D^2}}}{p} r^Q \\
 &= 2^{\frac{M}{4D^2}} c_7 r^Q,
 \end{aligned}$$

where $c_7 = \frac{49 \cdot 2^{\frac{-2}{4D^2} - 1} \left(2^{\frac{1}{4D^2(Q-1)}} - 1 \right)^{Q-1}}{12000 c_4 N C_1 (2A^2)^Q D^{Q+1} c_5^{Q-1}}$. This is a contradiction because when M is sufficiently big, the volume growth condition VG(B) will not hold. Consequently, if $k([x_{j+1}, x_j], \gamma_j) > M$ then our ρ -ball is in the quasihyperbolic metric k so big that $\text{dist}_k(w_j, B_\rho(w, 2r)) \leq M$. Thus the conclusion is that $\ell_\rho([x_{j+1}, x_j]) \leq C \ell_\rho(\gamma_j)$ for some $C = C(A, B, C_1, D, Q)$. \square

There is nothing special about the constant $\frac{1}{2}$ in condition HI(A) and the constants $\frac{1}{50}$ and 5 in Whitney covering. The only restriction in the Whitney covering is that if $\lambda_1 B_d(z_1, d(z_1)/\lambda_2) \cap \lambda_1 B_d(z_2, d(z_2)/\lambda_2) \neq \emptyset$, then $\lambda_1 B_d(z_1, d(z_1)/\lambda_2)$ must be included in some ball $B_d(z_2, d(z_2)/\lambda_3)$ on which the measure μ is doubling. Otherwise one can choose the constants as desired.

Acknowledgement. The results of this paper were obtained while the authors were visiting at Centre de Recerca Matemàtica, Universitat Autònoma de Barcelona in April 2009, during the research programme “Harmonic Analysis, Geometric Measure Theory and Quasiconformal Mappings”. They thanks CRM for its hospitality. The authors also thank Kevin Wildrick for carefully reading the manuscript.

REFERENCES

- [BB] Z.M. BALOGH AND S.M. BUCKLEY, *Geometric characterization of Gromov hyperbolicity*. Invent. Math. **153** (2003), 261–301.
- [BO] J. BJÖRN AND J. ONNINEN, *Orlicz capacities and Hausdorff measures on metric spaces*. Math. Z. **251** (2005), 131–146.
- [BHK] M. BONK, J. HEINONEN AND P. KOSKELA, *Uniformizing Gromov Hyperbolic spaces*. Astérisque **270** (2001), 1–99.
- [BHR] M. BONK, J. HEINONEN AND S. ROHDE, *Doubling conformal densities*. J. reine angew. Math. **541** (2001), 117–141.
- [BKR] M. BONK, P. KOSKELA AND S. ROHDE, *Conformal metrics on the unit ball in Euclidean space*. Proc. London Math. Soc. (3) **77** (1998), 635–664.
- [GH] F.W. GEHRING AND W.K. HAYMAN, *An inequality in the theory of conformal mapping*. J. Math. Pures Appl. (9) **41** 1962, 353–361.
- [GO] F.W. GEHRING AND B.G. OSGOOD, *Uniform domains and the quasi-hyperbolic metric*. J. Anal. Math. **36** (1979), 50–74.
- [GP] F.W. GEHRING AND B.P. PALKA, *Quasiconformally homogeneous domains*. J. Anal. Math. **30** (1976), 172–199.
- [Gr] M. GROMOV, *Hyperbolic Groups, Essays in Group Theory*. S. Gersten, Editor, MSRI Publication, Springer, 1987, 75–265.
- [HKT] P. HAJLÁSZ, P. KOSKELA AND H. TUOMINEN, *Measure density and extendability of Sobolev functions*. Rev. Mat. Iberoamericana **24(2)** (2008), 645–669.
- [HK] J. HEINONEN AND P. KOSKELA, *Quasiconformal maps in metric space with controlled geometry*. Acta Math. **181** (1998), 1–61.
- [HN] J. HEINONEN AND R. NÄKKI, *Quasiconformal distortion on arcs*. J. Anal. Math. **63** (1994), 19–53.
- [HR] J. HEINONEN AND S. ROHDE, *The Gehring–Hayman inequality for quasihyperbolic geodesics*. Math. Proc. Camp. Phil. Soc. **114** (1993), 393–404.
- [H1] D.A. HERRON, *Conformal deformations of uniform Loewner spaces*. Math. Proc. Camp. Phil. Soc. **136** (2004), 325–360.
- [H2] D.A. HERRON, *Quasiconformal deformations and volume growth*. Proc. London Math. Soc. (3) **92** (2006), 161–199.
- [Jo] P.W. JONES, *Extension Theorems for BMO*. Indiana Univ. Math. J. **29(1)** (1980), 41–66.
- [KL] P. KOSKELA AND J. LEHRBÄCK, *Quasihyperbolic Growth Conditions and Compact embeddings of Sobolev Spaces*. Michigan Math. J. **55** (2007), 183–193.

- [MS] O. MARTIO AND J. SARVAS, *Injectivity theorems in plane and space*. Ann. Acad. Sci. Fenn. Ser. A. I. Math. **4** (1979), 383–401.
- [Ma] P. MATTILA, *Geometry of Sets and Measures in Euclidean Spaces: Fractals and rectifiability*. Cambridge Univ. Press, Cambridge, 1985.

PEKKA KOSKELA AND PÄIVI LAMMI
DEPARTMENT OF MATHEMATICS AND STATISTICS
P.O.Box 35 (MAD)
FI-40014 UNIVERSITY OF JYVÄSKYLÄ
FINLAND

E-mail address: pekka.j.koskela@jyu.fi

E-mail address: paivi.e.lammi@jyu.fi