

NONPERSISTENCE OF RESONANT CAUSTICS IN PERTURBED ELLIPTIC BILLIARDS

SÔNIA PINTO-DE-CARVALHO AND RAFAEL RAMÍREZ-ROS

ABSTRACT. Caustics are curves with the property that a billiard trajectory, once tangent to it, stays tangent after every reflection at the boundary of the billiard table. When the billiard table is an ellipse, any nonsingular billiard trajectory has a caustic, which can be either a confocal ellipse or a confocal hyperbola. Resonant caustics —the ones whose tangent trajectories are closed polygons— are destroyed under generic perturbations of the billiard table. We prove that none of the resonant elliptical caustics persists under a large class of explicit perturbations of the original ellipse. This result follows from a standard Melnikov argument and the analysis of the complex singularities of certain elliptic functions.

1. INTRODUCTION AND MAIN RESULT

Birkhoff [3] introduced the problem of *convex billiard tables* more than 80 years ago as a way to describe the motion of a free particle inside a closed convex smooth curve. The particle is reflected at the boundary according to the law “angle of incidence equals angle of reflection”. Good modern starting points in the literature of the billiard problem are [11, 18].

Caustics —curves with the property that a billiard trajectory, once tangent to it, stays tangent after every reflection— are the most distinctive geometric objects inside billiard tables, since they are a geometric manifestation of the regularity of their tangent trajectories. For example, integrable billiards have a continuum of caustics, whereas the nonexistence of caustics inside a convex billiard table implies that there are some billiard trajectories whose past and future behaviours differ dramatically. See, for instance, [13]. Hence, the existence and persistence of caustics are two fundamental questions in billiards. Most of the literature deals with convex caustics, since they are easier to understand and related to ordered trajectories. Two exceptions are [8, §3] and [10].

We summarize the classical existence results as follows. On the one hand, if the boundary curve is smooth enough and strictly convex, then there exists a collection of smooth convex caustics close to the boundary of the table whose union has positive area [7, 12]. On the other hand, Mather [13] proved that there are no smooth convex caustics inside a convex billiard table when its boundary

Key words and phrases. Billiards, Caustics, Invariant curves, Melnikov method.

curve has some flat point. Gutkin and Katok [8] gave a quantitative version of Mather's theorem.

The robustness of a smooth convex caustic is closely related to the arithmetic properties of its *rotation number*, which measures the number of turns around the caustic per impact. Caustics with Diophantine rotation numbers persist under small perturbations of the boundary curve. This follows from standard KAM arguments [7, 12]. On the contrary, *resonant caustics* —the ones whose tangent trajectories are closed polygons, so that their rotation numbers are rational— are fragile structures that generically break up. See, for instance, [16].

This raises two complementary questions. First, to characterize the perturbations that preserve/destroy a given resonant caustic of a billiard table. Second, to determine all resonant caustics that are preserved/destroyed under a given perturbation of an integrable billiard table. These questions have been studied by several authors. Baryshnikov and Zarnitsky [2] proved that the perturbations preserving a given resonant caustic of a smooth convex billiard table form an infinite-dimensional Hilbert manifold. As a sample, we point out that this Hilbert manifold is given by the set of billiard tables with constant width when the rotation number of the unperturbed caustic is one half [10]. Concerning the second question, Ramírez-Ros [16] gave a sufficient condition for the break-up of the resonant circular caustics inside a circular billiard table, in terms of the Fourier coefficients of the perturbation, see Remark 3 below.

In this paper we tackle the second question when the billiard boundary is an ellipse. In that case, the billiard dynamics is integrable and any billiard trajectory has a caustic [18]. The caustics are the conics confocal to the original ellipse: confocal ellipses, confocal hyperbolas, and the foci. Poncelet [15] showed that if a billiard trajectory inside an ellipse is a closed polygon, then all the billiard trajectories sharing its caustic are also closed polygons. Even more, if a billiard trajectory tangent to one of the elliptical caustics is a (m, n) -gon —a closed polygon with n sides that makes m turns around its caustic—, then all the billiard trajectories sharing its caustic are also (m, n) -gons, and their caustic is called (m, n) -*resonant*. (These two definitions are not restricted to billiards inside ellipses.) We shall see in Section 4 that there is a unique (m, n) -resonant elliptical caustic for any relatively prime integers m and n such that $1 \leq m < n/2$. Our main result is that all these resonant elliptical caustics break up under a large class of explicit perturbations of the original ellipse, see Theorem 1.

The following notations are required to state the main result. Once fixed the ellipse

$$Q = \left\{ (x, y) \in \mathbb{R}^2 : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \right\}, \quad a > b > 0,$$

we consider its associated elliptic coordinates (μ, φ) given by the relations

$$x = c \cosh \mu \cos \varphi, \quad y = c \sinh \mu \sin \varphi,$$

where $c = \sqrt{a^2 - b^2}$ is the semifocal distance of Q . The equation of the ellipse Q in this elliptic coordinates is $\mu \equiv \mu_0$, where $\cosh \mu_0 = a/c$ and $\sinh \mu_0 = b/c$. Hence, any smooth perturbation Q_ϵ of the ellipse Q can be written in elliptic coordinates as

$$(1) \quad \mu = \mu_\epsilon(\varphi) = \mu_0 + \epsilon \mu_1(\varphi) + O(\epsilon^2),$$

for some 2π -periodic smooth function $\mu_\epsilon(\varphi)$.

Theorem 1. *Let $\mu_1(\varphi)$ be a 2π -periodic entire function. If $\mu_1(\varphi)$ is not constant (respectively, $\mu_1'(\varphi)$ is not π -antiperiodic), then none of the (m, n) -resonant elliptical caustics with odd n (respectively, even n) persists under the perturbation (1).*

Our proof is based on the study of the persistence of the resonant rotational invariant circles (resonant RICs) of some twist maps by means of a first-order Melnikov method. Only convex caustics can be related to the RICs of those twist maps. Thus, there is no direct way to extend the same procedure to the nonconvex caustic hyperbolas, but we believe that the same results hold for them.

Remark 1. If $\mu_\epsilon(\varphi)$ is constant, then the perturbed curves Q_ϵ are ellipses, so all caustics (resonant or not) are preserved. Hence, the hypothesis $\mu_1(\varphi)$ nonconstant is natural, since we are using a first-order method. Nevertheless, we can still state some results when this hypothesis fails. More precisely, let us assume that

$$\mu_\epsilon(\varphi) = \mu_0 + \epsilon \mu_1 + \dots + \epsilon^{i-1} \mu_{i-1} + \epsilon^i \mu_i(\varphi) + O(\epsilon^{i+1}),$$

for some $\mu_0, \dots, \mu_{i-1} \in \mathbb{R}$ and some nonconstant 2π -periodic entire function $\mu_i(\varphi)$. Then:

- If n is odd, all the (m, n) -resonant elliptical caustics with odd n break up. This result is a corollary of Theorem 1. It suffices to consider $\delta = \epsilon^i$ as the new perturbative parameter, $Q_\epsilon^* = \{\mu \equiv \mu_0 + \dots + \epsilon^{i-1} \mu_{i-1}\}$ as the unperturbed ellipse, and to realize that Q_ϵ is a $O(\delta)$ -perturbation of Q_ϵ^* whose first-order term in δ verifies the hypotheses of Theorem 1.
- If n is even, we believe that all (m, n) -resonant elliptical caustics also break up, even if $\mu_i'(\varphi)$ is π -antiperiodic, but we should use a second-order Melnikov method in order to prove it. Unfortunately, the computations become too cumbersome.

Remark 2. If we write the perturbed ellipse Q_ϵ in Cartesian coordinates as

$$x^2/a^2 + y^2/b^2 + \epsilon P_1(x, y) + O(\epsilon^2) = 1,$$

then $2(a^2 \sin^2 \varphi + b^2 \cos^2 \varphi) \mu_1(\varphi) + ab P_1(a \cos \varphi, b \sin \varphi) = 0$. In particular, the function $\mu_1(\varphi)$ is π -antiperiodic when $P_1(x, y)$ is odd.

Remark 3. The case of perturbed circular tables was studied using similar techniques in [16], but the final result was quite different. Let us recall it for comparison. Any billiard trajectory inside a circle of radius r_0 has some concentric circle of radius $\sqrt{r_0^2 - \lambda^2}$ as caustic, where $0 < \lambda < r_0$ plays the role of a caustic

parameter. If $\lambda = r_0 \sin(m\pi/n)$, then the circular caustic is (m, n) -resonant. Let us write the perturbed circle in polar coordinates (r, θ) as

$$(2) \quad r = r_\epsilon(\theta) = r_0(1 + \epsilon r_1(\theta) + O(\epsilon^2)),$$

for some smooth function $r_\epsilon: \mathbb{T} \rightarrow \mathbb{R}$. Let $\sum_{l \in \mathbb{Z}} \hat{r}_1^l e^{il\theta}$ be the Fourier expansion of $r_1(\theta)$ and $n \geq 2$. If there exists some $l \in n\mathbb{Z} \setminus \{0\}$ such that $\hat{r}_1^l \neq 0$, then the (m, n) -resonant circular caustics do not persist, see [16, Theorem 1]. In particular, it is not known if the (m, n) -resonant circular caustics with odd (respectively, even) n break up when $r_1(\theta)$ is not constant (respectively, $r_1'(\theta)$ is not π -antiperiodic).

We complete this introduction with a note on the organization. In Section 2 we develop a general Melnikov theory to study the persistence of resonant RICs of twist maps. The general setup is adapted to billiard maps in Section 3. Finally, Theorem 1 is proved in Section 4 by analysing the complex singularities of certain elliptic functions, an idea borrowed from [6].

2. BREAK-UP OF RESONANT INVARIANT CURVES IN TWIST MAPS

This section is a generalization of [16, §2], although several hypotheses have been weakened. Namely, the unperturbed map can be nonintegrable, the resonant invariant circle does not need to be horizontal, and the shift on the invariant circles can be nonconstant. In spite of it, the essential idea does not change. A similar theory is contained in [17]. For a general background on twist maps we refer to the book [9, §9.3] or to the review [14].

Let $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$, and $\pi_1: \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T}$ be the natural projection. Sometimes it is convenient to work in the universal cover \mathbb{R} of \mathbb{T} . We will use the coordinates (x, y) for both $\mathbb{T} \times \mathbb{R}$ and \mathbb{R}^2 . The lines of the form $x = \text{constant}$ and $y = \text{constant}$ will be called vertical and horizontal, respectively. A tilde will always denote the lift of a function or set to the universal cover. If g is a real-valued function, $\partial_i g$ denotes the derivative with respect to the i th variable. We will assume that all the considered objects are smooth. Here, smooth means C^∞ . In particular, all the dependences on the perturbative parameter ϵ are assumed to be smooth.

We will consider certain diffeomorphisms defined on an open cylinder of the form $Z = \mathbb{T} \times Y$, for some open bounded interval $Y = (y_-, y_+) \subset \mathbb{R}$. Then $\tilde{Z} = \mathbb{R} \times Y$ is an open strip of the plane. A diffeomorphism $f: Z \rightarrow Z$ is called an *area-preserving twist map* when it preserves area, orientation, and verifies the *twist condition*

$$\partial_2 \tilde{\pi}_1 \tilde{f}(x, y) \neq 0, \quad \forall (x, y) \in \tilde{Z}.$$

If the twist is positive (respectively, negative), then the first iterate of any vertical line tilts to the right (respectively, left). We also assume, although it is not essential, that f verifies some *rigid boundary conditions*. To be more precise, we suppose that the twist map f can be extended continuously to the closed cylinder $\mathbb{T} \times [y_-, y_+]$ as a rigid rotation on the boundaries. That is, there exist some *boundary frequencies* $\omega_\pm \in \mathbb{R}$, $\omega_- < \omega_+$, such that $\tilde{f}(x, y_\pm) = (x + \omega_\pm, y_\pm)$.

Let $D = \{(x, x') \in \mathbb{R}^2 : \omega_- < x' - x < \omega_+\}$. Then there exists a function $h: D \rightarrow \mathbb{R}$ such that $\tilde{f}(x, y) = (x', y')$ if and only if

$$(3) \quad y = -\partial_1 h(x, x'), \quad y' = \partial_2 h(x, x').$$

The function h is called the *generating function* of f . Besides, if $(x'', y'') = \tilde{f}(x', y')$, then

$$(4) \quad \partial_2 h(x, x') + \partial_1 h(x', x'') = 0.$$

We study the dynamics of f , but it is often more convenient to work with the lift \tilde{f} , so we will pass between the two without comment and, in what follows, the lift \tilde{f} remains fixed.

A closed curve $\Upsilon \subset Z$ is said to be a *rotational invariant circle (RIC)* of f when it is homotopically nontrivial and $f(\Upsilon) = \Upsilon$. Birkhoff proved that all RICs are graphs of Lipschitz functions. See, for instance, [14, §IV.C]. Let $v: \mathbb{T} \rightarrow Y$ be the Lipschitz function such that $\Upsilon = \text{graph } v := \{(x, v(x)) : x \in \mathbb{T}\}$. If v is smooth, we say that Υ is a *smooth RIC*.

Twist maps do not form a closed set under composition. For instance, the square of a twist map is not necessarily a twist map, and indeed typically it is not. Nevertheless, any power of a twist map is *locally twist* on its smooth RICs.

Lemma 2. *If $\Upsilon = \text{graph } v$ is a smooth RIC of an area-preserving twist map $f: Z \rightarrow Z$, then*

$$\partial_2 \tilde{\pi}_1 \tilde{f}^n(x, \tilde{v}(x)) \neq 0, \quad \forall x \in \mathbb{R}, \quad \forall n \geq 1.$$

Proof. Given any point $p = (x, \tilde{v}(x)) \in \tilde{\Upsilon}$, let $p_j = (x_j, \tilde{v}(x_j)) = \tilde{f}^j(p)$, $t_j = (1, \tilde{v}'(x_j))$, and $v_j = (0, 1)$. We identify the tangent planes $T_p \tilde{Z}$ with the Euclidean plane \mathbb{R}^2 . Thus, the vector t_j is tangent to $\tilde{\Upsilon}$ at the point p_j and v_j is a vertical vector

The linear map $d\tilde{f}^n(p): T_p \tilde{Z} \rightarrow T_{p_n} \tilde{Z}$ is the composition of the linear maps $d\tilde{f}(p_j): T_{p_j} \tilde{Z} \rightarrow T_{p_{j+1}} \tilde{Z}$ for $j = 0, \dots, n-1$. Let $a_j, b_j, c_j, d_j, \alpha_n, \beta_n, \gamma_n, \delta_n \in \mathbb{R}$ be the coefficients such that

$$\begin{aligned} d\tilde{f}(p_j): t_j &\mapsto a_j t_{j+1} + c_j v_{j+1}, & v_j &\mapsto b_j t_{j+1} + d_j v_{j+1} \\ d\tilde{f}^n(p): t_0 &\mapsto \alpha_n t_n + \gamma_n v_n, & v_0 &\mapsto \beta_n t_n + \delta_n v_n. \end{aligned}$$

We note that $b_j = \partial_2 \tilde{\pi}_1 \tilde{f}(p_j)$ and $\beta_n = \partial_2 \tilde{\pi}_1 \tilde{f}^n(p)$. Let us suppose that the twist is positive, so $b_j > 0$. We want to prove that $\beta_n > 0$ for any integer $n \geq 1$. The case of negative twist is completely analogous.

We deduce that $c_j = 0$ from the invariance of $\tilde{\Upsilon}$. Hence, $\beta_n = \sum_{j=0}^{n-1} D_0^{j-1} b_j A_{j+1}^{n-1}$, where $D_i^j = \prod_{k=i}^j d_k$ and $A_i^j = \prod_{k=i}^j a_k$. Besides, we note that $d_j > 0$ because the two components of $C \setminus \Upsilon$ are invariant. Finally, we get that $a_j > 0$ from the preservation of orientation. \square

Roughly speaking, a RIC is said to be *resonant* when all its points are periodic, but we need to be more precise. Let $(x, y) \in Z$ be a periodic point of the twist map f , and let n be its least period. Then there exists an integer m such that its lift verifies $\tilde{f}^n(x, y) = (x + 2\pi m, y)$. Obviously, $\omega_- < 2\pi m/n < \omega_+$. Such a periodic point is said to be of *type* (m, n) . A RIC is said to be (m, n) -*resonant* when all its points are periodic of type (m, n) .

Let f be an area-preserving twist map with a (m, n) -resonant smooth RIC $\Upsilon = \text{graph } v$. Considering area-preserving twist perturbations of the form $f_\epsilon = f + O(\epsilon)$, we prove in the following lemma that there exists two graphs $\Upsilon_\epsilon = \text{graph } v_\epsilon$ and $\Upsilon_\epsilon^* = \text{graph } v_\epsilon^*$ $O(\epsilon)$ -close to Υ and such that f_ϵ^n projects the first graph onto the second one along the vertical direction.

Lemma 3. *There exist two smooth functions $v_\epsilon, v_\epsilon^*: \mathbb{T} \rightarrow Y$ defined for $\epsilon \in (-\epsilon_0, \epsilon_0)$, $\epsilon_0 > 0$, such that:*

- (1) $v_\epsilon(x) = v(x) + O(\epsilon)$ and $v_\epsilon^*(x) = v(x) + O(\epsilon)$, uniformly in $x \in \mathbb{T}$; and
- (2) $f_\epsilon^n(x, v_\epsilon(x)) = (x, v_\epsilon^*(x))$, for all $x \in \mathbb{T}$.

Proof. We work with the lift of the maps. Once fixed an angle $x \in \mathbb{R}$, let $y_0 = \tilde{v}(x)$ and

$$\tilde{G}(y, \epsilon) := \tilde{\pi}_1 \tilde{f}_\epsilon^n(x, y) - x - 2\pi m.$$

This function $\tilde{G}(y, \epsilon)$ verifies the hypotheses of the Implicit Function Theorem at the point $(y, \epsilon) = (y_0, 0)$, since $\tilde{G}(y_0, 0) = 0$ and $\partial_1 \tilde{G}(y_0, 0) = \partial_2 \tilde{\pi}_1 \tilde{f}^n(x, \tilde{v}(x)) \neq 0$, see Lemma 2. Consequently, there exist $\epsilon_0, \eta > 0$ such that the equation $\tilde{G}(y, \epsilon) = 0$ has exactly one solution $y_\epsilon = y_0 + O(\epsilon)$ in the interval $(y_0 - \eta, y_0 + \eta)$ for all $\epsilon \in (-\epsilon_0, \epsilon_0)$. We recall that $\tilde{G}(y, \epsilon)$ had $x \in \mathbb{R}$ as an extra parameter, but it appeared in a 2π -periodic smooth way. Hence, ϵ_0 and η can be taken independent from x , the estimate $|y_\epsilon - y_0| = O(\epsilon)$ is uniform in x , and y_ϵ depends in a 2π -periodic smooth way on x . Finally, set $\tilde{v}_\epsilon(x) = y_\epsilon$ and then $\tilde{v}_\epsilon^*(x)$ is determined by means of relation $\tilde{f}_\epsilon^n(x, \tilde{v}_\epsilon(x)) = (x + 2\pi m, \tilde{v}_\epsilon^*(x))$. The functions $\tilde{v}_\epsilon, \tilde{v}_\epsilon^*: \mathbb{R} \rightarrow Y$ are 2π -periodic and smooth, so they can be projected to two smooth functions $v_\epsilon, v_\epsilon^*: \mathbb{T} \rightarrow Y$ that verify the two claimed properties by construction. \square

We say that a (m, n) -resonant smooth RIC Υ of a twist map f *persists* under an area-preserving twist perturbation $f_\epsilon = f + O(\epsilon)$ whenever the perturbed map has a (m, n) -resonant RIC Υ_ϵ for any small enough ϵ such that $\Upsilon_\epsilon = \Upsilon + O(\epsilon)$. The corollary below follows immediately from this definition.

Corollary 4. *The resonant RIC Υ persists under the perturbation f_ϵ if and only if $\Upsilon_\epsilon = \Upsilon_\epsilon^*$.*

Therefore, it is rather useful to quantify the separation between the graphs Υ_ϵ and Υ_ϵ^* .

Lemma 5. $v_\epsilon^*(x) - v_\epsilon(x) = L'_\epsilon(x)$, where $L_\epsilon: \mathbb{T} \rightarrow \mathbb{R}$ is a function whose lift is

$$(5) \quad \tilde{L}_\epsilon(x) = \sum_{j=0}^{n-1} h_\epsilon(\bar{x}_j(x; \epsilon), \bar{x}_{j+1}(x; \epsilon)), \quad \bar{x}_j(x; \epsilon) = \tilde{\pi}_1 \tilde{f}_\epsilon^j(x, \tilde{v}_\epsilon(x)),$$

and h_ϵ is the generating function of f_ϵ .

Proof. As long as confusion is avoided, we will omit the dependence on x and ϵ . We introduce the notations $(\bar{x}_j, \bar{y}_j) = \tilde{f}^j(x, \tilde{v}(x))$ and $\bar{w}_j = \partial \bar{x}_j / \partial x$ for $j = 0, \dots, n$. Then $\bar{x}_0 = x$ and $\bar{x}_n = x + 2\pi m$, so $\bar{w}_0 = \bar{w}_n = 1$. Besides, $\bar{y}_0 = \tilde{v}(x)$ and $\bar{y}_n = \tilde{v}^*(x)$. From the implicit equations (3), we get that $\partial_1 h(\bar{x}_0, \bar{x}_1) = -\bar{y}_0$, $\partial_2 h(\bar{x}_{n-1}, \bar{x}_n) = \bar{y}_n$, and $\partial_2 h(\bar{x}_{j-1}, \bar{x}_j) + \partial_1 h(\bar{x}_j, \bar{x}_{j+1}) = 0$ for $j = 1, \dots, n-1$. Therefore, $\tilde{L}'(x) = \partial_1 h(\bar{x}_0, \bar{x}_1) \bar{w}_0 + \sum_{j=1}^{n-1} (\partial_2 h(\bar{x}_{j-1}, \bar{x}_j) + \partial_1 h(\bar{x}_j, \bar{x}_{j+1})) \bar{w}_j + \partial_2 h(\bar{x}_{n-1}, \bar{x}_n) \bar{w}_n = \tilde{v}^*(x) - \tilde{v}(x)$. It is immediate to check that $\tilde{L}: \mathbb{R} \rightarrow \mathbb{R}$ is 2π -periodic, so it can be projected to a function $L: \mathbb{T} \rightarrow \mathbb{R}$. \square

Corollary 6. *The resonant RIC Υ persists under the perturbation f_ϵ if and only if $L'_\epsilon(x) \equiv 0$.*

We shall say that $L_\epsilon: \mathbb{T} \rightarrow \mathbb{R}$ is the *subharmonic potential* of the resonant RIC Υ under the twist perturbation f_ϵ . It is rather natural to extract information from the low-order terms of its expansion $L_\epsilon(x) = L_0(x) + \epsilon L_1(x) + O(\epsilon^2)$. This is the main idea behind any Melnikov approach to a perturbative problem. The zero-order term $L_0(x)$ is constant (and so useless), since $L'_0(x) = v_0^*(x) - v_0(x) = v(x) - v(x) \equiv 0$. We shall say that the first-order term $L_1(x)$ is the *subharmonic Melnikov potential* of the resonant RIC Υ under the twist perturbation f_ϵ . The proposition below provides a closed formula for its computation.


Proposition 7. *If $h_\epsilon = h + \epsilon h_1 + O(\epsilon^2)$, then the lift of $L_1(x)$ is*

$$\tilde{L}_1(x) = \sum_{j=0}^{n-1} h_1(x_j, x_{j+1}), \quad x_j = \tilde{\pi}_1 \tilde{f}^j(x, \tilde{v}(x)).$$

Proof. Given any $x \in \mathbb{R}$, we set $x_j = x_j(x) := \bar{x}_j(x; 0)$ and $z_j = z_j(x) := \partial_2 \bar{x}_j(x; 0)$ for $j = 0, \dots, n$. Then the $O(\epsilon)$ -term of (5) is

$$\begin{aligned} \tilde{L}_1(x) &= \partial_1 h(x_0, x_1) z_0 + \sum_{j=1}^{n-1} \left(\partial_1 h(x_j, x_{j+1}) + \partial_2 h(x_{j-1}, x_j) \right) z_j + \\ &\quad + \partial_2 h(x_{n-1}, x_n) z_n + \sum_{j=0}^{n-1} h_1(x_j, x_{j+1}). \end{aligned}$$

Using the implicit equations (3) for the unperturbed twist map, the first summation vanishes. The terms $\partial_1 h(x_0, x_1) z_0$ and $\partial_2 h(x_{n-1}, x_n) z_n$ also vanish, since $\bar{x}_0(x; \epsilon) = x$ and $\bar{x}_n(x; \epsilon) = x + 2\pi m$ for all $\epsilon \in (-\epsilon_0, \epsilon_0)$. Besides, $x_j = x_j(x) = \bar{x}_j(x; 0) = \tilde{\pi}_1 \tilde{f}^j(x, v(x))$. \square

FIGURE 1. The billiard map $f(\varphi, \vartheta) = (\varphi', \vartheta')$.

The following corollary displays the most important property of the subharmonic Melnikov potential in relation with the goals of this paper.

Corollary 8. *If $L_1(x)$ is not constant, then the resonant RIC Υ does not persist under the perturbation f_ϵ .*

Proof. It follows directly from Corollary 6 and the estimate $L_\epsilon = \text{constant} + \epsilon L_1 + O(\epsilon^2)$. \square

3. BREAK-UP OF RESONANT CAUSTICS IN PERTURBED BILLIARD TABLES

Let Q be a closed strictly convex smooth curve in the plane. Let $\gamma: \mathbb{T} \rightarrow Q$ be a counterclockwise parametrization. Let $Z = \mathbb{T} \times (0, \pi)$ be an open cylinder. We can model the billiard dynamics inside Q by means of a map $f: Z \rightarrow Z$, $f(\varphi, \vartheta) = (\varphi', \vartheta')$, defined as follows. If the particle hits Q at a point $\gamma(\varphi)$ under an angle of incidence $\vartheta \in (0, \pi)$ with the tangent vector at $\gamma(\varphi)$, then, as the motion is free inside Q , the next impact point is $\gamma(\varphi')$, the intersection point with the boundary and the next angle of incidence is $\vartheta' \in (0, \pi)$, as in Figure 1. A straightforward computation shows that $f(\varphi, \vartheta) = (\varphi', \vartheta')$ if and only if

$$(6) \quad |\gamma'(\varphi)| \cos \vartheta = -\partial_1 h(\varphi, \varphi'), \quad |\gamma'(\varphi')| \cos \vartheta' = \partial_2 h(\varphi, \varphi'),$$

where $h: \mathbb{T}^2 \setminus \{\varphi' \neq \varphi\} \rightarrow \mathbb{R}$ is given by $h(\varphi, \varphi') = |\gamma(\varphi) - \gamma(\varphi')|$. Besides, the twist condition holds: $\partial \varphi' / \partial \vartheta = h(\varphi, \varphi') / |\gamma'(\varphi')| \sin \vartheta' > 0$. Finally, it is geometrically clear that f verifies the rigid boundary conditions with $\omega_- = 0$ and $\omega_+ = 2\pi$.

A remark is in order. Equations (6) differ slightly from equations (3), but identity (4) still holds and so the theory developed in the previous section still applies.




FIGURE 2. Left: A $(1,4)$ -resonant convex smooth caustic C . Right: Its two smooth RICs $\Upsilon^- = \text{graph } \vartheta^-$ and $\Upsilon^+ = \text{graph } \vartheta^+$ in the phase space $Z = \mathbb{T} \times (0, \pi)$.

Obviously, one could write the map in the canonical coordinates —arclength parameter for the boundary and $\cos \vartheta$ as its conjugate— in order to have h as a generating function, but this is not a wise choice when dealing with ellipses.

Let us assume that there exists a closed convex smooth caustic C contained in the region enclosed by Q . Then the billiard map $f: Z \rightarrow Z$ has two smooth RICs $\Upsilon^\pm = \text{graph } \vartheta^\pm \subset Z$. The functions $\vartheta^\pm: \mathbb{T} \rightarrow (0, \pi)$ are easy to understand: $\vartheta^+(\varphi)$ and $\vartheta^-(\varphi)$ are the angles determined by the two tangent lines to the caustic C from the point $\gamma(\varphi) \in Q$, see Figure 2. In particular, $\vartheta^-(\varphi) + \vartheta^+(\varphi) = \pi$. To fix ideas, we will assume that Υ^- and Υ^+ correspond to the billiard motion around C in the counterclockwise and clockwise senses, respectively. Hence, $0 < \vartheta^-(\varphi) < \pi/2 < \vartheta^+(\varphi) < \pi$. There is an explicit formula relating the parametrization of the billiard curve Q , the parametrization of the caustic C , and the functions ϑ^\pm . See, for instance, [7, 10].

Let Q be a closed strictly convex smooth billiard boundary with a (m, n) -resonant convex caustic C , so that its RIC Υ^- is (m, n) -resonant and its RIC Υ^+ is $(n-m, n)$ -resonant. We say that C *persists* under a perturbation $Q_\epsilon = Q + O(\epsilon)$ whenever the perturbed billiard curve has a (m, n) -resonant caustic C_ϵ for any small enough ϵ such that $C_\epsilon = C + O(\epsilon)$.

Let f_ϵ be the billiard map inside Q_ϵ and $L_1^-(\varphi)$ and $L_1^+(\varphi)$ be the subharmonic Melnikov potentials of the resonant RICs Υ^- and Υ^+ under the area-preserving twist perturbation f_ϵ . Both potentials coincide, due to the time reversibility of the billiard dynamics. Therefore, we can skip the \pm signs. In this context, we will say that $L_1(\varphi)$ is the subharmonic Melnikov potential of the resonant caustic C for the perturbation Q_ϵ .

Corollary 9. *If $L_1(\varphi)$ is not constant, then the resonant caustic C does not persist under the perturbation Q_ϵ .*

4. BREAK-UP OF RESONANT CAUSTICS IN PERTURBED ELLIPTIC BILLIARD TABLES

From now on, we will assume that the unperturbed billiard boundary is the ellipse

$$Q = \left\{ q = (x, y) \in \mathbb{R}^2 : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \right\}, \quad a > b > 0.$$

It is known that the convex caustics of the billiard inside Q are the confocal ellipses

$$C_\lambda = \left\{ q = (x, y) \in \mathbb{R}^2 : \frac{x^2}{a^2 - \lambda^2} + \frac{y^2}{b^2 - \lambda^2} = 1 \right\}, \quad 0 < \lambda < b.$$

Let $\rho(\lambda)$ be the rotation number of the elliptical caustic C_λ . Then $\rho: (0, b) \rightarrow \mathbb{R}$ is an analytic increasing function such that $\rho(0) = 0$ and $\rho(b) = 1/2$. See, for instance, [4]. Thus, there is a unique (m, n) -resonant elliptical caustic for any relatively prime integers m and n such that $1 \leq m < n/2$. We shall see that the caustic parameter $\lambda \in (0, b)$ of the (m, n) -resonant caustic is implicitly determined by means of an equation containing a couple of elliptic integrals, see equation (10).

The following lemma on elliptic billiards is useful to simplify the expression of the subharmonic Melnikov potential later on.

Lemma 10. *Let $(q_j)_{j \in \mathbb{Z}}$ be any billiard trajectory inside the ellipse Q with caustic C_λ . Let $p_j = (q_{j+1} - q_j)/|q_{j+1} - q_j|$ be the unit inward velocities of the trajectory. Then*

$$ab \langle p_{j-1} - p_j, D^{-2} q_j \rangle = 2\lambda, \quad \forall j \in \mathbb{Z},$$

where $D = \text{diag}(a, b)$ is the diagonal matrix such that $Q = \{q \in \mathbb{R}^2 : \langle q, D^{-2} q \rangle = 1\}$.

Proof. We shall prove that given any point $q = (x, y) \in Q$ and any unit inward vector $p = (u, v) \in \mathbb{S}^1$, the line $\ell = \{q + \tau p : \tau \in \mathbb{R}\}$ is tangent to the conic C_λ if and only if

$$\lambda = -(bxu/a + ayv/b) = -ab \langle p, D^{-2} q \rangle.$$

To begin with, we note that the line ℓ is tangent to the conic C_λ if and only if the equation of second order in the variable τ given by

$$(x + \tau u)^2/(a^2 - \lambda^2) + (y + \tau v)^2/(b^2 - \lambda^2) - 1 = 0$$

has zero discriminant, which is equivalent to the equation

$$\left(\frac{xu}{a^2 - \lambda^2} + \frac{yv}{b^2 - \lambda^2} \right)^2 = \left(\frac{u^2}{a^2 - \lambda^2} + \frac{v^2}{b^2 - \lambda^2} \right) \left(\frac{x^2}{a^2 - \lambda^2} + \frac{y^2}{b^2 - \lambda^2} - 1 \right).$$

After some simplifications, we can rewrite this equation as

$$(xv - yu)^2 = (b^2 - \lambda^2)u^2 + (a^2 - \lambda^2)v^2 = a^2v^2 + b^2u^2 - \lambda^2,$$

since $u^2 + v^2 = 1$. Next, using that $x^2/a^2 + y^2/b^2 = 1$, we obtain that

$$\lambda^2 = (a^2v^2 + b^2u^2)(x^2/a^2 + y^2/b^2) - (xv - yu)^2 = (bxu/a + ayv/b)^2.$$

Thus, we have two possibilities: $\lambda = ab\langle p, D^{-2}q \rangle$ or $\lambda = -ab\langle p, D^{-2}q \rangle$. The first one is discarded, because $\lambda > 0$ and $\langle p, D^{-2}q \rangle < 0$. The second inequality follows from the fact that the vector p points inward Q at q , whereas $D^{-2}q$ is an outward normal vector to Q at q .

Finally, we note that $-p_{j-1} = (q_{j-1} - q_j)/|q_{j-1} - q_j|$ and $p_j = (q_{j+1} - q_j)/|q_{j+1} - q_j|$ are the two unit vectors that point inward Q at the impact point q_j and give the two tangent directions to the caustic C_λ . Therefore, $\lambda = ab\langle p_{j-1}, D^{-2}q_j \rangle = -ab\langle p_j, D^{-2}q_j \rangle$. \square

Proposition 11. *Let C_λ be the (m, n) -resonant elliptical caustic confocal to the ellipse Q . Given any angle $\varphi \in \mathbb{T}$, let $q_j = (a \cos \varphi_j, b \sin \varphi_j)$ be the vertexes of the (m, n) -gon inscribed in Q and circumscribed around C_λ such that $q_0 = (a \cos \varphi, b \sin \varphi)$. Then the subharmonic Melnikov potential of the caustic C_λ for the perturbed ellipse (1) is*

$$(7) \quad L_1(\varphi) = 2\lambda \sum_{j=0}^{n-1} \mu_1(\varphi_j).$$

Proof. The parametrization of the perturbed ellipse (1) is given by

$$\gamma_\epsilon(\varphi) = (c \cosh \mu_\epsilon(\varphi) \cos \varphi, c \sinh \mu_\epsilon(\varphi) \sin \varphi) = \gamma_0(\varphi) + \epsilon \gamma_1(\varphi) + O(\epsilon^2),$$

where $\gamma_0(\varphi) = (a \cos \varphi, b \sin \varphi)$, $\gamma_1(\varphi) = ab\mu_1(\varphi)D^{-2}\gamma_0(\varphi)$, and $D = \text{diag}(a, b)$ as above. The generating function of the billiard map inside the perturbed ellipse is

$$h_\epsilon(\varphi, \varphi') = |\gamma_\epsilon(\varphi') - \gamma_\epsilon(\varphi)| = h_0(\varphi, \varphi') + \epsilon h_1(\varphi, \varphi') + O(\epsilon^2).$$

The first terms of this expansion verify the identities $h_0(\varphi, \varphi') = |\gamma_0(\varphi') - \gamma_0(\varphi)|$ and $h_0(\varphi, \varphi')h_1(\varphi, \varphi') = \langle \gamma_0(\varphi') - \gamma_0(\varphi), \gamma_1(\varphi') - \gamma_1(\varphi) \rangle$.

Let $(q_j)_{j \in \mathbb{Z}}$ be the billiard trajectory inside the ellipse Q with caustic C_λ such that $q_j = \gamma_0(\varphi_j)$ and $\varphi_0 = \varphi$. The unit inward velocities of this trajectory are

$$p_j = \frac{q_{j+1} - q_j}{|q_{j+1} - q_j|} = \frac{\gamma_0(\varphi_{j+1}) - \gamma_0(\varphi_j)}{h_0(\varphi_j, \varphi_{j+1})}.$$

It follows from Proposition 7 that the subharmonic Melnikov potential is

$$\begin{aligned} L_1(\varphi) &= \sum_{j=0}^{n-1} h_1(\varphi_j, \varphi_{j+1}) \\ &= \sum_{j=0}^{n-1} \langle p_j, \gamma_1(\varphi_{j+1}) - \gamma_1(\varphi_j) \rangle \\ &= ab \sum_{j=0}^{n-1} \langle p_j, \mu_1(\varphi_{j+1})D^{-2}q_{j+1} - \mu_1(\varphi_j)D^{-2}q_j \rangle \end{aligned}$$

$$\begin{aligned}
&= ab \sum_{j=0}^{n-1} \langle p_{j-1} - p_j, D^{-2} q_j \rangle \mu_1(\varphi_j) \\
&= 2\lambda \sum_{j=0}^{n-1} \mu_1(\varphi_j).
\end{aligned}$$

We have used the periodicity in the fourth equality and Lemma 10 in the last one. \square

Next, we give a couple of sufficient conditions for the subharmonic Melnikov potential to be constant. These conditions are trivial. Nevertheless, they play a key role in our problem. Concretely, we shall check later on that they are also necessary conditions in the class of 2π -periodic entire functions $\mu_1(\varphi)$.

Corollary 12. *Let $\mu_1(\varphi)$ be any 2π -periodic smooth function.*

- (1) *If the period n is odd, then $\mu_1(\varphi)$ constant $\Rightarrow L_1(\varphi)$ constant.*
- (2) *If the period n is even, then $\mu'_1(\varphi)$ π -antiperiodic $\Rightarrow L_1(\varphi)$ constant.*

Proof. The case n odd is obvious. If n is even, the (m, n) -gons inscribed in Q and circumscribed around C_λ are symmetric with respect to the origin, so $\varphi_{j+n/2} = \varphi_j + \pi$ and

$$L'_1(\varphi) = 2\lambda \sum_{j=0}^{n-1} \mu'_1(\varphi_j) = 2\lambda \sum_{j=0}^{n/2-1} (\mu'_1(\varphi_j) + \mu'_1(\varphi_j + \pi)).$$

In particular, n even and $\mu'_1(\varphi)$ π -antiperiodic $\Rightarrow L'_1(\varphi) \equiv 0 \Rightarrow L_1(\varphi)$ constant. \square

The subharmonic Melnikov potential of the (m, n) -resonant caustic for the perturbed circle (2) is

$$(8) \quad L_1(\theta) = 2r_0 \sin(m\pi/n) \sum_{j=0}^{n-1} r_1(\theta_j), \quad \theta_j = \theta + 2\pi m j/n,$$

see [16, Proposition 10]. We recall that $\lambda = r_0 \sin(m\pi/n)$ is the (m, n) -resonant caustic parameter of the circle of radius r_0 . Besides, all the (m, n) -gons inscribed in the circle of radius r_0 and circumscribed around the circle of radius $\lambda = r_0 \sin(m\pi/n)$ are regular, so their vertexes are of the form $q_j = (r_0 \cos \theta_j, r_0 \sin \theta_j)$ with $\theta_j = \theta + 2\pi m j/n$. Hence, the function (8) is the limit of function (7) when both a and b tend to r_0 .

Although functions (7) and (8) look quite similar, they hide a crucial difference. There is a simple formula for the θ_j angles, but not for the φ_j ones. This has to do with the fact that the billiard trajectories inside a circle of radius r_0 sharing a circular caustic with radius $\lambda = r_0 \sin(\delta/2)$ have a rigid angular dynamics of the form $\theta \mapsto \theta + \delta$. On the contrary, such a rigid angular dynamics does not take place for elliptic tables when the angle φ is considered, which is a source

of technical difficulties in the study of the subharmonic Melnikov potential (7). Nevertheless, it is possible to define a new angular parameter t over the ellipse Q in such a way that all billiard trajectories inside Q sharing the elliptical caustic C_λ have a rigid angular dynamics of the form $t \mapsto t + \delta$, for some constant shift $\delta = \delta(\lambda)$.

We need some notations on elliptic functions in order to define this angular parameter t . We refer to [1, 19] for a general background on elliptic functions. Given a quantity $k \in (0, 1)$, called the *modulus*, then $K = K(k) = \int_0^{\pi/2} (1 - k^2 \sin^2 \phi)^{-1/2} d\phi$ is the *complete elliptic integral of the first kind*. We also write $K' = K'(k) = K(\sqrt{1 - k^2})$. The *amplitude* function $\varphi = \text{am } t$ is defined through the inversion of the integral

$$t = \int_0^\varphi (1 - k^2 \sin^2 \phi)^{-1/2} d\phi.$$

Then the *elliptic sinus* and the *elliptic cosinus* are defined by the trigonometric relations

$$\text{sn } t = \sin \varphi, \quad \text{cn } t = \cos \varphi,$$

respectively. Dependence on the modulus is denoted by a comma preceding it, so we can write $\text{am}(t, k)$, $\text{sn}(t, k)$, and $\text{cn}(t, k)$ to avoid any confusion. In the following lemma it is stated that the angular dynamics becomes rigid in the angular parameter t given by $\varphi = \text{am}(t, k)$. It suffices to find the suitable modulus k for each elliptical caustic C_λ .

Lemma 13. *Once fixed any caustic parameter $\lambda \in (0, b)$, we set the modulus $k \in (0, 1)$ and the constant shift $\delta \in (0, 2K)$ by the formulae*

$$(9) \quad k^2 = \frac{a^2 - b^2}{a^2 - \lambda^2}, \quad \delta/2 = \int_0^{\vartheta/2} (1 - k^2 \sin^2 \phi)^{-1/2} d\phi,$$

where $\vartheta \in (0, \pi)$ is the angle such that $\sin(\vartheta/2) = \lambda/b$. Let

$$q_j = (a \cos \varphi_j, b \sin \varphi_j) = (a \text{cn}(t_j, k), b \text{sn}(t_j, k))$$

be any billiard trajectory inside the ellipse Q with caustic C_λ . Then $t_{j+1} = t_j + \delta$.

Proof. By definition, $\varphi_j = \text{am}(t_j, k)$, so $t_{j+1} - t_j = \int_{\varphi_j}^{\varphi_{j+1}} (1 - k^2 \sin^2 \phi)^{-1/2} d\phi$. These integrals are equal to a constant δ that depends only on C_λ , see [5, page 1543]). The formula for the constant shift is given in [5, page 1540]. \square

Remark that if $a = b = r_0$ then the modulus k is equal to zero, the complete elliptic integral K is equal to $\pi/2$, the amplitude function is the identity, the elliptic sinus/cosinus are the usual sinus/cosinus, the shift $\delta \in (0, \pi)$ is given by $\lambda = r_0 \sin(\delta/2)$, and the dynamical relation $t_{j+1} = t_j + \delta$ becomes $\varphi_{j+1} = \varphi_j + \delta$. Thus, we recover the known rigid angular dynamics for circular tables as a limit of the formulae for elliptic tables.

From now on, k and δ will denote the modulus and the constant shift defined in (9). Thus, we shall skip the dependence of the elliptic functions on the modulus.

We note that C_λ has eccentricity k . Besides, C_λ is the (m, n) -resonant elliptical caustic if and only if

$$(10) \quad n\delta = 4Km.$$

This identity has the following geometric interpretation. When a billiard trajectory makes one turn around C_λ , the old angular variable φ changes by 2π , so the new angular variable t changes by $4K$. On the other hand, we have seen that the variable t changes by δ when a billiard trajectory bounces once. Hence, a billiard trajectory inscribed in Q and circumscribed around C_λ makes exactly m turns around C_λ after n bounces if and only if (10) holds.

Proposition 14. *Let $\mu_1(\varphi)$ be any 2π -periodic entire function.*

- (1) *If the period n is odd, then $L_1(\varphi)$ constant $\Leftrightarrow \mu_1(\varphi)$ constant.*
- (2) *If the period n is even, then $L_1(\varphi)$ constant $\Leftrightarrow \mu_1'(\varphi)$ π -antiperiodic.*

Proof. Let $\Delta = 2K + 2K'i$ and $z(t) = \operatorname{cn} t + i \operatorname{sn} t$. If $\varphi = \operatorname{am} t$, then

$$\begin{aligned} e^{i\varphi} &= \cos \varphi + i \sin \varphi = \operatorname{cn} t + i \operatorname{sn} t = z(t), \\ e^{-i\varphi} &= \cos \varphi - i \sin \varphi = \operatorname{cn} t - i \operatorname{sn} t = z(t + \Delta). \end{aligned}$$

We have used that the elliptic cosinus is Δ -periodic, but the elliptic sinus is Δ -antiperiodic. We also recall that the elliptic cosinus/sinus are $2K$ -antiperiodic meromorphic functions on the whole complex plane whose unique singularities are the points of the form

$$\tau_{r,s} = 2Kr + (1 + 2s)K'i, \quad r, s \in \mathbb{Z}.$$

Besides, these singularities are just simple poles whose residues are

$$\operatorname{res}(\operatorname{cn}; \tau_{r,s}) = (-1)^{r+s+1}i/k, \quad \operatorname{res}(\operatorname{sn}; \tau_{r,s}) = (-1)^r/k.$$

Thus, $z(t)$ is a $2K$ -antiperiodic meromorphic function whose unique singularities are the points of the set

$$P = \{\tau_{r,2s+1} : r, s \in \mathbb{Z}\} = \tau_* + 2K\mathbb{Z} + 4K'i\mathbb{Z}, \quad \tau_* = \tau_{0,-1} = -K'i.$$

As before, these singularities are just simple poles.

Let $\sum_{l \in \mathbb{Z}} \hat{\mu}_l e^{il\varphi}$ be the Fourier expansion of $\mu_1(\varphi)$. Then

$$\mu_1(\operatorname{am} t) = \mu_1(\varphi) = \sum_{l \in \mathbb{Z}} \hat{\mu}_l e^{il\varphi} = \hat{\mu}_-(z(t + \Delta)) + \hat{\mu}_0 + \hat{\mu}_+(z(t)),$$

where $\hat{\mu}_-(z) = \sum_{l=1}^{\infty} \hat{\mu}_{-l} z^l$ and $\hat{\mu}_+(z) = \sum_{l=1}^{\infty} \hat{\mu}_l z^l$. We note that the functions $\hat{\mu}_\pm(z)$ are entire, because $\mu_1(\varphi)$ is entire. Besides,

$$(11) \quad L_1(\operatorname{am} t) = L_1(\varphi) = 2\lambda \sum_{j=0}^{n-1} \mu_1(\varphi_j) = 2\lambda (L_-(t) + n\hat{\mu}_0 + L_+(t)),$$

where $L_-(t) = \sum_{j=0}^{n-1} \hat{\mu}_-(z(t + \Delta + j\delta))$ and $L_+(t) = \sum_{j=0}^{n-1} \hat{\mu}_+(z(t + j\delta))$. Let us study the behaviour of these two functions around the point $\tau_* = -K'i$.

Concretely, we shall prove that $L_-(t)$ is analytic at $t = \tau_*$, whereas $L_+(t)$ has a nonremovable singularity at $t = \tau_*$ provided $\mu_1(\varphi)$ is nonconstant and n is odd, or provided $\mu'_1(\varphi)$ is not π -antiperiodic and n is even.

We begin with a couple of simple observations. If $j \in \{0, \dots, n-1\}$, then:

- a) $\Im(\tau_* + \Delta + j\delta) = K'$, so $\tau_* + \Delta + j\delta \notin P$; and
- b) $\tau_* + j\delta \in P \Leftrightarrow 4Km/n = j\delta \in 2K\mathbb{Z} \Leftrightarrow 2jm \in n\mathbb{Z} \Leftrightarrow 2j \in n\mathbb{Z} \Leftrightarrow j \in \{0, n/2\}$. Here, we have used that $\delta \in \mathbb{R}$, equation (10), and $\gcd(m, n) = 1$. Besides, we stress that the equality $j = n/2$ only can take place when n is even.

We deduce the following results from the above observations.

- 1) $L_-(t)$ is analytic at $t = \tau_*$, because so are $z(t + \Delta + j\delta)$ for $j = 0, \dots, n-1$.
- 2) If n is odd and $\mu_1(\varphi)$ is nonconstant, then:
 - The function $\hat{\mu}_+(z)$ is nonconstant and entire;
 - The function $L_+(t) - \hat{\mu}_+(z(t)) = \sum_{j=1}^{n-1} \hat{\mu}_+(z(t + j\delta))$ is analytic at $t = \tau_*$;
 - The composition $\hat{\mu}_+(z(t))$ has a nonremovable singularity at $t = \tau_*$; and
 - The function (11) is nonconstant, since it has a nonremovable singularity at $t = \tau_*$.
- 3) If n is even and $\mu'_1(\varphi)$ is not π -antiperiodic, then:
 - The sum $\hat{\sigma}(z) = \hat{\mu}_+(z) + \hat{\mu}_+(-z) = 2 \sum_{l=1}^{\infty} \hat{\mu}_{2l} z^{2l}$ is a nonconstant entire function;
 - $z(t + n\delta/2) = z(t + 2Km) = (-1)^m z(t) = -z(t)$, since m is odd;
 - $\hat{\mu}_+(z(t)) + \hat{\mu}_+(z(t + n\delta/2)) = \hat{\sigma}(z(t))$;
 - The function $L_+(t) - \hat{\sigma}(z(t))$ is analytic at $t = \tau_*$;
 - The composition $\hat{\sigma}(z(t))$ has a nonremovable singularity at $t = \tau_*$; and
 - The function (11) is nonconstant, since it has a nonremovable singularity at $t = \tau_*$.

Therefore, the proof follows by combining the above results with Corollary 12. \square

Finally, we note that our main result (namely, Theorem 1 stated in the introduction) follows directly from Corollary 9 and Proposition 14.

Acknowledgements: SP-de-C was partially supported by the CRM and Brazilian agencies CNPq and FAPEMIG. RR-R was supported in part by MICINN-FEDER Grant MTM2009-06973 (Spain) and CUR-DIUE Grant 2009SGR859 (Catalonia).

This work was completed while SP-de-C was a visitor at the CRM at Barcelona (Spain). Useful conversations with Pablo S. Casas, Amadeu Delshams, Vadim Kaloshin, and Vassilios Rothos are gratefully acknowledged.

REFERENCES

- [1] Abramowitz M. and Stegun I., *Handbook of Mathematical Functions*, (New York: Dover) 1972.
- [2] Baryshnikov Yu and Zharnitsky V., Sub-Riemannian geometry and periodic orbits in classical billiards, *Math. Res. Lett.* **13** (2006), 587–598.
- [3] Birkhoff G D 1927 *Dynamical Systems* (*Am. Math. Soc. Coll. Pub. vol. 9*) (Providence RI: American Mathematical Society).
- [4] Casas P S and Ramírez-Ros R., The frequency map for elliptic billiards, *SIAM J. Appl. Dyn. Syst.*, **10** (2011), 278–324
- [5] Chang S-J. and Friedberg R., Elliptical billiards and Poncelet’s theorem, *J. Math. Phys.*, **29** (1988), 1537–50.
- [6] Delshams A and Ramírez-Ros R 1996 Poincaré-Melnikov-Arnold method for analytic planar maps, *Nonlinearity*, **9** /1996), 1–26.
- [7] Douady R., *Applications du théorème des tores invariantes* (Paris: PhD. These Université Paris VII) (1982).
- [8] Gutkin E., and Katok A., Caustics for inner and outer billiards, *Comm. Math. Phys.*, **173** (1995), 101–133.
- [9] Katok A and Hasselblatt B., *Introduction to the Modern Theory of Dynamical Systems*, Cambridge: Cambridge Univ. Press, (1995).
- [10] Knill O., On nonconvex caustics of convex billiards, *Elem. Math.*, **53** (1998), 89–106.
- [11] Kozlov V V and Treshchëv D., *Billiards: a Genetic Introduction to the Dynamics of Systems with Impacts* (*Transl. Math. Monographs 89*), Providence, RI: American Mathematical Society, (1991).
- [12] Lazutkin V.F., The existence of caustics for a billiard problem in a convex domain, *Math. USSR Izvestija*, **7** (1973), 185–214.
- [13] Mather J., Glancing billiards, *Ergodic Theory Dyn. Syst.*, **2** (1982), 397–403.
- [14] Meiss J.D., Symplectic maps, variational principles, and transport, *Rev. Mod. Phys.*, **64** (1992), 795–848.
- [15] Poncelet J V 1822 *Traité des propriétés projectives des figures* (Mett-Paris)
- [16] Ramírez-Ros R., Break-up of resonant invariant curves in billiards and dual billiards associated to perturbed circular tables, *Physica D.*, **214** (2006), 78–87.
- [17] Rothos V., Subharmonic bifurcations of localized solutions of a discrete NLS equation, *Discrete Contin. Dyn. Syst.*, supplement volume (2005), 756–67.
- [18] Tabachnikov S., Billiards, *Panor. Synth.*, **1** (1995), vi+142.
- [19] Whittaker E.T. and Watson G.N., *A Course of Modern Analysis*, Cambridge: Cambridge University Press, (1927).

SÔNIA PINTO-DE-CARVALHO
 DEPARTAMENTO DE MATEMÁTICA
 ICEx
 UNIVERSIDADE FEDERAL DE MINAS GERAIS
 CP 702, 30123-970, BELO HORIZONTE
 MG, BRAZIL
E-mail address: sonia@mat.ufmg.br

RAFAEL RAMÍREZ-ROS
DEPARTAMENT DE MATEMÀTICA APLICADA I
UNIVERSITAT POLITÈCNICA DE CATALUNYA
DIAGONAL 647
08028 BARCELONA
SPAIN
E-mail address: `Rafael.Ramirez@upc.edu`