

Moduli of smoothness and growth properties of Fourier transforms: two-sided estimates

D. Gorbachev and S. Tikhonov

ABSTRACT. We prove two-sided inequalities between the integral moduli of smoothness of a function on $\mathbb{R}^d/\mathbb{T}^d$ and the weighted tail-type integrals of its Fourier transform/series. Sharpness of obtained results in particular is given by the equivalence results for functions satisfying certain regular conditions. Applications include a quantitative form of the Riemann–Lebesgue lemma as well as several other questions in approximation theory and the theory of function spaces.

1. Introduction

This paper studies the interrelation between the smoothness of a function and growth properties of Fourier transforms/coefficients. Let us first recall the classical Riemann–Lebesgue lemma: $|\widehat{f}_n| \rightarrow 0$ as $|n| \rightarrow \infty$, where $f \in L^1(\mathbb{T}^d)$. Its quantitative version, the Lebesgue type estimate for the Fourier coefficients, is well known [Zy, Vol. I, Ch. 4, § 4] and given by

$$|\widehat{f}_n| \lesssim \omega_l \left(f, \frac{1}{|n|} \right)_1, \quad f \in L^1(\mathbb{T}^d), \quad (1.1)$$

where the modulus of smoothness $\omega_l(f, \delta)_p$ of a function $f \in L^p(X)$ is defined by

$$\omega_l(f, \delta)_p = \sup_{|h| \leq \delta} \|\Delta_h^l f(x)\|_{L^p(X)}, \quad 1 \leq p \leq \infty, \quad (1.2)$$

and

$$\Delta_h^l f(x) = \Delta_h^{l-1}(\Delta_h f(x)), \quad \Delta_h f(x) = f(x+h) - f(x).$$

For the Fourier transform, the estimate similar to (1.1) can be found in, e.g., [Tr1]

$$|\widehat{f}(\xi)| \lesssim \omega_l \left(f, \frac{1}{|\xi|} \right)_1, \quad f \in L^1(\mathbb{R}^d), \quad (1.3)$$

where the Fourier transform is given by

$$\widehat{f}(\xi) = \int_{\mathbb{R}^d} f(x) e^{i\xi x} dx, \quad \xi \in \mathbb{R}^d. \quad (1.4)$$

However, unlike (1.1) the inequality (1.3) cannot be extended for the range $p > 1$ (see section 7.2 below).

Very recently, Bray and Pinsky [BP1, BP2] and Ditzian [Di] (see also Gioev’s paper [Gi]) extended Lebesgue type estimate for the Fourier transform/coefficients. We will need the following

2000 *Mathematics Subject Classification.* 42B10, 26A15.

Key words and phrases. Fourier transforms, moduli of smoothness, Pitt’s inequality.

This research was partially supported by the MTM 2011-27637, RFFI 10-01-00564, RFFI 12-01-00170, 2009 SGR 1303. The results of the paper were presented at the conference on Function spaces in CRM (Barcelona) in September 2011.

average function. For a locally integrable function f the average on a sphere in \mathbb{R}^d of radius $t > 0$ is given by

$$V_t f(x) := \frac{1}{m_t} \int_{|y-x|=t} f(y) dy \quad \text{with} \quad V_t 1 = 1, \quad d \geq 2.$$

For $l \in \mathbb{N}$ we define

$$V_{l,t} f(x) := \frac{-2}{\binom{2l}{l}} \sum_{j=1}^l (-1)^j \binom{2l}{l-j} V_{j,t} f(x).$$

THEOREM A. *Let $f \in L^p(\mathbb{R}^d)$, $d \geq 2$, and $1 \leq p \leq 2$, $1/p + 1/p' = 1$. Then for $t > 0$, $l \in \mathbb{N}$,*

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{2l} |\widehat{f}(\xi)| \right]^{p'} d\xi \right)^{1/p'} \lesssim \|f - V_{l,t} f\|_p, \quad 1 < p \leq 2, \quad (1.5)$$

and

$$\sup_{\xi \in \mathbb{R}^d} \left[\min(1, t|\xi|)^{2l} |\widehat{f}(\xi)| \right] \lesssim \|f - V_{l,t} f\|_1. \quad (1.6)$$

Similar results were also proved for moduli of smoothness of functions on \mathbb{R} and \mathbb{T}^d (see [Di]). In the rest of the paper we will assume that $t > 0$, $l \in \mathbb{N}$, and

$$\Omega_l(f, t)_p = \|f - V_{l,t} f\|_p, \quad \theta = 2, \quad (1.7)$$

if $d \geq 2$ and

$$\Omega_l(f, t)_p = \omega_l(f, t)_p, \quad \theta = 1 \quad (1.8)$$

if $d = 1$.

The main goal of this paper is to extend inequalities (1.5) and (1.6) in the following sense. First, we prove sharper estimates by considering the weighted L^q norm of $\min(1, t|\xi|)^{\theta l} |\widehat{f}(\xi)|$, that is,

$$\left\| \min(1, t|\xi|)^{\theta l} |\widehat{f}(\xi)| \right\|_{L^q(u)} \lesssim \Omega_l(f, t)_p, \quad p \leq q \quad (1.9)$$

with the certain weight function u . Then varying the parameter q gives us the better bound from below of $\Omega_l(f, t)_p$. In particular, if $q = p'$ we arrive at (1.5) and (1.6).

Second, we prove the reverse inequalities showing how smoothness of a function depends on the average decay of its Fourier transform:

$$\Omega_l(f, t)_p \lesssim \left\| \min(1, t|\xi|)^{\theta l} |\widehat{f}(\xi)| \right\|_{L^q(u)}, \quad q \leq p, \quad (1.10)$$

Third, we define the class of general monotone functions and prove that for this class the equivalence result holds:

$$\Omega_l(f, t)_p \asymp \left\| \min(1, t|\xi|)^{\theta l} |\widehat{f}(\xi)| \right\|_{L^p(u)}. \quad (1.11)$$

Note that for $p = 2$, this follows from (1.9) and (1.10) in the general case (see also [BP1, Gi]).

The paper is organized as follows. In Section 2, we prove inequalities (1.9) and (1.10) when $1 < p \leq 2$ and $p \geq 2$ respectively. In Section 3 we study inequalities (1.9) and (1.10) in the case of radial functions and we show that, with a fixed p , the range of the parameter q is extended. In Section 4 we deal with the general monotone functions. Again, we prove inequalities (1.9) and (1.10) under wider range of the parameter q than in the case of radial functions. Moreover, we show equivalence (1.11) in this case. Section 5 studies inequalities (1.9) and (1.10) for functions on \mathbb{T}^d , $d \geq 1$. In Section 6 we obtain the equivalence result of type (1.11) for periodic functions whose sequence of Fourier coefficients is general monotone. Section 7 considers several application of obtained results in approximation theory (sharp relations between best approximations and moduli of smoothness) and functional analysis (embedding theorems, characterization of the Lipschitz/Besov spaces in terms of the Fourier transforms).

Finally, we remark that inequalities between moduli of smoothness and the Fourier transform in the Lebesgue and Lorentz spaces were studied earlier in [Cl] and [GK].

2. Growth of Fourier transforms via moduli of smoothness. The general case

The following theorem is the main result of this section.

THEOREM 2.1. *Let $f \in L^p(\mathbb{R}^d)$, $d \geq 1$.*

(A) *Let $1 < p \leq 2$. Then for $p \leq q \leq p'$ we have $|\xi|^{d(1-1/p-1/q)} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$, and*

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{\theta l} |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)| \right]^q d\xi \right)^{1/q} \lesssim \Omega_l(f, t)_p. \quad (2.1)$$

(B) *Let $2 \leq p < \infty$, $|\xi|^{d(1-1/p-1/q)} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$, $q > 1$, and $\max\{q, q'\} \leq p$. Then*

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{\theta l} |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)| \right]^q d\xi \right)^{1/q} \gtrsim \Omega_l(f, t)_p. \quad (2.2)$$

REMARK. Theorem A follows from Theorem 2.1 (A) (take $q = p'$). In part (B) we assume that for $f \in L^p(\mathbb{R}^d)$ the Fourier transform \widehat{f} is well defined and such that $|\xi|^{d(1-1/p-1/q)} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$ for a certain $q > 1$ satisfying $\max\{q, q'\} \leq p$.

PROOF OF THEOREM 2.1. We will use the following Pitt's inequality [**BH**] (see also [**GLT**]):

$$\left(\int_{\mathbb{R}^d} (|\xi|^{-\gamma} |\widehat{g}(\xi)|)^q d\xi \right)^{1/q} \lesssim \left(\int_{\mathbb{R}^d} (|x|^\beta |g(x)|)^p dx \right)^{1/p}, \quad (2.3)$$

where

$$\beta - \gamma = d \left(1 - \frac{1}{p} - \frac{1}{q} \right), \quad \max \left\{ 0, d \left(\frac{1}{p} + \frac{1}{q} - 1 \right) \right\} \leq \gamma < \frac{d}{q}, \quad 1 < p \leq q < \infty. \quad (2.4)$$

Here the Fourier transform \widehat{g} is understood in the usual sense of weighted Fourier inequality (2.3); see, e.g., [**BL**, Sect. 1, 2].

Let us write inequality (2.3) with change of parameters $\widehat{g} \leftrightarrow f$, $p \leftrightarrow q$, $\beta \leftrightarrow -\gamma$. Let $|\xi|^{-\gamma} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$, then

$$\left(\int_{\mathbb{R}^d} (|\xi|^{-\gamma} |\widehat{f}(\xi)|)^q d\xi \right)^{1/q} \gtrsim \left(\int_{\mathbb{R}^d} (|x|^\beta |f(x)|)^p dx \right)^{1/p}, \quad (2.5)$$

where

$$\beta - \gamma = d \left(1 - \frac{1}{p} - \frac{1}{q} \right), \quad \max \left\{ 0, d \left(\frac{1}{p} + \frac{1}{q} - 1 \right) \right\} \leq -\beta < \frac{d}{p}, \quad 1 < q \leq p < \infty. \quad (2.6)$$

The case of $d \geq 2$. Then by (1.7), $\Omega_l(f, t)_p = \|f - V_{l,t} f\|_p$, $\theta = 2$. Let us write the left-hand side in (2.1) and (2.2) as

$$I := \left\| \min(1, t|\xi|)^{2l} h(\xi) \right\|_q, \quad h(\xi) = |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)|.$$

In [**DD**, Cor. 2.3, Th. 3.1], it is shown that for $f \in L^p(\mathbb{R}^d)$, $1 \leq p \leq \infty$, $t > 0$, and integer l ,

$$\|f - V_{l,t} f\|_p \asymp K_l(f, \Delta, t^{2l})_p \asymp R_l(f, \Delta, t^{2l})_p, \quad (2.7)$$

where

$$K_l(f, \Delta, t^{2l})_p := \inf \left\{ \|f - g\|_p + t^{2l} \|\Delta^l g\|_p : \Delta^l g \in L^p(\mathbb{R}^d) \right\},$$

the Laplacian is given by $\Delta = \frac{\partial^2}{\partial x_1^2} + \cdots + \frac{\partial^2}{\partial x_d^2}$,

$$R_l(f, \Delta, t^{2l})_p := \|f - R_{\lambda,l,b}(f)\|_p + t^{2l} \|\Delta^l R_{\lambda,l,b}(f)\|_p, \quad (2.8)$$

$$\lambda = 1/t, \quad b \geq d + 2.$$

Here (see [**DD**, Sec. 2])

$$R_{\lambda,l,b}(f)(x) = (G_{\lambda,l,b} * f)(x), \quad G_{\lambda,l,b}(x) = \lambda^d G_{l,b}(\lambda x), \quad \widehat{G_{l,b}}(\xi) = \eta_{l,b}(|\xi|),$$

where

$$\eta_{l,b}(s) = (1 - s^{2l})_+^b, \quad s = |\xi| \geq 0, \quad (2.9)$$

and

$$\begin{aligned} [R_{\lambda,l,b}(f)]^\wedge(\xi) &= \eta_{l,b}(t|\xi) \widehat{f}(\xi), \quad [f - R_{\lambda,l,b}(f)]^\wedge(\xi) = [1 - \eta_{l,b}(t|\xi)] \widehat{f}(\xi), \\ [\Delta^l R_{\lambda,l,b}(f)]^\wedge(\xi) &= (-1)^l |\xi|^{2l} [R_{\lambda,l,b}(f)]^\wedge(\xi) = (-1)^l |\xi|^{2l} \eta_{l,b}(t|\xi) \widehat{f}(\xi). \end{aligned} \quad (2.10)$$

Also,

$$\|G_{\lambda,l,b}(x)\|_1 = \|G_{l,b}\|_1 < \infty. \quad (2.11)$$

Taking into account that, for $b > 0$,

$$\eta_{l,b}(s) \sim 1 - bs^{2l}, \quad s \rightarrow 0, \quad \eta_{l,b}(s) = 0, \quad s \geq 1,$$

we obtain

$$1 - \eta_{l,b}(s) \asymp \min(1, s)^{2l}, \quad s \geq 0. \quad (2.12)$$

Changing variables $b \leftrightarrow b + 1$ gives

$$\min(1, s)^{2l} \asymp 1 - \eta_{l,b+1}(s) = 1 - (1 - s^{2l})\eta_{l,b}(s) = 1 - \eta_{l,b}(s) + s^{2l}\eta_{l,b}(s).$$

Therefore,

$$I = \|\min(1, t|\xi|)^{2l} h(\xi)\|_q \asymp \|[1 - \eta_{l,b}(t|\xi|) + (t|\xi|)^{2l} \eta_{l,b}(t|\xi|)] h(\xi)\|_q. \quad (2.13)$$

Define

$$h_1(\xi) = [1 - \eta_{l,b}(t|\xi|)] h(\xi), \quad h_2(\xi) = (t|\xi|)^{2l} \eta_{l,b}(t|\xi|) h(\xi). \quad (2.14)$$

Note that both h_1 and h_2 are non-negative. For non-negative functions we have

$$\|h_1 + h_2\|_q \asymp \|h_1\|_q + \|h_2\|_q, \quad 1 \leq q \leq \infty. \quad (2.15)$$

This, (2.13), and (2.14) yield

$$I \asymp \left\| |\xi|^{d(1-1/p-1/q)} [1 - \eta_{l,b}(t|\xi|)] \widehat{f}(\xi) \right\|_q + \left\| |\xi|^{d(1-1/p-1/q)} (t|\xi|)^{2l} \eta_{l,b}(t|\xi|) \widehat{f}(\xi) \right\|_q,$$

or, by (2.10),

$$I \asymp \left\| |\xi|^{d(1-1/p-1/q)} [f - R_{\lambda,l,b}(f)]^\wedge(\xi) \right\|_q + t^{2l} \left\| |\xi|^{d(1-1/p-1/q)} [\Delta^l R_{\lambda,l,b}(f)]^\wedge(\xi) \right\|_q. \quad (2.16)$$

Now to prove (A), we assume that $p \leq q$ and we use (2.16) and Pitt's inequality (2.3) with $\beta = 0$. In this case $\gamma = d\left(\frac{1}{p} + \frac{1}{q} - 1\right)$ and $\gamma \geq 0$ (see (2.4)). The latter is ensured by $q \leq p'$. Then $|\xi|^{d(1-1/p-1/q)} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$ and

$$I \lesssim \|f - R_{\lambda,l,b}(f)\|_p + t^{2l} \|\Delta^l R_{\lambda,l,b}(f)\|_p.$$

Combining this with (2.7), and (2.8) we get (A).

In part (B) we assume that $q \leq p$. Inequality (2.2) follows from (2.16) and inequality (2.5) for $\beta = 0$. In this case, by (2.6), $\gamma = d\left(\frac{1}{p} + \frac{1}{q} - 1\right)$ and $\max\{0, \gamma\} \leq 0$, i.e., $\gamma \leq 0$. The latter is $q \geq p'$ or, equivalently, $q' \leq p$.

The case of $d = 1$. According to (1.8), we have $\Omega_l(f, t)_p = \omega_l(f, t)_p$ and $\theta = 1$. The proof of key steps is similar to the proof in the case of $d \geq 2$. The only difference is the realization result ([DHI]) given by

$$\omega_l(f, t)_p \asymp \inf \left(\|f - g\|_p + t^l \|g^{(l)}\|_p : g^{(l)} \in E_\lambda \cap L^p(\mathbb{R}) \right) \asymp \|f - g_\lambda\|_p + t^l \|g_\lambda^{(l)}\|_p, \quad \lambda = 1/t,$$

where E_λ is the collection of all entire functions of exponential type λ and $g_\lambda \in E_\lambda$ is such that

$$\|f - g_\lambda\|_p \lesssim E_\lambda(f)_p := \inf_{g \in E_\lambda} \|f - g\|_p.$$

Since $\|g_\lambda^{(l)}\|_p \asymp \|H g_\lambda^{(l)}\|_p$, $1 < p < \infty$, where H is the Hilbert transform [Tit, Ch. 5], then $\omega_l(f, t)_p \asymp \|f - g_\lambda\|_p + t^l \|D_l g_\lambda\|_p$, where $D_l = (id/dx)^l$ for even l and $D_l = -iH(id/dx)^l$ for odd l .

Let $\chi_\lambda := \chi_{[0, \lambda]}$. As Hille and Tamarkin [HT] showed, if $S_\lambda(f)$ is the partial Fourier integral of f , i.e.,

$$[S_\lambda(f)]^\wedge(\xi) = \chi_\lambda(|\xi|) \widehat{f}(\xi), \quad (2.17)$$

we have

$$\|S_\lambda(f)\|_p \lesssim \|f\|_p, \quad 1 < p < \infty.$$

Then (see also [Tim]) g_λ can be taken as $S_\lambda(f)$, that is, $\|f - S_\lambda(f)\|_p \lesssim E_\lambda(f)_p$. Therefore, for $1 < p < \infty$,

$$\omega_l(f, t)_p \asymp \|f - S_\lambda(f)\|_p + t^l \|S_\lambda^{(l)}(f)\|_p \asymp \|f - S_\lambda(f)\|_p + t^l \|D_l S_\lambda(f)\|_p, \quad (2.18)$$

where

$$[S_\lambda^{(l)}(f)]^\wedge(\xi) = (-i\xi)^l \chi_\lambda(|\xi|) \widehat{f}(\xi), \quad [D_l S_\lambda(f)]^\wedge(\xi) = |\xi|^l \chi_\lambda(|\xi|) \widehat{f}(\xi). \quad (2.19)$$

For $s \geq 0$ we have $\min(1, s)^l = 1 - \chi_1(s) + s^l \chi_1(s)$ and $\chi_1(ts) = \chi_\lambda(s)$, which gives

$$\min(1, ts)^l = 1 - \chi_\lambda(s) + (ts)^l \chi_\lambda(s). \quad (2.20)$$

This, (2.15), (2.17), and (2.19) imply

$$\begin{aligned} I &:= \left\| \min(1, t|\xi|)^l |\xi|^{1-1/p-1/q} |\widehat{f}(\xi)| \right\|_q = \left\| [1 - \chi_\lambda(|\xi|) + (t|\xi|)^l \chi_\lambda(|\xi|)] |\xi|^{1-1/p-1/q} |\widehat{f}(\xi)| \right\|_q \\ &\asymp \left\| |\xi|^{1-1/p-1/q} [1 - \chi_\lambda(|\xi|)] |\widehat{f}(\xi)| \right\|_q + \left\| |\xi|^{1-1/p-1/q} (t|\xi|)^l \chi_\lambda(|\xi|) |\widehat{f}(\xi)| \right\|_q \\ &= \left\| |\xi|^{1-1/p-1/q} [(f - S_\lambda(f))^\wedge(\xi)] \right\|_q + t^l \left\| |\xi|^{1-1/p-1/q} [D_l S_\lambda(f)]^\wedge(\xi) \right\|_q, \end{aligned} \quad (2.21)$$

which is an analogue of (2.16). Then as in the case of $d \geq 2$ we continue by using Pitt's inequality (2.3) and its corollary (2.5) with $\beta = 0$ and $d = 1$. This concludes the proof of the case $d = 1$. \square

3. Growth of Fourier transforms via moduli of smoothness. The case of radial functions

Theorem 2.1 was proved under the condition $1 < p \leq q \leq p' < \infty$ (A) and $1 < \max\{q, q'\} \leq p < \infty$ (B). When $d \geq 2$ these conditions can be extended if we restrict ourselves to radial functions

$$f(x) = f_0(|x|).$$

The Fourier transform of a radial function is also radial $\widehat{f}(\xi) = F_0(|\xi|)$ (see [SW, Ch. 4]) and it can be written as the Fourier–Hankel transform

$$F_0(s) = |S^{d-1}| \int_0^\infty f_0(t) j_{d/2-1}(st) t^{d-1} dt,$$

where $j_\alpha(t) = \Gamma(\alpha + 1)(t/2)^{-\alpha} J_\alpha(t)$ is the normalized Bessel function ($j_\alpha(0) = 1$), $\alpha \geq -1/2$. Useful properties of J_α can be found in, e.g., [AS, Ch. 9]; see also [GLT] for some properties of j_α .

THEOREM 3.1. *Let $f \in L^p(\mathbb{R}^d)$ be a radial function and $d \geq 2$.*

(A) *Let $1 < p \leq q < \infty$. Then, for $p \leq \frac{2d}{d+1}$, $q < \infty$ or $\frac{2d}{d+1} < p \leq 2$, $p \leq q \leq \left(\frac{d+1}{2} - \frac{d}{p}\right)^{-1}$,*

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{2l} |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)| \right]^q d\xi \right)^{1/q} \lesssim \|f - V_{l,t} f\|_p.$$

(B) *Let $2 \leq p < \infty$, $|\xi|^{d(1-1/p-1/q)} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$, $q > 1$ and $\max\left\{q, d \left(\frac{d+1}{2} - \frac{1}{q}\right)^{-1}\right\} \leq p$. Then*

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{2l} |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)| \right]^q d\xi \right)^{1/q} \gtrsim \|f - V_{l,t} f\|_p.$$

REMARK. 1. Formally, when $d = 1$ conditions in Theorems 3.1 and 2.1 coincide. However, note that no regularity condition was assumed in Theorem 2.1.

2. The range of conditions on p and q in Theorem 3.1 is wider than the corresponding range in Theorem 2.1 for $d \geq 2$.

Indeed, in Theorem 2.1 (A) we assume the following conditions: $1 < p \leq 2$ and $p \leq q \leq p'$. If $p \leq \frac{2d}{d+1}$, in Theorem 3.1 (A) conditions are $p \leq q < \infty$. If $\frac{2d}{d+1} < p \leq 2$, then $\left(\frac{d+1}{2} - \frac{d}{p}\right)^{-1} \geq p'$. Thus, the conditions $p \leq q \leq \left(\frac{d+1}{2} - \frac{d}{p}\right)^{-1}$ are less restrictive than $p \leq q \leq p'$.

In its turn, in Theorem 2.1 (B) we assume that $2 \leq p < \infty$ and $\max\{q, q'\} \leq p$. If $q < 2$, then $p \geq q'$ and $\max\left\{q, d\left(\frac{d+1}{2} - \frac{1}{q}\right)^{-1}\right\} = d\left(\frac{d+1}{2} - \frac{1}{q}\right)^{-1} < q'$. If $2 \leq q$, then $\max\left\{q, d\left(\frac{d+1}{2} - \frac{1}{q}\right)^{-1}\right\} = q$. Hence, we get $\max\left\{q, d\left(\frac{d+1}{2} - \frac{1}{q}\right)^{-1}\right\} \leq \max\{q, q'\}$.

PROOF OF THEOREM 3.1. The proof is similar to the proof of Theorem 2.1 but we use Pitt's inequality for radial functions. We also remark that for a radial function f , functions $f - R_{\lambda, l, b}(f)$ and $\Delta^l R_{\lambda, l, b}(f)$ are radial as well.

De Carli [DC] proved Pitt's inequality for the Hankel transform. In particular, this gives inequality (2.3) for radial functions. As it was shown in [DC], in this case the condition on γ is as follows

$$\frac{d}{q} - \frac{d+1}{2} + \max\left\{\frac{1}{p}, \frac{1}{q'}\right\} \leq \gamma < \frac{d}{q}, \quad 1 < p \leq q < \infty. \quad (3.1)$$

Therefore, (2.5) for radial functions holds under the condition

$$\frac{d}{p} - \frac{d+1}{2} + \max\left\{\frac{1}{q}, \frac{1}{p'}\right\} \leq -\beta < \frac{d}{p}, \quad 1 < q \leq p < \infty. \quad (3.2)$$

We will use (3.1) and (3.2) with $\beta = 0$ and $\gamma = d\left(\frac{1}{p} + \frac{1}{q} - 1\right)$.

To show (A), we assume (3.1), that is, the following two conditions hold simultaneously

$$\frac{d-1}{2} + \frac{1}{p} \leq \frac{d}{p}, \quad \frac{d-1}{2} + \frac{1}{q'} \leq \frac{d}{p}.$$

If $d \geq 2$, the first condition is equivalent to $p \leq 2$. If $p \leq \frac{2d}{d+1}$, then the second condition is $q < \infty$. If $\frac{2d}{d+1} < p \leq 2$, then respectively $q \leq \left(\frac{d+1}{2} - \frac{d}{p}\right)^{-1}$.

Let us verify all conditions in (B). We assume (3.2), or, equivalently,

$$\frac{d}{p} - \frac{d+1}{2} + \frac{1}{q} \leq 0, \quad \frac{d}{p} - \frac{d+1}{2} + \frac{1}{p'} \leq 0.$$

If $d \geq 2$, the second inequality is equivalent to the condition $p \geq 2$. The first inequality can be rewritten as $p \geq d\left(\frac{d+1}{2} - \frac{1}{q}\right)^{-1}$. Since also $p \geq q$, we finally arrive at condition $\max\left\{q, d\left(\frac{d+1}{2} - \frac{1}{q}\right)^{-1}\right\} \leq p$, under which needed Pitt's inequality holds. \square

4. Growth of Fourier transforms via moduli of smoothness. The case of general monotone functions

The following equivalence holds for $p = 2$ (see [BP1], [Di], [Gi] and Theorem 2.1 (A, B)):

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{\theta_l} |\widehat{f}(\xi)|\right]^p d\xi\right)^{1/p} \asymp \Omega_l(f, t)_p, \quad (4.1)$$

where $\Omega_l(f, t)_p$ and θ are given by (1.7) and (1.8).

In this section we show that similar two sided inequalities also hold for $\frac{2d}{d+1} < p < \infty$ provided \widehat{f} is radial, nonnegative and regular in a certain sense.

4.1. General monotone functions and the \widehat{GM}^d class. A function $\varphi(z)$, $z > 0$, is called *general monotone* ($\varphi \in GM$), if it is locally of bounded variation on $(0, \infty)$, vanishes at infinity, and for some constant $c > 1$ depending on φ , the following is true

$$\int_z^\infty |d\varphi(u)| \lesssim \int_{z/c}^\infty \frac{|\varphi(u)|}{u} du < \infty, \quad z > 0 \quad (4.2)$$

(see [GLT]). Any monotone function vanishing at infinity satisfies GM-condition. Note also that (4.2) implies

$$|\varphi(z)| \lesssim \int_{z/c}^\infty \frac{|\varphi(u)|}{u} du. \quad (4.3)$$

In particular, the latter gives, for any $b > 1$,

$$|\varphi(z)| \lesssim \int_{z/(bc)}^{\infty} u^{-1} \left(\int_{u/b}^{bu} \frac{|\varphi(v)|}{v} dv \right) du. \quad (4.4)$$

We will also use the following result on multipliers of general monotone functions.

LEMMA 4.1. *Let $\varphi \in GM$ and a function $\alpha(z)$ be locally of bounded variation on $(0, \infty)$ such that $\lim_{z \rightarrow 0} \alpha(z) = 0$ and*

$$\int_0^{cu} |d\alpha(v)| \lesssim |\alpha(u)|, \quad u > 0.$$

Then $\varphi_1 = \alpha\varphi \in GM$.

PROOF. By definition of GM, it is sufficient to verify

$$I := \int_z^{\infty} |d\varphi_1(u)| \lesssim \int_{z/c}^{\infty} \frac{|\varphi_1(u)|}{u} du, \quad z > 0. \quad (4.5)$$

First,

$$I \lesssim \int_z^{\infty} |\varphi(u)| |d\alpha(u)| + \int_z^{\infty} |\alpha(u)| |d\varphi(u)| =: I_1 + I_2,$$

and, by (4.3), we get

$$I_1 = \int_z^{\infty} |\varphi(u)| |d\alpha(u)| \lesssim \int_z^{\infty} \left(\int_{u/c}^{\infty} \frac{|\varphi(v)|}{v} dv \right) |d\alpha(u)| = \int_{z/c}^{\infty} \left(\int_z^{cu} |d\alpha(u)| \right) \frac{|\varphi(v)|}{v} dv.$$

To estimate I_2 , using

$$|\alpha(u)| = \left| \alpha(z) + \int_z^u d\alpha(v) \right| \lesssim |\alpha(z)| + \int_z^u |d\alpha(v)|, \quad u > z,$$

and condition (4.2), we have

$$\begin{aligned} I_2 &\lesssim |\alpha(z)| \int_z^{\infty} |d\varphi(u)| + \int_z^{\infty} \left(\int_z^u |d\alpha(v)| \right) |d\varphi(u)| \\ &\lesssim |\alpha(z)| \int_{z/c}^{\infty} \frac{|\varphi(v)|}{v} dv + \int_z^{\infty} \left(\int_v^{\infty} |d\varphi(u)| \right) |d\alpha(v)| \\ &\lesssim |\alpha(z)| \int_{z/c}^{\infty} \frac{|\varphi(v)|}{v} dv + \int_z^{\infty} \left(\int_{v/c}^{\infty} \frac{|\varphi(u)|}{u} du \right) |d\alpha(v)| \\ &= |\alpha(z)| \int_{z/c}^{\infty} \frac{|\varphi(v)|}{v} dv + \int_{z/c}^{\infty} \left(\int_z^{cu} |d\alpha(v)| \right) \frac{|\varphi(u)|}{u} du. \end{aligned}$$

Therefore, since

$$|\alpha(z)| = \left| \int_0^z d\alpha(v) \right| \leq \int_0^z |d\alpha(v)|,$$

we arrive at

$$I \lesssim I_1 + I_2 \lesssim \int_{z/c}^{\infty} \left(|\alpha(z)| + \int_z^{cu} |d\alpha(v)| \right) \frac{|\varphi(v)|}{v} dv \leq \int_{z/c}^{\infty} \left(\int_0^{cu} |d\alpha(v)| \right) \frac{|\varphi(v)|}{v} dv.$$

Finally, the integral condition on α concludes the proof of (4.5). \square

Let \widehat{GM}^d , $d \geq 1$, be the collection of all radial functions $f(x) = f_0(|x|)$, $x \in \mathbb{R}^d$, which are defined in terms of the inverse Fourier–Hankel transform

$$f_0(z) = \frac{|S^{d-1}|}{(2\pi)^d} \int_0^{\infty} F_0(s) j_{d/2-1}(zs) s^{d-1} ds, \quad (4.6)$$

where the function $F_0 \in GM$ and satisfies the following condition

$$\int_0^1 s^{d-1} |F_0(s)| ds + \int_1^{\infty} s^{(d-1)/2} |dF_0(s)| < \infty. \quad (4.7)$$

Applying Lemma 1 from the paper [GLT] to F_0 , we obtain that the integral in (4.6) converges in the improper sense and therefore $f_0(z)$ is continuous for $z > 0$. In addition, F_0 is a radial component of the Fourier transform of the function f , that is, $\widehat{f}(\xi) = F_0(|\xi|)$, $\xi \in \mathbb{R}^d$.

Let us give some examples of functions from the class \widehat{GM}^d .

EXAMPLE 1. Let $f \in C(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$, where $1 \leq p < 2d/(d+1)$ for $d \geq 2$ and $p = 1$ for $d = 1$, be a radial positive-definite function such that $F_0 \in GM$. Then $f \in \widehat{GM}^d$. Indeed, \widehat{f} is continuous function vanishing at infinity and $\widehat{f} \geq 0$ [SW, Ch. 1]. From continuity of f at zero we get $\widehat{f} \in L^1(\mathbb{R}^d)$ [SW, Cor. 1.26], i.e., $\int_0^\infty s^{d-1}|F_0(s)| ds < \infty$. Since any GM-function F_0 satisfies ([GLT, p. 111])

$$\int_1^\infty s^\sigma |dF_0(s)| \lesssim \int_{1/c}^\infty s^{\sigma-1} |F_0(s)| ds, \quad \sigma \geq 0,$$

then, using $(d-1)/2 - 1 < d-1$, we get

$$\int_0^1 s^{d-1} |F_0(s)| ds + \int_1^\infty s^{(d-1)/2} |dF_0(s)| \lesssim \int_0^\infty s^{d-1} |F_0(s)| ds < \infty.$$

Therefore, condition (4.7) holds, that is, $f \in \widehat{GM}^d$. As an example of such function we can take $f(x) = (1 + |x|^2)^{-(d+1)/2}$ and the corresponding $F_0(s) = c_d e^{-s}$.

EXAMPLE 2. Take $f(x) = j_{d/2}(|x|)$ (for $d = 1$, $f(x) = \frac{\sin x}{x}$). Then $F_0(s) = c\chi_1(s) \in GM$ and condition(4.7) holds, i.e., $f \in \widehat{GM}^d$. Moreover, we have (see, e.g., [GLT])

$$j_{d/2}(z) \asymp 1, \quad 0 \leq z \leq 1, \quad |j_{d/2}(z)| \lesssim z^{-(d+1)/2}, \quad z \geq 1,$$

and

$$|j_{d/2}(z)| \gtrsim z^{-(d+1)/2}, \quad z \in \bigcup_{k=1}^\infty [\rho_{d/2,k} + \varepsilon, \rho_{d/2,k+1} - \varepsilon],$$

where $\rho_{\alpha,k}$ positive zeros of the Bessel function J_α , $\inf_{k \geq 1} (\rho_{d/2,k+1} - \rho_{d/2,k}) \geq 3\varepsilon > 0$. This implies $f \in L^p(\mathbb{R}^d)$ if $p > \frac{2d}{d+1}$.

EXAMPLE 3. Let $F_0(s) \in GM$ and $|\xi|^{d(1-1/p-1/q)} F_0(|\xi|) \in L^q(\mathbb{R}^d)$, $1 < q \leq p < \infty$, $\frac{2d}{d+1} < p$. Then, using statement (A.1) below, condition (4.7) for F_0 holds, f is defined by (4.6), and $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$. The fact that $f \in L^p(\mathbb{R}^d)$ follows from Pitt's inequality (4.8) (take $\beta = 0$).

4.2. Two-sided inequalities.

THEOREM 4.1. Let $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$, $d \geq 1$.

(A) If $\widehat{f} \geq 0$ and $1 < p \leq q < \infty$, then

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{\theta l} |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)| \right]^q d\xi \right)^{1/q} \lesssim \Omega_l(f, t)_p.$$

(B) If $|\xi|^{d(1-1/p-1/q)} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$, $1 < q \leq p < \infty$, $\frac{2d}{d+1} < p$, then

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{\theta l} |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)| \right]^q d\xi \right)^{1/q} \gtrsim \Omega_l(f, t)_p.$$

REMARK. Conditions on p and q in Theorem 4.1 (A, B) are less restrictive than corresponding conditions in Theorem 3.1. It is clear for (A). Since $\frac{2d}{d+1} \leq d \left(\frac{d+1}{2} - \frac{1}{q} \right)^{-1}$, conditions $q \leq p$ and $\frac{2d}{d+1} < p$ in Theorem 4.1 (B) are weaker than $\max \left\{ 2, q, d \left(\frac{d+1}{2} - \frac{1}{q} \right)^{-1} \right\} \leq p$, which is the corresponding condition in Theorem 3.1 (B).

In case of $p = q$ Theorem 4.1 gives the following equivalence result.

COROLLARY 4.1. If $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$, $d \geq 1$, $\widehat{f} \geq 0$, $\frac{2d}{d+1} < p < \infty$, then

$$\left(\int_{\mathbb{R}^d} \left[\min(1, t|\xi|)^{\theta l} |\xi|^{d(1-2/p)} |\widehat{f}(\xi)| \right]^p d\xi \right)^{1/p} \asymp \Omega_l(f, t)_p.$$

EXAMPLE. Take $f(x) = j_{d/2}(|x|)$ (see Example 2). By Corollary 4.1, for $0 < t < 1$ and $\frac{2d}{d+1} < p < \infty$, we have

$$\Omega_t(f, t)_p \asymp \left\| \min(1, t|\xi|)^{\theta t} |\xi|^{d(1-2/p)} \chi_1(|\xi|) \right\|_p \asymp t^{\theta t}.$$

4.3. Weighted Fourier inequalities. To prove Theorem 4.1, we will use several auxiliary results from the paper [GLT].

Let $d \geq 1$, $1 < p, q < \infty$, $\beta - \gamma = d \left(1 - \frac{1}{p} - \frac{1}{q}\right)$, $g(x) = g_0(|x|)$, and $\widehat{g}(\xi) = G_0(|\xi|)$.

(A.1) If $g_0 \in GM$, $p \leq q$, and

$$\frac{d}{q} - \frac{d+1}{2} < \gamma < \frac{d}{q},$$

then the following Pitt's inequality holds [GLT, Th. 2 (A)]

$$\left\| |\xi|^{-\gamma} \widehat{g}(\xi) \right\|_q \lesssim \left\| |x|^\beta g(x) \right\|_p.$$

Then changing variables $g \leftrightarrow \widehat{f}$, $p \leftrightarrow q$, and $\beta \leftrightarrow -\gamma$, we get

$$\left\| |x|^\beta f(x) \right\|_p \lesssim \left\| |\xi|^{-\gamma} \widehat{f}(\xi) \right\|_q, \quad \frac{d}{p} - \frac{d+1}{2} < -\beta < \frac{d}{p}, \quad q \leq p. \quad (4.8)$$

Here $\widehat{f}(\xi) = F_0(|\xi|)$ and $F_0 \in GM$. Note [GLT, Sect. 5.1] that the condition $|\xi|^{-\gamma} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$ implies condition (4.7).

(A.2) Let $g_0 \in GM$, $g_0 \geq 0$ and g_0 satisfy condition (4.7). Then if $q \leq p$ and

$$\frac{d}{q} - \frac{d+1}{2} < \gamma,$$

then [GLT, Th. 2(B)]

$$\left\| |\xi|^{-\gamma} \widehat{g}(\xi) \right\|_q \gtrsim \left\| |x|^\beta g(x) \right\|_p.$$

Again, changing variables $g \leftrightarrow \widehat{f}$, $p \leftrightarrow q$, and $\beta \leftrightarrow -\gamma$, we arrive at

$$\left\| |x|^\beta f(x) \right\|_p \gtrsim \left\| |\xi|^{-\gamma} \widehat{f}(\xi) \right\|_q, \quad \frac{d}{p} - \frac{d+1}{2} < -\beta, \quad p \leq q. \quad (4.9)$$

Here $\widehat{f}(\xi) = F_0(|\xi|) \geq 0$ and $F_0 \in GM$.

From (A.1) and (A.2) (see also [GLT, Th. 1]), for a non-negative GM-function F_0 satisfying condition (4.7), we have

$$\left\| |\xi|^{d(1-2/p)} \widehat{f}(\xi) \right\|_p \asymp \|f(x)\|_p, \quad \frac{2d}{d+1} < p < \infty. \quad (4.10)$$

(A.3) Let $g_0 \geq 0$. For $z > 0$ we get (see [GLT, formula (53)])

$$\int_{z/(bc)}^\infty u^{-1} \left(\int_{u/b}^{bu} \frac{g_0(v)}{v} dv \right) du \lesssim \int_0^{2bc/z} u^{(d-1)/2-1} \left(\int_0^u v^{(d-1)/2} |G_0(v)| dv \right) du, \quad (4.11)$$

where $1 < b < \rho_{d/2,1}$.

(A.4) The following inequality was shown in [GLT, pp. 115-116]

$$\begin{aligned} & \left[\int_0^\infty u^{-\gamma p + dp/q - dp - 1} \left(\int_0^u v^{(d-1)/2-1} \left(\int_0^v z^{(d-1)/2} |G_0(z)| dz \right) dv \right)^p du \right]^{1/p} \\ & \lesssim \left(\int_{\mathbb{R}^d} [|x|^{-\gamma} |\widehat{g}(x)|]^q dx \right)^{1/q}, \quad \frac{d}{q} - \frac{d+1}{2} < \gamma, \quad q \leq p. \end{aligned}$$

Noting $u^{-\gamma p + dp/q - dp - 1} = u^{-p\beta - d - 1}$ and changing variables $\widehat{g} \leftrightarrow f$, $p \leftrightarrow q$, $\beta \leftrightarrow -\gamma$, we obtain

$$\begin{aligned} & \left[\int_0^\infty u^{q\gamma - d - 1} \left(\int_0^u v^{(d-1)/2-1} \left(\int_0^v z^{(d-1)/2} |f_0(z)| dz \right) dv \right)^q du \right]^{1/q} \\ & \lesssim \left(\int_{\mathbb{R}^d} [|x|^\beta |f(x)|]^p dx \right)^{1/p}, \quad \frac{d}{p} - \frac{d+1}{2} < -\beta, \quad p \leq q. \quad (4.12) \end{aligned}$$

4.4. Proof of Theorem 4.1 in the case $d \geq 2$. Let $t > 0$, $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$, $f(x) = f_0(|x|)$, and $\widehat{f}(\xi) = F_0(|\xi|)$. Note that $F_0 \in GM$. We use notations from the proof of Theorem 2.1.

First, we prove (B). Let $|\xi|^{d(1-1/p-1/q)} \widehat{f}(\xi) \in L^q(\mathbb{R}^d)$. We have

$$\begin{aligned} I &= \left\| \min(1, t|\xi|)^{2l} |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)| \right\|_q \\ &\asymp \left\| |\xi|^{d(1-1/p-1/q)} [1 - \eta_{l,b}(t|\xi|)] |\widehat{f}(\xi)| \right\|_q + \left\| |\xi|^{d(1-1/p-1/q)} (t|\xi|)^{2l} \eta_{l,b}(t|\xi|) |\widehat{f}(\xi)| \right\|_q =: I_1 + I_2. \end{aligned}$$

Then inequalities

$$\begin{aligned} \int_0^{cu} |d(1 - \eta_{l,b}(tv))| &\asymp t^{2l} \int_0^{cu} v^{2l-1} \eta_{l,b-1}(tv) dv \leq t^{2l} \int_0^{\min(cu, 1/t)} v^{2l-1} dv \\ &\asymp \min(1, ctu)^{2l} \asymp 1 - \eta_{l,b}(tu), \quad b > 1, \end{aligned}$$

and Lemma 4.1 imply that the function $[1 - \eta_{l,b}(ts)] F_0(s) = [1 - \eta_{l,b}(t|\xi|)] \widehat{f}(\xi)$ is a GM-function. Using Pitt's inequality (4.8) for $\beta = 0$ and $\gamma = d \left(\frac{1}{p} + \frac{1}{q} - 1 \right)$ yields

$$I_1 = \left\| |\xi|^{d(1-1/p-1/q)} [f - R_{\lambda,l,b}(f)]^\wedge(\xi) \right\|_q \gtrsim \|f - R_{\lambda,l,b}(f)\|_p \quad (4.13)$$

for

$$p > \frac{2d}{d+1}, \quad q \leq p. \quad (4.14)$$

Since $\eta_{l,b}(s) = 0$ when $s \geq 1$, then $(ts)^{2l} \eta_{l,b}(ts) = \min(1, ts)^{2l} \eta_{l,b}(ts)$. This and (2.10) give

$$(-1)^l t^{2l} [\Delta^l R_{\lambda,l,b}(f)]^\wedge(\xi) = \eta_{l,b}(ts) \min(1, ts)^{2l} F_0(s), \quad s = |\xi|.$$

Also, since $\eta_{l,b}(t|\xi|) = \widehat{G_{l,\lambda,b}}(\xi)$, then

$$(-1)^l t^{2l} \Delta^l R_{\lambda,l,b}(f) = G_{\lambda,l,b} * h, \quad \widehat{h}(\xi) = \min(1, t|\xi|)^{2l} F_0(|\xi|).$$

Using Young's convolution inequality, we obtain

$$\|t^{2l} \Delta^l R_{\lambda,l,b}(f)\|_p \leq \|G_{\lambda,l,b}\|_1 \|h\|_p = \|G_{l,b}\|_1 \|h\|_p \lesssim \|h\|_p.$$

We remark that

$$\min(1, ts)^{2l} F_0(s) \in GM. \quad (4.15)$$

This follows from the estimate

$$\int_0^{cu} |d \min(1, tv)^{2l}| \asymp t^{2l} \int_0^{\min(cu, 1/t)} v^{2l-1} dv \asymp \min[(ctu)^{2l}, 1] \asymp \min(1, tu)^{2l},$$

and Lemma 4.1.

Using again Pitt's inequality (4.8), we have

$$I = \left\| |\xi|^{d(1-1/p-1/q)} \widehat{h}(\xi) \right\|_q \gtrsim \|h\|_p \gtrsim \|t^{2l} \Delta^l R_{\lambda,l,b}(f)\|_p. \quad (4.16)$$

Adding estimates (4.13) and (4.16), we get

$$\|f - V_{l,t} f\|_p \asymp \|f - R_{\lambda,l,b}(f)\|_p + t^{2l} \|\Delta^l R_{\lambda,l,b}(f)\|_p \lesssim I_1 + I \lesssim I.$$

This and (4.14) give the part (B) of the theorem.

Let us now prove the part (A). If $p \leq \frac{2d}{d+1}$, the proof follows from Theorem 3.1. Suppose $\widehat{f}(\xi) = F_0(|\xi|) \geq 0$. By [DD, Lemma 3.4],

$$[f - V_{l,t} f]^\wedge(\xi) = [1 - m_l(t|\xi|)] \widehat{f}(\xi),$$

where the function $m_l(s)$ satisfies for $d \geq 2$ the following conditions

$$0 < C_1 s^{2l} \leq 1 - m_l(s) \leq C_2 s^{2l}, \quad 0 < s \leq \pi, \quad 0 < m_l(s) \leq v_{d,l} < 1, \quad s \geq \pi.$$

This gives

$$1 - m_l(s) \asymp \min(1, s)^{2l}, \quad s \geq 0. \quad (4.17)$$

Define $h(x) = f(x) - V_{l,t}f(x)$ and its radial component by $h_0 := G_0$. Using (4.11) for the non-negative function $g_0(s) = [1 - m_l(ts)]F_0(s)$, we obtain

$$J(z) := \int_{z/(bc)}^{\infty} u^{-1} \left(\int_{u/b}^{bu} \frac{g_0(v)}{v} dv \right) du \lesssim \int_0^{2bc/z} u^{(d-1)/2-1} \left(\int_0^u v^{(d-1)/2} |h_0(v)| dv \right) du. \quad (4.18)$$

Using (4.17), we get

$$J(z) \asymp \int_{z/(bc)}^{\infty} u^{-1} \left(\int_{u/b}^{bu} \frac{\min(1, tv)^{2l} F_0(v)}{v} dv \right) du, \quad z > 0.$$

where, by (4.15), $\min(1, tv)^{2l} F_0(v) \in GM$. Therefore, (4.4) for $z > 0$ yields

$$\min(1, tz)^{2l} F_0(z) \lesssim J(z).$$

Further, the latter and (4.18) imply

$$\begin{aligned} I &= \left\| \min(1, t|\xi|)^{2l} |\xi|^{d(1-1/p-1/q)} |\widehat{f}(\xi)| \right\|_q \asymp \left(\int_0^{\infty} \left[z^{d(1-1/p-1/q)} \min(1, tz)^{2l} F_0(z) \right]^q z^{d-1} dz \right)^{1/q} \\ &\lesssim \left(\int_0^{\infty} \left[z^{d(1-1/p-1/q)} J(z) \right]^q z^{d-1} dz \right)^{1/q} \\ &\lesssim \left(\int_0^{\infty} \left[z^{d(1-1/p-1/q)} \left(\int_0^{2bc/z} u^{(d-1)/2-1} \left(\int_0^u v^{(d-1)/2} |h_0(v)| dv \right) du \right) \right]^q z^{d-1} dz \right)^{1/q}. \end{aligned}$$

Changing variables $2bc/z \rightarrow z$, we obtain

$$I \lesssim \left(\int_0^{\infty} z^{-qd(1-1/p-1/q)-d-1} \left[\int_0^z u^{(d-1)/2-1} \left(\int_0^u v^{(d-1)/2} |h_0(v)| dv \right) du \right]^q dz \right)^{1/q}. \quad (4.19)$$

Let us now use (4.12) for $\beta = 0$ and $\gamma = d \left(\frac{1}{p} + \frac{1}{q} - 1 \right)$. Since, in this case

$$z^{-qd(1-1/p-1/q)-d-1} = z^{q\gamma-d-1}$$

inequalities (4.19) and (4.12) give

$$I \lesssim \left(\int_{\mathbb{R}^d} |h(x)|^p dx \right)^{1/p} = \|f - V_{l,t}f\|_p. \quad (4.20)$$

when $\frac{d}{p} - \frac{d+1}{2} < 0$ and $p \leq q$. The latter is $\frac{2d}{d+1} < p \leq q$. The proof of (A) is now complete. \square

4.5. Proof of Theorem 4.1 in the case $d = 1$. We follow the proof of Theorem 2.1. We have

$$\omega_l(f, t)_p \asymp \inf \left(\|f - g\|_p + t^l \|g^{(l)}\|_p : g^{(l)} \in E_\lambda \cap L^p(\mathbb{R}) \right), \quad \lambda = 1/t.$$

To show the estimate of $\omega_l(f, t)_p$ from above, that is, to prove (B), we take $g_\lambda(x)$ such that

$$\widehat{g}_\lambda(\xi) = [1 - (t|\xi|)^l]_+^b \widehat{f}(\xi), \quad b \geq 3.$$

Note that the function g_λ is analogues to the Riesz-type means $R_{\lambda,l,b}(f)$ and satisfies all required properties (2.9)–(2.12) with l in place of $2l$. In particular, $1 - [1 - (ts)^l]_+^b \asymp \min(1, ts)^l$. Proceeding similarly to the proof of (B) in the case $d \geq 2$, we arrive at the statement (B) in the case $d = 1$.

Let us now show (A). Let $\frac{2d}{d+1} < p \leq q < \infty$ and $\widehat{f} \geq 0$. Equivalence (2.18) gives

$$\omega_l(f, t)_p \asymp \|f - S_\lambda(f)\|_p + t^l \|D_l S_\lambda(f)\|_p \geq \|h\|_p,$$

where $h = f - S_\lambda(f) + t^l D_l S_\lambda(f)$. Moreover, $\widehat{h}(\xi) = [1 - \chi_\lambda(|\xi|) + (t|\xi|)^l \chi_\lambda(|\xi|)] \widehat{f}(\xi)$. Because of (2.20) and (4.15) with $s \geq 0$, we have $\widehat{h}(\xi) = \min(1, ts)^l F_0(s) \in GM$. Using then (4.9) with $\beta = 0$, we obtain

$$\omega_l(f, t)_p \gtrsim \|h\|_p \gtrsim \left\| |\xi|^{1-1/p-1/q} \widehat{h}(\xi) \right\|_p = \left\| \min(1, t|\xi|)^l |\xi|^{1-1/p-1/q} \widehat{f}(\xi) \right\|_p. \quad \square$$

5. Growth of Fourier coefficients via moduli of smoothness. The case of functions on \mathbb{T}^d

Let $f \in L^p(\mathbb{T}^d)$, $1 < p < \infty$, and

$$\widehat{f}_n = \int_{\mathbb{T}^d} f(x) e^{inx} dx, \quad n \in \mathbb{Z}^d, \quad \|\widehat{f}_n\|_{l^q(\mathbb{Z}^d)} := \left(\sum_{n \in \mathbb{Z}^d} |\widehat{f}_n|^q \right)^{1/q}.$$

In the paper [Di, Th. 4.1] the following was proved

$$\left\| \min(1, t|n|)^{\theta_l} |\widehat{f}_n| \right\|_{l^{p'}(\mathbb{Z}^d)} \lesssim \Omega_l(f, t)_p, \quad 1 < p \leq 2,$$

where $\Omega_l(f, t)_p$ is given by (1.7) and (1.8) with $\|\cdot\|_p = \|\cdot\|_{L^p(\mathbb{T}^d)}$.

The goal of the section is to obtain the generalization of this result which is a periodic analogue of inequalities (2.1)–(2.2).

THEOREM 5.1. *Let $f \in L^p(\mathbb{T}^d)$, $d \geq 1$, $1 < q < \infty$ and $\gamma = d\left(\frac{1}{p} + \frac{1}{q} - 1\right)$.*

(A) *Let $1 < p \leq 2$. Then for $p \leq q \leq p'$ we have $\{(1 + |n|)^{-\gamma} \widehat{f}_n\} \in l^q(\mathbb{Z}^d)$, and*

$$\left\| \min(1, t|n|)^{\theta_l} (1 + |n|)^{-\gamma} |\widehat{f}_n| \right\|_{l^q(\mathbb{Z}^d)} \lesssim \Omega_l(f, t)_p. \quad (5.1)$$

(B) *Let $2 \leq p < \infty$, $\{(1 + |n|)^{-\gamma} \widehat{f}_n\} \in l^q(\mathbb{Z}^d)$, and $\max\{q, q'\} \leq p$. Then*

$$\left\| \min(1, t|n|)^{\theta_l} (1 + |n|)^{-\gamma} |\widehat{f}_n| \right\|_{l^q(\mathbb{Z}^d)} \gtrsim \Omega_l(f, t)_p. \quad (5.2)$$

The proof of this theorem is similar to the proof of estimates (2.1)–(2.2) from Theorem 2.1. The key points are Pitt's inequalities of form

$$\left\| \widehat{f}_n (1 + |n|)^{-\gamma} \right\|_{l^q(\mathbb{Z}^d)} \lesssim \|f\|_{L^p(\mathbb{T}^d)}, \quad 1 < p \leq 2 \quad (5.3)$$

and

$$\left\| \widehat{f}_n (1 + |n|)^{-\gamma} \right\|_{l^q(\mathbb{Z}^d)} \gtrsim \|f\|_{L^p(\mathbb{T}^d)}, \quad p \geq 2, \quad (5.4)$$

under the corresponding conditions on q , as well as the realization results for the K -functionals in the periodic case (see [Di] and [DHI]).

PROOF OF (5.3). Let us show that the proof of (5.3) follows from Pitt's inequality for functions on \mathbb{R}^d . Note that $\gamma \geq 0$. Let f_* be the function on \mathbb{R}^d such that $f_* = f$ on $(-\pi, \pi]^d$ and $f_* = 0$ outside $(-\pi, \pi]^d$. Then

$$\|f_*\|_{L^p(\mathbb{R}^d)} = \|f\|_{L^p(\mathbb{T}^d)}, \quad \widehat{f}_*(\xi) = \int_{\mathbb{T}^d} f(x) e^{i\xi x} dx, \quad \xi \in \mathbb{R}^d, \quad \widehat{f}_*(n) = \widehat{f}_n, \quad n \in \mathbb{Z}^d.$$

Further, we use the results from [Ni, Ch. 3]. For an entire function g of exponential type σe , $\sigma > 0$, we have

$$\|g\|_{l^q(\mathbb{Z}^d)} \leq (1 + \sigma)^d \|g\|_{L^q(\mathbb{R}^d)}, \quad q \geq 1. \quad (5.5)$$

Note that the function \widehat{f}_* is an entire function of exponential type $\pi \bar{e}$, where $\bar{e} = (1, \dots, 1) \in \mathbb{R}^d$. We cannot use (5.5) since the weight function $|\xi|^{-\gamma}$, $\gamma \geq 0$, is not an entire function. However, it is possible to construct a positive radial entire function of exponential (spherical) type such that for $|\xi| \geq 1$ this function is equivalent to $|\xi|^{-\gamma}$.

We consider

$$\psi_\gamma(u) = j_\nu\left(\frac{u+i}{2}\right) j_\nu\left(\frac{u-i}{2}\right), \quad u \in \mathbb{C}, \quad 2\nu + 1 = \gamma \geq 0,$$

where j_ν is the normalized Bessel function. The function ψ_γ is an even positive entire function of type 1. Positivity of ψ_γ follows from the fact that all its zeros lie on lines $t \pm i$, $t \in \mathbb{R}$. The asymptotic expansion of Bessel functions [AS, formula 9.2.1] yields, for $|z| \rightarrow \infty$,

$$j_\nu(z) = \frac{C_\nu}{z^{\nu+1/2}} (\cos(z - c_\nu) + O(|z|^{-1})), \quad \operatorname{Re} z \geq 0, \quad |\operatorname{Im} z| \lesssim 1.$$

This and $\psi_\gamma(0) > 0$ give $\psi_\gamma(u) \asymp (1 + |u|)^{-\gamma}$, $u \in \mathbb{R}$.

Let us now consider the radial function $\psi_\gamma(|\xi|)$, $\xi \in \mathbb{R}^d$, which is an entire function of (spherical) type 1, and therefore, of type \bar{e} . Also,

$$\psi_\gamma(|\xi|) \asymp (1 + |\xi|)^{-\gamma}, \quad \xi \in \mathbb{R}^d. \quad (5.6)$$

Define $g(\xi) = \widehat{f}_*(\xi)\psi_\gamma(|\xi|)$, which is an entire function of type $(\pi + 1)\bar{e}$. Using (5.6), we get

$$\begin{aligned} \|g\|_{l^q(\mathbb{Z}^d)} &= \left(\sum_{n \in \mathbb{Z}^d} \left| \widehat{f}_*(n)\psi_\gamma(|n|) \right|^q \right)^{1/q} \asymp \left(\sum_{n \in \mathbb{Z}^d} \left| \widehat{f}_n(1 + |n|)^{-\gamma} \right|^q \right)^{1/q}, \\ \|g\|_{L^q(\mathbb{R}^d)} &= \left(\int_{\mathbb{R}^d} \left| \widehat{f}_*(\xi)\psi_\gamma(|\xi|) \right|^q d\xi \right)^{1/q} \lesssim \left(\int_{\mathbb{R}^d} \left| \widehat{f}_*(\xi)|\xi|^{-\gamma} \right|^q d\xi \right)^{1/q}. \end{aligned}$$

Then by (5.5) and Pitt's inequality for function on \mathbb{R}^d , we have

$$\begin{aligned} \left\| \widehat{f}_n(1 + |n|)^{-\gamma} \right\|_{l^q(\mathbb{Z}^d)} &\asymp \|g\|_{l^q(\mathbb{Z}^d)} \leq (\pi + 2)^d \|g\|_{L^q(\mathbb{R}^d)} \lesssim \left\| \widehat{f}_*(\xi)|\xi|^{-\gamma} \right\|_{L^q(\mathbb{R}^d)} \\ &\lesssim \|f_*\|_{L^p(\mathbb{R}^d)} = \|f\|_{L^p(\mathbb{T}^d)}. \end{aligned}$$

Thus we have proved the Pitt inequality (5.3) for function on \mathbb{T}^d . \square

PROOF OF (5.4). The following inequality is a consequence of [Nu, Th. 7] and Hardy's inequality for rearrangements:

$$\|f\|_{L^p} \lesssim \left(\sum_{k \in \mathbb{Z}^d} \prod_{j=1}^d (|k_j| + 1)^{q/p' - 1} |\widehat{f}_k|^q \right)^{1/q}, \quad \max\{q, q'\} \leq p. \quad (5.7)$$

The latter immediately gives (5.4). We would like to thank Erlan Nursultanov for drawing our attention to his result (5.7), which simplifies the proof. \square

6. An equivalence result for periodic functions

A complex null-sequence $a = \{a_n\}_{n \in \mathbb{N}}$ is said to be *general monotone*, written $a \in GM$, if (see [DT]) there exists $c > 1$ such that $(\Delta a_k = a_k - a_{k+1})$

$$\sum_{k=n}^{\infty} |\Delta a_k| \lesssim \sum_{k=\lfloor n/c \rfloor}^{\infty} \frac{|a_k|}{k}, \quad n \in \mathbb{N}.$$

THEOREM 6.1. *Let $f \in L^p(\mathbb{T})$, $1 < p < \infty$, and*

$$f(x) \sim \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx),$$

where nonnegative $\{a_n\}_{n \in \mathbb{N}}$, $\{b_n\}_{n \in \mathbb{N}}$ are general monotone sequences. Then

$$\omega_l(f, t)_p \asymp \left(\sum_{\nu=1}^{\infty} \min(1, \nu t)^{lp} \nu^{p-2} (a_\nu^p + b_\nu^p) \right)^{1/p}. \quad (6.1)$$

We will use the following lemma (see [AW]).

LEMMA 6.1. *Let $1 < p < \infty$ and let $\sum_{\nu=1}^{\infty} a_\nu \cos \nu x$ be the Fourier series of $f \in L^1(\mathbb{T})$.*

(A) *If the sequences $\{a_n\}$ and $\{\beta_n\}$ are such that*

$$\sum_{k=\nu}^{\infty} |\Delta a_k| \lesssim \beta_\nu, \quad \nu \in \mathbb{N}, \quad (6.2)$$

then

$$\|f\|_p^p \lesssim \sum_{\nu=1}^{\infty} \nu^{p-2} \beta_\nu^p. \quad (6.3)$$

(B) *If $a = \{a_n\}$ is a nonnegative sequence, then*

$$\sum_{n=1}^{\infty} \left(\sum_{k=\lfloor n/2 \rfloor}^n a_k \right)^p n^{-2} \lesssim \|f\|_p^p. \quad (6.4)$$

PROOF OF THEOREM 6.1. First, we remark that since $1 < p < \infty$ it is sufficient to prove that

$$\omega_l^p\left(f, \frac{1}{n}\right)_p \asymp I_1 + I_2,$$

where

$$I_1 = n^{-lp} \sum_{\nu=1}^n a_\nu^p \nu^{(l+1)p-2}, \quad I_2 = \sum_{\nu=n+1}^{\infty} a_\nu^p \nu^{p-2},$$

$$f(x) \sim \sum_{n=1}^{\infty} a_n \cos nx, \quad \{a_n\}_{n \in \mathbb{N}} \in GM.$$

We will also use the realization result for the modulus of smoothness (see [DHI]), that is,

$$\omega_l^p\left(f, \frac{1}{n}\right)_p \asymp \|f(x) - T_n(x)\|_p^p + n^{-lp} \|T_n^{(l)}(x)\|_p^p, \quad (6.5)$$

where $T_n(f)$ is the n -th almost best approximant, i.e., $\|f(x) - T_n(x)\|_p \lesssim E_n(f)_p$. In particular we can take T_n as $S_n = S_n(f)$, i.e., the n -th partial sum of $\sum_{k=1}^{\infty} a_k \cos kx$.

Let us prove estimate of I_1 and I_2 from above. Since $\{a_n\} \in GM$, we have

$$a_\nu \leq \sum_{l=\nu}^{\infty} |\Delta a_l| \lesssim \sum_{l=[\nu/c]}^{\infty} \frac{a_l}{l}, \quad (6.6)$$

then Hölder's inequality yields

$$\begin{aligned} I_1 &\lesssim n^{-lp} \sum_{\nu=1}^n \left(\sum_{j=[\nu/c]}^{\infty} \frac{a_j}{j} \right)^p \nu^{(l+1)p-2} \\ &\lesssim n^{-lp} \sum_{\nu=1}^n \left(\sum_{j=[\nu/c]}^n \frac{a_j}{j} \right)^p \nu^{(l+1)p-2} + n^{p-1} \left(\sum_{j=n}^{\infty} \frac{a_j}{j} \right)^p \\ &\lesssim n^{-lp} \sum_{\nu=1}^n \left(\sum_{j=\nu}^n \frac{a_j}{j} \right)^p \nu^{(l+1)p-2} + \sum_{j=n+1}^{\infty} a_j^p j^{p-2} =: I_3 + I_2. \end{aligned}$$

To estimate I_2 and I_3 , we are going to use the following inequalities

$$\sum_{s=n}^{\infty} a_s \lesssim \sum_{s=n}^{\infty} \frac{1}{s} \sum_{m=[s/2]}^s a_m \quad \text{and} \quad \sum_{s=1}^n a_s \lesssim \sum_{s=1}^{2n} \frac{1}{s} \sum_{m=[s/2]}^s a_m. \quad (6.7)$$

Then by Hardy's inequality [HLP], we have

$$\begin{aligned} I_3 &\lesssim n^{-lp} \sum_{\nu=1}^n \left(\sum_{j=\nu}^{2n} \frac{1}{j^2} \sum_{m=[j/2]}^l a_m \right)^p \nu^{(l+1)p-2} \\ &\lesssim n^{-lp} \sum_{j=1}^{2n} \left(\sum_{m=[j/2]}^j a_m \right)^p j^{lp-2}. \end{aligned}$$

Then Lemma 6.1 (B) and (6.5) yield

$$I_3 \lesssim n^{-lp} \left\| \sum_{\nu=1}^{2n} \nu^l a_\nu \cos \nu x \right\|_p^p \asymp n^{-lp} \|S_{2n}^{(l)}(f)\|_p^p \lesssim \omega_l^p\left(f, \frac{1}{2n}\right)_p \lesssim \omega_l^p\left(f, \frac{1}{n}\right)_p.$$

Further, using (6.6), (6.7), and Hardy' inequality, we have

$$\begin{aligned} I_2 &\lesssim \sum_{j=n+1}^{\infty} j^{p-2} \left(\sum_{s=[j/c]}^{\infty} \frac{a_s}{s} \right)^p \lesssim \sum_{j=n+1}^{\infty} j^{p-2} \left(\sum_{s=[j/c]}^{\infty} \frac{1}{s^2} \sum_{m=[s/2]}^s a_m \right)^p \\ &\lesssim \sum_{s=[n/c]}^{\infty} s^{-2} \left(\sum_{m=[s/2]}^s a_m \right)^p \lesssim \sum_{s=2n}^{\infty} s^{-2} \left(\sum_{m=[s/2]}^s a_m \right)^p + n^{-lp} \sum_{s=1}^{2n} s^{lp-2} \left(\sum_{m=[s/2]}^s a_m \right)^p. \end{aligned}$$

The last sum was estimated above. Again, by Lemma 6.1 (B) and (6.5),

$$\sum_{s=2n}^{\infty} s^{-2} \left(\sum_{m=\lfloor s/2 \rfloor}^s a_m \right)^p \lesssim \left\| \sum_{\nu=n}^{\infty} a_{\nu} \cos \nu x \right\|_p^p \lesssim \omega_l^p \left(f, \frac{1}{n} \right)_p.$$

So, we showed that

$$I_1 + I_2 \lesssim \omega_l^p \left(f, \frac{1}{n} \right)_p.$$

To prove the reverse, we use Lemma 6.1 (A), the definition of the GM class, Hölder's and Hardy's inequalities:

$$\begin{aligned} \|f - S_n\|_p^p &\lesssim \sum_{j=1}^{\infty} \beta_j^{p-2} j^{p-2} \lesssim n^{p-1} \left(\sum_{s=n}^{\infty} |\Delta a_s| \right)^p + \sum_{j=n}^{\infty} j^{p-2} \left(\sum_{s=l}^{\infty} |\Delta a_s| \right)^p \\ &\lesssim n^{p-1} \left(\sum_{s=\lfloor n/c \rfloor}^{\infty} \frac{a_s}{s} \right)^p + \sum_{j=n}^{\infty} j^{p-2} \left(\sum_{s=\lfloor j/c \rfloor}^{\infty} \frac{a_s}{s} \right)^p \lesssim \sum_{j=\lfloor n/c \rfloor}^{\infty} a_j^p j^{p-2} \lesssim I_1 + I_2, \end{aligned}$$

where $\beta_j = \sum_{s=\max(j,n)}^{\infty} |\Delta a_s|$. Similarly,

$$n^{-lp} \|S_n^{(l)}(f)\|_p^p \lesssim n^{-lp} \left\| \sum_{\nu=1}^n \nu^l a_{\nu} \cos \nu x \right\|_p^p \lesssim n^{-lp} \sum_{\nu=1}^n \nu^{p-2} \left(\sum_{s=\nu}^n |\Delta(s^l a_s)| \right)^p.$$

Further,

$$\sum_{s=\nu}^n |\Delta(s^l a_s)| \lesssim \sum_{s=\nu}^n s^{l-1} a_s + \sum_{s=\nu}^n s^l |\Delta a_s| \lesssim \sum_{s=\nu}^n s^{l-1} a_s + \sum_{s=\nu}^n |\Delta a_s| \left(\sum_{m=\nu}^s m^{l-1} + \nu^l \right),$$

and after routine calculations, we arrive at

$$\sum_{s=\nu}^n |\Delta(s^l a_s)| \lesssim \sum_{s=\lfloor \nu/c \rfloor}^n s^{l-1} a_s + n^l \sum_{m=n}^{\infty} \frac{a_m}{m}.$$

Using this and Hardy's inequality, we get $n^{-lp} \|S_n^{(l)}(f)\|_p^p \lesssim I_1 + I_2$. Finally, by (6.5),

$$\omega_l^p \left(f, \frac{1}{n} \right)_p \lesssim I_1 + I_2. \quad \square$$

7. Discussion and applications

7.1. Riemann–Lebesgue-type results. From Theorem A and [Di, Th. 2.2], one has the following estimate of the Fourier transform

$$t^{\theta l} \left(\int_{|\xi| < 1/t} |\xi|^{\theta l p'} |\widehat{f}(\xi)|^{p'} d\xi \right)^{1/p'} + \left(\int_{1/t \leq |\xi|} |\widehat{f}(\xi)|^{p'} d\xi \right)^{1/p'} \lesssim \Omega_l(f, t)_p, \quad 1 < p \leq 2. \quad (7.1)$$

On the other hand, Theorem 2.1 gives ($p \leq q \leq p'$, $1 < p \leq 2$)

$$t^{\theta l} \left(\int_{|\xi| < 1/t} |\xi|^{\theta l q + dq(1-1/p-1/q)} |\widehat{f}(\xi)|^q d\xi \right)^{1/q} + \left(\int_{1/t \leq |\xi|} |\xi|^{dq(1-1/p-1/q)} |\widehat{f}(\xi)|^q d\xi \right)^{1/q} \lesssim \Omega_l(f, t)_p. \quad (7.2)$$

If $q = p'$ (7.2) reduces to (7.1). The following example shows that (7.2), in general, provides better estimates than (7.1).

EXAMPLE. Let $\widehat{f}(\xi) = F_0(|\xi|)$,

$$F_0(s) = \frac{s^{-d/p'}}{\ln^{2/p}(2+s)}, \quad \frac{2d}{d+1} < p < \infty.$$

Note that F_0 is decreasing to zero and therefore $F_0 \in GM$. Also, it is easy to see that $|\xi|^{d(1-2/p)} \widehat{f}(\xi) \in L^p(\mathbb{R}^d)$. Hence, as in Example 3 (for $q = p$) we get $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$.

We have

$$t^{\theta l} \left(\int_{|\xi| < 1/t} |\xi|^{\theta l q + dq(1-1/p-1/q)} |\widehat{f}(\xi)|^q d\xi \right)^{1/q} + \left(\int_{|\xi| \geq 1/t} |\xi|^{dq(1-1/p-1/q)} |\widehat{f}(\xi)|^q d\xi \right)^{1/q} \\ \asymp [\ln(2 + 1/t)]^{-2/p+1/q}.$$

Then (7.1) gives

$$[\ln(2 + 1/t)]^{1-3/p} \lesssim \Omega_l(f, t)_p, \quad p \leq 2,$$

and (7.2) implies (with $q = p$)

$$[\ln(2 + 1/t)]^{-1/p} \lesssim \Omega_l(f, t)_p, \quad p \leq 2.$$

The latter estimate is stronger. Moreover, it is sharp since by Corollary 4.1 we in fact have

$$[\ln(2 + 1/t)]^{-1/p} \asymp \Omega_l(f, t)_p, \quad \frac{2d}{d+1} < p < \infty. \quad \square$$

7.2. Pointwise Riemann–Lebesgue-type results. For $f \in L^1(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$, $1 < p \leq 2$, the Riemann–Lebesgue inequality

$$|\widehat{f}(\xi)| \lesssim \Omega_l(f, 1/|\xi|)_p \quad (7.3)$$

does not hold in general.

Let us consider the case of $d = 1$ and $l \geq 2$. Define

$$f(x) = \sum_{n \in \mathbb{Z}} a_n \psi_n(x), \quad \psi_n(x) = \varepsilon_n \varphi(\varepsilon_n x) e^{-inx}, \\ \varphi(x) = (2\pi)^{-1} \left(\frac{\sin(x/2)}{x/2} \right)^2, \quad \widehat{\varphi}(\xi) = (1 - |\xi|)_+, \\ a_n = (1 + |n|)^{-3/2}, \quad \varepsilon_n = (1 + |n|)^{-\alpha p'}, \quad 1 < \alpha < 3/2.$$

Changing variables, we have

$$\|\varphi(\varepsilon_n x)\|_q = \varepsilon_n^{-1/q} \|\varphi\|_q \asymp \varepsilon_n^{-1/q}.$$

Hence

$$\|f\|_q \leq \sum_{n \in \mathbb{Z}} a_n \varepsilon_n \|\varphi(\varepsilon_n x)\|_q \asymp \sum_{n \in \mathbb{Z}} a_n \varepsilon_n^{1/q'} \leq \sum_{n \in \mathbb{Z}} (1 + |n|)^{-3/2} < \infty, \quad q \geq 1. \quad (7.4)$$

This implies $f \in L^1(\mathbb{R}) \cap L^p(\mathbb{R})$. The Fourier transform of f is written as

$$\widehat{f}(\xi) = \sum_{n \in \mathbb{Z}} a_n \widehat{\psi}_n(\xi), \quad \widehat{\psi}_n(\xi) = \widehat{\varphi} \left(\frac{\xi - n}{\varepsilon_n} \right).$$

Let us estimate $\omega_l(f, 1/t)_p$ from above. We will use the realization result (see (2.18)) given by

$$\omega_l(f, 1/t)_p \asymp \|f - S_t(f)\|_p + t^{-l} \|S_t^{(l)}(f)\|_p, \quad t > 0, \quad 1 < p < \infty. \quad (7.5)$$

Since $\text{supp } \widehat{\psi}_n \subset [n - \varepsilon_n, n + \varepsilon_n]$, then

$$S_t(f)(x) = \sum_{|n| \leq [t]} a_n \psi_n(x), \quad f(x) - S_t(f)(x) = \sum_{|n| > [t]} a_n \psi_n(x).$$

The function φ and its derivatives are given by

$$\varphi^{(l)}(x) = \frac{1}{2\pi} \int_{-1}^1 (1 - |\xi|) (-i\xi)^l e^{-i\xi x} d\xi, \quad l \in \mathbb{Z}_+.$$

Then $|\varphi^{(l)}(x)| \leq 1$, $x \in \mathbb{R}$. For $|x| \geq 1$ we get

$$|\varphi^{(l)}(x)| = \left| (2\pi)^{-1} \sum_{j=0}^l \binom{l}{j} [\sin^2(x/2)]^{(l-j)} [(x/2)^{-2}]^{(j)} \right| \lesssim \frac{1}{x^2}.$$

Thus, $|\varphi^{(l)}(x)| \lesssim (1 + x^2)^{-1}$ and then

$$|\psi_n^{(l)}(x)| = \varepsilon_n \left| \sum_{j=0}^l \binom{l}{j} [e^{-inx}]^{(l-j)} [\varphi(\varepsilon_n x)]^{(j)} \right| = \varepsilon_n \left| \sum_{j=0}^l \binom{l}{j} (-in)^{l-j} \varepsilon_n^j \varphi^{(j)}(\varepsilon_n x) \right|$$

$$\lesssim \frac{\varepsilon_n}{1 + (\varepsilon_n x)^2} \sum_{j=0}^l \binom{l}{j} |n|^{l-j} \varepsilon_n^j = \frac{\varepsilon_n (|n| + \varepsilon_n)^l}{1 + (\varepsilon_n x)^2}.$$

Then we arrive at

$$\|\psi_n^{(l)}\|_q \lesssim (|n| + \varepsilon_n)^l \varepsilon_n^{1/q'} \lesssim (1 + |n|)^l \varepsilon_n^{1/q'}.$$

Using these relations and proceeding similarly to (7.4), we get

$$\begin{aligned} \|f - S_t(f)\|_p &\leq \sum_{|n|>[t]} a_n \varepsilon_n^{1/p'} = \sum_{|n|>[t]} (1 + |n|)^{-3/2-\alpha} \lesssim t^{-1/2-\alpha}, \\ t^{-l} \|S_t^{(l)}(f)\|_p &\lesssim t^{-l} \sum_{|n|\leq[t]} a_n (1 + |n|)^l \varepsilon_n^{1/p'} \lesssim t^{-l} \sum_{|n|\leq[t]} (1 + |n|)^{l-3/2-\alpha}. \end{aligned}$$

For $l \geq 2$, $\alpha < 3/2$ we get $l - 3/2 - \alpha > -1$. Then $t^{-l} \|S_t^{(l)}(f)\|_p \lesssim t^{-1/2-\alpha}$. Finally, (7.5) implies

$$\omega_l(f, 1/t)_p \lesssim t^{-1/2-\alpha}.$$

For any large enough $t \in \mathbb{N}$

$$\widehat{f}(t) = a_t \widehat{\varphi}(0) = a_t \asymp t^{-3/2}.$$

Since $\varepsilon = \alpha - 1 > 0$, we finally get

$$\widehat{f}(t) \gtrsim t^\varepsilon \omega_l(f, 1/t)_p. \quad \square$$

However, let us remark that for functions from the class \widehat{GM}^d class, it is possible to obtain the pointwise bound of the Fourier transform.

COROLLARY 7.1. *Let $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$, $d \geq 1$, $\widehat{f}(\xi) = F_0(|\xi|) \geq 0$, and $\frac{2d}{d+1} < p < \infty$. Then*

$$F_0(t) \lesssim t^{-d/p'} \Omega_l(f, 1/t)_p. \quad (7.6)$$

PROOF. Since $f \in \widehat{GM}^d$, using (4.3) and Hölder's inequality, we get

$$\begin{aligned} F_0(s) &\lesssim \int_{s/c}^\infty \frac{F_0(u)}{u} du = \int_{s/c}^\infty F_0(u) u^{d-d/p-1/p} u^{-d+(d+1-p)/p} du \\ &\lesssim s^{d/p-d} \left(\int_{s/c}^\infty F_0^p(u) u^{dp-d-1} du \right)^{1/p}. \end{aligned} \quad (7.7)$$

Then using Corollary 4.1, we have

$$\Omega_l^p(f, t)_p \asymp t^{\theta l p} \int_0^{1/t} s^{\theta l p + dp - d - 1} F_0^p(s) ds + \int_{1/t}^\infty s^{dp - d - 1} F_0^p(s) ds \quad (7.8)$$

and by (7.7), we finally get

$$F_0(t) \lesssim t^{d/p-d} \left(\int_{t/c}^\infty F_0^p(u) u^{dp-d-1} du \right)^{1/p} \lesssim t^{-d/p'} \Omega_l(f, 1/t)_p. \quad \square$$

7.3. Moduli of smoothness and best approximations: sharp relations. The following direct and inverse theorems of trigonometric approximation are well known (see e.g. [DL, p. 210], [DDT, Intr.]):

$$\frac{1}{n^l} \left(\sum_{\nu=0}^n (\nu+1)^{\tau l - 1} E_\nu^\tau(f)_p \right)^{1/\tau} \lesssim \omega_l\left(f, \frac{1}{n}\right)_p \lesssim \frac{1}{n^l} \left(\sum_{\nu=0}^n (\nu+1)^{q l - 1} E_\nu^q(f)_p \right)^{1/q}, \quad (7.9)$$

where $f \in L^p(\mathbb{T})$, $1 < p < \infty$, $l, n \in \mathbb{N}$, $q = \min(2, p)$, $\tau = \max(2, p)$, $E_n(f)_p$ denotes the n -th best trigonometric approximation of f in L^p , and $\omega_l(f, \delta)_p$ is the L^p -modulus of smoothness, see (1.2) with $X = \mathbb{T}$.

We remark that (7.9) is the sharp version of classical Jackson and weak-type inequalities ([DL, p. 205, 208]) and it can be written equivalently as follows ([DDT]):

$$t^l \left(\int_t^1 u^{-\tau l - 1} \omega_{l+1}^\tau(f, u)_p du \right)^{1/\tau} \lesssim \omega_l(f, t)_p \lesssim t^l \left(\int_t^1 u^{-q l - 1} \omega_{l+1}^q(f, u)_p du \right)^{1/q}. \quad (7.10)$$

Constructing individual functions shows ([**DDT**]) that the parameters $q = \min(2, p)$ and $\tau = \max(2, p)$ are optimal in (7.9) and (7.10). For functions on $[-1, 1]$ inequalities of type (7.9) and (7.10) were obtained in [**To, DDT**].

For functions on $L^p(\mathbb{R}^d)$, similar results were also proved for $\Omega_k(f, t)_p$ and $E_n(f)_p$, i.e., the best L^p -approximation by functions of exponential type n (see [**DDT**]). For example, an analogue of (7.9) is given by

$$\frac{1}{2^{\theta l n}} \left(\sum_{\nu=0}^n 2^{\theta l \tau \nu} E_{2^\nu}^\tau(f)_p \right)^{1/\tau} \lesssim \Omega_l \left(f, \frac{1}{2^n} \right)_p \lesssim \frac{1}{2^{\theta l n}} \left(\sum_{\nu=0}^n 2^{\theta l q \nu} E_{2^\nu}^q(f)_p \right)^{1/q}, \quad \|\cdot\|_p = \|\cdot\|_{L^p(\mathbb{R}^d)}.$$

Below we show that for functions from the class \widehat{GM}^d we can completely solve the problem of description of relationships between $\Omega_l(f, t)_p$ and $E_n(f)_p$ as well as $\Omega_l(f, t)_p$ and $\Omega_{l+1}(f, t)_p$.

THEOREM 7.1. *If $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$, $d \geq 1$, $\widehat{f} \geq 0$, and $\frac{2d}{d+1} < p < \infty$, then*

$$\Omega_l(f, t)_p \asymp \left(t^{\theta l p} \int_t^1 u^{-\theta l p} \Omega_{l+1}^p(f, u)_p \frac{du}{u} \right)^{1/p} + t^{\theta l} A(f)_p, \quad 0 < t < \frac{1}{2}, \quad (7.11)$$

where $A(f)_p := \|\widehat{|\xi|^{\theta l + d(1-2/p)} \chi_1(|\xi|) \widehat{f}(\xi)}\|_p \lesssim \Omega_l(f, 1)_p$. In particular, we have

$$\left(t^{\theta l p} \int_t^1 u^{-\theta l p} \Omega_{l+1}^p(f, u)_p \frac{du}{u} \right)^{1/p} \lesssim \Omega_l(f, t)_p \lesssim \left(t^{\theta l p} \int_t^1 u^{-\theta l p} \Omega_{l+1}^p(f, u)_p \frac{du}{u} \right)^{1/p} + t^{\theta l} \|f\|_p$$

and

$$\frac{1}{2^{\theta l n}} \left(\sum_{\nu=0}^n 2^{\theta l p \nu} E_{2^\nu}^p(f)_p \right)^{1/p} \lesssim \Omega_l \left(f, \frac{1}{2^n} \right)_p \lesssim \frac{1}{2^{\theta l n}} \left(\sum_{\nu=0}^n 2^{\theta l p \nu} E_{2^\nu}^p(f)_p \right)^{1/p} + \frac{1}{2^{\theta l n}} \|f\|_p. \quad (7.12)$$

REMARK 1. In (7.11) one cannot drop $t^{\theta l} A(f)_p$. Indeed, consider

$$F_0^p(s) = s^{-(dp-d-1)} \chi_{1/n}(s).$$

Then

$$\Omega_l^p(f, t)_p \asymp t^{\theta l p} \int_0^{1/n} s^{\theta l p + dp - d - 1} F_0^p(s) ds \asymp t^{\theta l p} \int_0^{1/n} s^{\theta l p} ds \asymp t^{\theta l p} n^{-\theta l p - 1}.$$

Using this,

$$t^{\theta l p} \int_t^1 u^{-\theta l p} \Omega_{l+1}^p(f, u)_p \frac{du}{u} \asymp t^{\theta l p} \int_t^1 u^{\theta l p} n^{-\theta(l+1)p-1} \frac{du}{u} \asymp t^{\theta l p} n^{-\theta(l+1)p-1}.$$

Hence, writing

$$t^{\theta l} n^{-\theta l - 1/p} \lesssim \Omega_l(f, t)_p \lesssim t^{\theta l} \left(\int_t^1 u^{-\theta l p} \Omega_{l+1}^p(f, u)_p \frac{du}{u} \right)^{1/p} \lesssim n^{-\theta} t^{\theta l} n^{-\theta l - 1/p}$$

we arrive at a contradiction as $n \rightarrow \infty$.

PROOF OF THEOREM 7.1. Using Corollary 4.1, we get

$$\Omega_l^p(f, t)_p \asymp t^{\theta l p} \int_0^{1/t} s^{\theta l p + dp - d - 1} F_0^p(s) ds + \int_{1/t}^\infty s^{dp-d-1} F_0^p(s) ds =: J_1(t) + J_2(t)$$

and

$$\begin{aligned} t^{\theta l p} \int_t^1 u^{-\theta l p} \Omega_{l+1}^p(f, u)_p \frac{du}{u} &\asymp t^{\theta l p} \int_1^{1/t} u^{-\theta p - 1} \left[\int_0^u s^{\theta(l+1)p + dp - d - 1} F_0^p(s) ds \right] du \\ &\quad + t^{\theta l p} \int_1^{1/t} u^{\theta l p - 1} \left[\int_u^\infty s^{dp-d-1} F_0^p(s) ds \right] du =: I_1(t) + I_2(t). \end{aligned}$$

Then

$$I_1(t) = t^{\theta l p} \int_1^{1/t} u^{-\theta p - 1} \left[\left(\int_0^1 + \int_1^u \right) s^{\theta(l+1)p + dp - d - 1} F_0^p(s) ds \right] du$$

$$\begin{aligned} &\asymp t^{\theta lp} \int_0^1 s^{\theta(l+1)p+dp-d-1} F_0^p(s) ds + t^{\theta lp} \int_1^{1/t} s^{\theta(l+1)p+dp-d-1} F_0^p(s) \int_s^{1/t} u^{-\theta p-1} du ds \\ &\lesssim J_1(t) \end{aligned}$$

and

$$\begin{aligned} I_2(t) &= t^{\theta lp} \int_1^{1/t} u^{\theta lp-1} \left[\left(\int_u^{1/t} + \int_{1/t}^\infty \right) s^{dp-d-1} F_0^p(s) ds \right] du \\ &\asymp t^{\theta lp} \int_1^{1/t} s^{dp-d-1} F_0^p(s) \int_1^s u^{\theta lp-1} du ds + \int_{1/t}^\infty s^{dp-d-1} F_0^p(s) ds \\ &\lesssim J_1(t) + J_2(t). \end{aligned}$$

Using again Corollary 4.1,

$$\begin{aligned} A(f)_p &\asymp \left(\int_0^1 s^{\theta lp+dp-d-1} F_0^p(s) ds \right)^{1/p} \lesssim \left(\int_0^\infty s^{dp-d-1} \min(1, s)^{\theta lp} F_0^p(s) ds \right)^{1/p} \\ &\asymp \left\| \min(1, |\xi|)^{\theta l} |\xi|^{d(1-2/p)} \widehat{f}(\xi) \right\|_p \asymp \Omega_l(f, 1)_p. \end{aligned}$$

Moreover, $A^p(f)_p \lesssim J_1(t)$. Thus,

$$I_1(t) + I_2(t) + t^{\theta lp} A^p(f)_p \lesssim J_1(t) + J_2(t).$$

To prove the inverse inequality, we first remark that $s^{-\theta p} \lesssim \int_s^{1/t} u^{-\theta p-1} du$, $1 < s < 1/(2t)$ and therefore using (4.10),

$$\begin{aligned} J_1(2t) &\lesssim t^{\theta lp} \int_0^1 s^{\theta lp+dp-d-1} F_0^p(s) ds + t^{\theta lp} \int_1^{1/2t} s^{\theta(l+1)p+dp-d-1} F_0^p(s) \left(\int_s^{1/t} u^{-\theta p-1} du \right) ds \\ &\lesssim t^{\theta lp} A^p(f)_p + I_1(t). \end{aligned}$$

Also,

$$\begin{aligned} J_2(2t) &\lesssim \int_{1/(2t)}^\infty s^{dp-d-1} F_0^p(s) ds \\ &\lesssim \int_{1/(2t)}^{1/t} s^{dp-d-1} F_0^p(s) ds + t^{\theta lp} \int_{1/(2t)}^{1/t} u^{\theta lp-1} \int_u^\infty s^{dp-d-1} F_0^p(s) ds du \\ &\lesssim I_1(t) + I_2(t). \end{aligned}$$

Finally, to verify (7.12), we apply [DDT, (5.7) and (5.8)]. □

Using (6.1), we state the analogous result for periodic functions; compare with (7.9) and (7.10).

THEOREM 7.2. *Let $f \in L^p(\mathbb{T})$, $1 < p < \infty$, and*

$$f(x) \sim \sum_{n=1}^\infty (a_n \cos nx + b_n \sin nx),$$

where non-negative $\{a_n\}_{n \in \mathbb{N}}$, $\{b_n\}_{n \in \mathbb{N}}$ are general monotone sequences. Then

$$\omega_l(f, t)_p \asymp \left(t^{lp} \int_t^1 u^{-lp} \omega_{l+1}^p(f, u)_p \frac{du}{u} \right)^{1/p}, \quad 0 < t < \frac{1}{2}.$$

In particular,

$$\omega_l(f, 1/n)_p \asymp \left(n^{-lp} \sum_{\nu=0}^n (\nu+1)^{lp-1} E_\nu^p(f)_p \right)^{1/p},$$

where $E_\nu(f)_p$ is the best L^p -approximation of f by trigonometric polynomials of degree ν .

Note that similar equivalence results for continuous functions were obtained in [Tik, Ths. 5.1, 5.2].

7.4. A characterization of the Besov spaces. For $1 \leq p \leq \infty$ and $\tau, r > 0$, define the Besov space $B_{p,\tau}^r(\mathbb{R}^d)$ as the collection of functions $f \in L^p(\mathbb{R}^d)$ such that

$$\|f\|_{B_{p,\tau}^r(\mathbb{R}^d)} = \|f\|_{L^p(\mathbb{R}^d)} + \left(\int_0^1 \left(\frac{\Omega_l(f,t)_p}{t^r} \right)^\tau \frac{dt}{t} \right)^{1/\tau} < \infty,$$

where $0 < r < \theta l$. Similarly we define the Lipschitz space $\text{Lip}_p^r(\mathbb{R}^d) \equiv B_{p,\infty}^r(\mathbb{R}^d)$, i.e.,

$$\|f\|_{\text{Lip}_p^r(\mathbb{R}^d)} = \|f\|_{L^p(\mathbb{R}^d)} + \sup_t \frac{\Omega_l(f,t)_p}{t^r}, \quad 0 < r < \theta l.$$

It turns out that it is possible to characterize functions from the Besov space $B_{p,\tau}^r(\mathbb{R}^d)$ in terms of growth properties of their Fourier transforms.

THEOREM 7.3. *Let $d \geq 1$, $1 < \tau \leq \infty$, and $\frac{2d}{d+1} < p \leq \tau$. If $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$ and $\widehat{f} \geq 0$, then a necessary and sufficient condition for $f \in B_{p,\tau}^r(\mathbb{R}^d)$ is*

$$\int_0^\infty s^{r\tau+d\tau-d\tau/p-1} F_0^\tau(s) ds < \infty \quad \text{if } 1 < \tau < \infty \quad (7.13)$$

and

$$\sup_s s^{r+d-d/p} F_0(s) < \infty \quad \text{if } \tau = \infty. \quad (7.14)$$

PROOF. *The case of $1 < \tau < \infty$.* Let first (7.13) hold. By (7.8), we get

$$\begin{aligned} |f|_{B_{p,\tau}^r} &\asymp K_1 + K_2 + K_3 := \int_0^1 t^{(\theta l - r)\tau - 1} \left(\int_0^1 s^{\theta l p + d p - d - 1} F_0^p(s) ds \right)^{\tau/p} dt \\ &+ \int_1^\infty t^{(r - \theta l)\tau - 1} \left(\int_1^t s^{\theta l p + d p - d - 1} F_0^p(s) ds \right)^{\tau/p} dt + \int_0^1 t^{r\tau - 1} \left(\int_{1/t}^\infty s^{d p - d - 1} F_0^p(s) ds \right)^{\tau/p} dt. \end{aligned}$$

Then by Hölder's inequality with parameters $\alpha = \tau/p$ and α' , we get

$$K_1 \lesssim \int_0^1 s^{r\tau+d\tau-d\tau/p-1} F_0^\tau(s) ds.$$

By Hardy's inequalities (see e.g. [BSH, p. 124]), we have

$$K_2 + K_3 \lesssim \int_1^\infty s^{r\tau+d\tau-d\tau/p-1} F_0^\tau(s) ds.$$

Hence, if (7.13) holds, $f \in B_{p,\tau}^r(\mathbb{R}^d)$.

Let $f \in B_{p,\tau}^r(\mathbb{R}^d)$. By (7.7),

$$F_0(s)^\tau \lesssim s^{d\tau/p-d\tau} \left(\int_{s/c}^\infty F_0^p(u) u^{d p - d - 1} du \right)^{\tau/p}.$$

Therefore, making use of this, we have

$$\begin{aligned} \int_0^\infty s^{r\tau+d\tau-d\tau/p-1} F_0^\tau(s) ds &\lesssim \int_1^\infty s^{r\tau-1} \left(\int_s^\infty F_0^p(u) u^{d p - d - 1} du \right)^{\tau/p} ds \\ &+ \int_0^1 s^{r\tau-1} \left(\int_s^\infty F_0^p(u) u^{d p - d - 1} du \right)^{\tau/p} ds \lesssim |f|_{B_{p,\tau}^r} + \left\| |\xi|^{d(1-2/p)} \widehat{f}(\xi) \right\|_p^\tau \int_0^1 s^{r\tau-1} ds. \end{aligned}$$

Finally, since $\left\| |\xi|^{d(1-2/p)} \widehat{f}(\xi) \right\|_p \lesssim \|f\|_p$ (see (4.10)), (7.13) holds.

The case of $\tau = \infty$. Let first (7.14) hold. Then by (7.8), (7.14) yields

$$\Omega_l^p(f,t)_p \lesssim t^{\theta l p} \int_0^{1/t} s^{\theta l p - r p - 1} ds + \int_{1/t}^\infty s^{-r p - 1} ds \lesssim t^{r p},$$

i.e., $f \in \text{Lip}_p^r(\mathbb{R}^d)$.

On the other hand, if $f \in \text{Lip}_p^r(\mathbb{R}^d)$, we use (7.7) and (7.8)

$$F_0^p(s) \lesssim s^{d-dp} \int_{s/c}^\infty F_0^p(u) u^{dp-d-1} du \lesssim s^{d-dp} \Omega_l^p(f, 1/s)_p \lesssim s^{d-dp-rp},$$

which is (7.14). □

7.5. Embedding theorems. The following Sobolev-type embedding result for the Besov space with the limiting smoothness parameter is well known: $B_{p,q}^r \hookrightarrow L^q$, $r = d(\frac{1}{p} - \frac{1}{q})$ (see, e.g., [Pe, (8.2)]). Theorem 7.3 gives the sharpness of this result in the following sense.

COROLLARY 7.2. *Let $d \geq 1$ and $\frac{2d}{d+1} < p < q < \infty$. If $f \in \widehat{GM}^d \cap L^p(\mathbb{R}^d)$ and $\widehat{f} \geq 0$, then*

$$f \in B_{p,q}^r(\mathbb{R}^d), \quad r = d\left(\frac{1}{p} - \frac{1}{q}\right) \iff f \in L^q(\mathbb{R}^d). \tag{7.15}$$

PROOF. To show (7.15), we combine Theorem 7.3 and $\|\|\xi\|^{d(1-2/p)} \widehat{f}(\xi)\|_p \asymp \|f\|_p$, $\frac{2d}{d+1} < p < \infty$ (see (4.10)). □

Note that the embedding $B_{p,q}^r \hookrightarrow L^q$ is equivalent to the sharp (Ul’yanov) inequalities for moduli of smoothness in different metrics, as recently shown in [Tr2, Th. 2.4].

References

[AS] M. Abramowitz, I. A. Stegun, eds., *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, New York: Dover Publications, 1972.

[AW] R. Askey, S. Wainger, *Integrability theorems for Fourier series*, Duke Math. J., **33** (1966), 223–228.

[BH] J. J. Benedetto, H. P. Heinig, *Weighted Fourier Inequalities: New Proofs and Generalizations*, J. Fourier Anal. Appl., **9** (2003), 1–37.

[BL] J. J. Benedetto, J. D. Lakey, *The Definition of the Fourier Transform for Weighted Inequalities*, J. Funct. Anal., **120**, no. 2 (1994), 403–439.

[BSh] C. Bennett, R. Sharpley, *Interpolation of operators*, Academic Press, 1988.

[BP1] W. Bray, M. Pinsky, *Growth properties of Fourier transforms via moduli of continuity*, J. Funct. Anal., **255**, no. 9 (2008), 2265–2285.

[BP2] W. Bray, M. Pinsky, *Growth properties of Fourier transforms*, arXiv:0910.1115

[Cl] D. B. H. Cline, *Regularly varying rates of decrease for moduli of continuity and Fourier transforms of functions on \mathbb{R}^d* , J. Math. Anal. Appl. **159** (1991), 507–519.

[DD] F. Dai, Z. Ditzian, *Combinations of multivariate averages*, J. Approx. Theory, **131**, no. 2 (2004), 268–283.

[DDT] F. Dai, Z. Ditzian, S. Tikhonov, *Sharp Jackson inequality*, J. Approx. Theory, **151**, no. 1 (2008), 86–112.

[DC] L. De Carli, *On the L^p - L^q norm of the Hankel transform and related operators*, J. Math. Anal. Appl., **348**, no. 1 (2008), 366–382.

[DL] R. A. DeVore, G. G. Lorentz, *Constructive approximation*, Berlin: Springer-Verlag, 1993.

[Di] Z. Ditzian, *Smoothness of a function and the growth of its Fourier transform or its Fourier coefficients*, J. Approx. Theory, **162**, no. 5 (2010), 980–986.

[DHI] Z. Ditzian, V. H. Hristov, K. G. Ivanov, *Moduli of smoothness and K -functionals in L_p , $0 < p < 1$* , Constr. Approx., **11**, no. 1 (1995), 67–83.

[DT] M. I. Dyachenko, S. Tikhonov, *Convergence of trigonometric series with general monotone coefficients*, C.R. Acad. Sci. Paris, Ser. I, **345** (2007), 123–126.

[GK] J. García-Cuerva, V. Kolyada, *Rearrangement estimates for Fourier transforms in L^p and H^p in terms of moduli of continuity*, Math. Nachr., **228** (2001), 123–144.

[Gi] D. Goev, *Moduli of continuity and average decay of Fourier transforms: Two-sided estimates*, in: J. Baik, T. Kriecherbauer, L. Li, K. D. McLaughlin, C. Tomei (Eds.), *Integrable Systems and Random Matrices: In Honor of Percy Deift*, in: Contemp. Math., vol. 458, 377–392, Amer. Math. Soc., 2008.

[GLT] D. Gorbachev, E. Liflyand, S. Tikhonov, *Weighted Fourier inequalities: Boas’ conjecture in \mathbb{R}^n* , J. d’Analyse Math., **114** (2011), 99–120.

[HLP] G. H. Hardy, J. E. Littlewood, G. Pólya, *Inequalities*, 2nd ed. Cambridge University Press, 1952.

[HT] E. Hille, J. D. Tamarkin, *On the theory of Fourier transforms*, Bull. Amer. Math. Soc, **39** (1933), 768–774.

[Ni] S. M. Nikolskii, *Approximation of functions of several variables and imbedding theorems*, Berlin; Heidelberg; New York: Springer, 1975.

[Nu] E. D. Nursultanov, *On the coefficients of multiple Fourier series in L_p -spaces*, Izv. RAN Ser. Mat., **64**, (1) (2000), 95–122.

[Pe] J. Peetre, *Espaces d’interpolation et théorème de Soboleff*, Ann. Inst. Fourier (Grenoble), **16** (1966), 279–317.

[SW] E. M. Stein, G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton, N. J., 1971.

[Tik] S. Tikhonov, *Best approximation and moduli of smoothness: computation and equivalence theorems*, J. Approx. Th., **153** (2008), 19–39.

- [Tim] M. F. Timan, *Best approximation and modulus of smoothness of functions prescribed on the entire real axis*, Izv. Vyssh. Uchebn. Zaved. Mat., no. 6 (1961), 108–120 (in Russian).
- [Tit] E. Titchmarsh, *Introduction to the theory of Fourier integrals*, 2nd ed., Clarendon Press, Oxford University, 1948.
- [To] V. Totik, *Sharp converse theorem of L^p polynomial approximation*, Constr. Approx., **4** (1988), 419–433.
- [Tr1] W. Trebels, *Estimates for moduli of continuity of functions given by their Fourier transform*, Lecture Notes in Math., vol. 571, 277–288, Springer, Berlin, 1977.
- [Tr2] W. Trebels, *Inequalities for moduli of smoothness versus embeddings of function spaces*, Arch. Math. 94 (2010), 155–164.
- [Zy] A. Zygmund, *Trigonometric series*, vol. I, II, 3th ed., Cambridge, 2002.

D. GORBACHEV

TULA STATE

300600 TULA, RUSSIA

E-mail address: `dvgmail@mail.ru`

S. TIKHONOV

ICREA AND CENTRE DE RECERCA MATEMÀTICA

08193 BELLATERRA, BARCELONA, SPAIN

E-mail address: `stikhonov@crm.cat`