

PARAMETRIZATION OF SPECTRAL SURFACES OF A CLASS OF PERIODIC 5-DIAGONAL MATRICES

IONELA MOALE AND PETER YUDITSKII

ABSTRACT. The main result of this work is a parametric description of the spectral surfaces of a class of periodic 5-diagonal matrices, related to the strong moment problem. This class is a self-adjoint twin of the class of CMV matrices. Jointly they form the simplest possible classes of 5-diagonal matrices.

1. INTRODUCTION

The main result of this work is a parametric description of the spectral surfaces of a class of periodic 5-diagonal matrices. This class is a self-adjoint twin of the quite popular now class of CMV matrices [5, 6]. Jointly they form the simplest possible classes of 5-diagonal matrices, we explain this claim in details in Section 2. We take a risk to introduce an abbreviation and call them SMP (Strong Moment Problem) matrices, see e.g. [3] and the remark below.

In a generic case SMP matrix is defined as a ratio of two 3-diagonal matrices $A = A_2 A_1^{-1}$, where

$$A_1 = \begin{bmatrix} \ddots & & & & & \\ & \ddots & \bar{\pi}_{-1} & & & \\ & & \ddots & \sigma_{-1} & 0 & \\ & & & \pi_0 & 1 & \bar{\pi}_1 \\ & & & & 0 & \sigma_1 & 0 \\ & & & & & \pi_2 & 1 & \ddots \\ & & & & & & 0 & \ddots \\ & & & & & & & \ddots \end{bmatrix}, \quad A_2 = \begin{bmatrix} \ddots & & & & & \\ & \ddots & 0 & & & \\ & & \ddots & 1 & \bar{p}_0 & \\ & & & 0 & q_0 & 0 \\ & & & & p_1 & 1 & \bar{p}_2 \\ & & & & & 0 & q_2 & \ddots \\ & & & & & & p_3 & \ddots \\ & & & & & & & \ddots \end{bmatrix},$$

A_1 and A_2 are invertible, equivalently $\sigma_{2n-1} \neq 0$ and $q_{2n} \neq 0$ for all n . Due to the symmetry $A = A^*$, the generating coefficient sequences are subject to the restrictions

$$\frac{p_{2n-1}}{q_{2n-2}} = -\frac{\pi_{2n-1}}{\sigma_{2n-1}}, \quad \frac{p_{2n}}{q_{2n}} = -\frac{\pi_{2n}}{\sigma_{2n-1}}.$$

2010 *Mathematics Subject Classification.* 30E05, 30F15, 47B39, 46E22.

Key words and phrases. Strong Moment Problem, periodic CMV matrices, Hardy spaces on Riemann surfaces, conformal mappings, comb domains, reproducing kernels.

This work was supported by Austrian Science Fund FWF, project no: P22025-N18.

Note that A_1^{-1} and A_2^{-1} are also 3-diagonal, thus A and A^{-1} are 5-diagonal matrices represented as products of 3-diagonal matrices. Note also that degenerations are possible, see Remark 4.3.

Let $\{e_n\}$ be the standard basis in the two-sided ℓ^2 . According to the above definition

$$\begin{aligned} Ae_{2n} &= A_2 e_{2n} = \bar{p}_{2n} e_{2n-1} + q_{2n} e_{2n} + p_{2n+1} e_{2n+1} \\ A^{-1} e_{2n-1} &= A_1 e_{2n-1} = \bar{\pi}_{2n-1} e_{2n-2} + \sigma_{2n-1} e_{2n-1} + \pi_{2n} e_{2n} \end{aligned}$$

Thus, constructing A we follow the procedure, which is similar to the CMV matrices case: having e_{-1} and e_0 as the generators of the cyclic subspace we form the whole space applying A on the even step and A^{-1} on the odd step, however, as it was mentioned, the operator is not unitary, but self-adjoint, that is, the spectrum is not on the unit circle but on the real axis. From this point of view the situation is similar to the orthogonalization procedure in the strong moment problem construction [3], however because of periodicity we are interested in two-sided matrices; it is very essential: we do not assume that the operator A is positive!

A can be represented as a *two dimensional perturbation* of a block orthogonal matrix

$$A = \begin{bmatrix} A_- & 0 \\ 0 & A_+ \end{bmatrix} + e_{-1} \langle \cdot, \tilde{e}_0 \rangle \tilde{p}_0 + \tilde{e}_0 \langle \cdot, e_{-1} \rangle \tilde{p}_0,$$

where

$$\tilde{p}_0 = \|P_+ A e_{-1}\|, \quad \tilde{e}_0 = \frac{1}{\tilde{p}_0} P_+ A e_{-1},$$

$A_{\pm} = P_{\pm} A P_{\pm}$ are restrictions of A to the positive and negative half-axis according to the orthogonal decomposition $\ell^2 = \ell_-^2 \oplus \ell_+^2$. For this reason certain general facts from its spectral theory can be reduced to the spectral theory of Jacobi matrices. However it is quite different as soon as we pose the problem:

Problem 1.1. Describe the spectral sets of periodic SMP matrices.

Recall that for periodic Jacobi matrices the spectrum is a system of intervals, which possesses the following parametric description, see [1] and references therein. For a system of nonnegative parameters $\{h_k\}_{k=1}^{n-1}$, let $D = D(h_1, \dots, h_{n-1})$ be the region obtained from the half-strip



$$\{w : -\pi n < \operatorname{Re} w < 0, \operatorname{Im} w > 0\}$$

by removing vertical intervals

$$\{w : \operatorname{Re} w = -\pi k, 0 < \operatorname{Im} w \leq h_k\}, \quad k = 1, \dots, n-1.$$

Let θ be the conformal map of the upper half-plane \mathbb{H} to D normalized by the conditions $\theta(a_0) = 0$, $\theta(b_0) = -\pi n$, $\theta(\infty) = \infty$. Denote by $E(h_1, \dots, h_{n-1})$ the full preimage of the interval $[-\pi n, 0] \subset \partial D$, i.e.:

$$E(h_1, \dots, h_{n-1}) := \theta^{-1}([-\pi n, 0]).$$

FIGURE 1. Π region for $\omega_0 \neq \pi\ell$ FIGURE 2. Π region for $\omega_0 = \pi\ell_0$.

A system of intervals $E = [b_0, a_0] \setminus \cup_{j \geq 1} (a_j, b_j)$ is the spectrum of a periodic Jacobi matrix if and only if $E = E(h_1, \dots, h_{n-1})$ for a certain system of parameters $\{h_k\}_{k=1}^{n-1}$.

The spectral sets for periodic CMV matrices are given by conformal mappings onto similar *periodic comb-like domains*. To formulate our main result we define comb regions of a new kind.

For integers k and $m \in [0, k]$ and parameters $h_0 > 0$ and ω_0 , $0 \leq \omega_0 \leq \pi m$ consider the region $\Pi_k^m(h_0, \omega_0)$ bounded by the hyperbolic curves

$$(1.1) \quad (\operatorname{Im} u)^2 < (\operatorname{Re} u)^2 + h_0^2,$$

and the orthogonal systems of hyperbolas

$$(1.2) \quad \begin{aligned} \operatorname{Im} u &< \frac{\pi m - \omega_0}{2 \operatorname{Re} u}, \quad \operatorname{Im} u > \frac{\pi(m - k) - \omega_0}{2 \operatorname{Re} u}, \quad \text{for } \operatorname{Re} u > 0, \\ \operatorname{Im} u &< \frac{-\omega_0}{2 \operatorname{Re} u}, \quad \operatorname{Im} u > \frac{\pi(k) - \omega_0}{2 \operatorname{Re} u}, \quad \text{for } \operatorname{Re} u < 0. \end{aligned}$$

If $\omega_0 \neq \pi\ell$, $0 \leq \ell \leq m$, by Π we denote the region which is obtained from $\Pi_k^m(h_0, \omega_0)$ by removing pieces of hyperbolic curves

$$(1.3) \quad \operatorname{Im} u = \frac{\pi\ell - \omega_0}{2\operatorname{Re} u}, \quad \text{for } \operatorname{Im} u > 0, \quad 1 \leq \ell \leq m-1$$

and

$$(1.4) \quad \operatorname{Im} u = \frac{\pi(\ell - k) - \omega_0}{2\operatorname{Re} u}, \quad \text{for } \operatorname{Im} u < 0, \quad m+1 \leq \ell \leq 2k-1$$

of length h_ℓ , $h_\ell \geq 0$, see Fig. 1. If $\omega_0 = \pi\ell_0$ then the hyperbolic curves related to $\ell = \ell_0$ in (1.3) and $\ell = k + \ell_0$ in (1.4) degenerate. In this case the corresponding cuts are pieces of the imaginary axis, as soon as

$$(1.5) \quad h_{\ell_0} < h_0 \text{ and } h_{k+\ell_0} < h_0.$$

Otherwise one of them still satisfies (1.5), and another one has T-shape, see Fig. 2, consisting of the piece of the imaginary axis

$$(1.6) \quad 0 \leq \operatorname{Im} u \leq h_0 \text{ or } -h_0 \leq \operatorname{Im} u \leq 0,$$

respectively, and of the real interval

$$(1.7) \quad -h_-^{(\ell_0)} \leq \operatorname{Re} u \leq h_+^{(\ell_0)}, \quad h_\pm^{(\ell_0)} \geq 0.$$

Theorem 1.2. *For a region Π described by the conditions (1.1)–(1.7), let $\theta : \mathbb{H} \rightarrow \Pi$ be a conformal map such that 0 and ∞ correspond to the infinite points in Π . A system of intervals E , $0, \infty \notin E$, is the spectral set of a periodic matrix of SMP class if and only if it corresponds to the preimage of the part of the boundary given by the condition (1.1) for a certain Π .*

The structure of the work is as follows: the simplest possible spectral surfaces of periodic 5-diagonal matrices are discussed in Section 2. In Section 3 we prove our main theorem. The functional model for periodic SMP matrices is given in Section 4.

The second author is thankful to the organizers of the program *Hilbert spaces of entire functions and spectral theory of self-adjoint differential operators*, at CRM, Barcelona, 2011, and to Alex Eremenko for stimulating discussions. In a sense this paper is an addition to their joint work [1].

2. SPECTRAL SURFACES WITH THE MAXIMAL NUMBER OF BOUNDARY OVALS

Let J be a 5-diagonal self-adjoint matrix of period d

$$(2.1) \quad J = rS^2 + pS + q + S^{-1}\bar{p} + S^{-2}r,$$

where S is the shift operator and p, r, q are diagonal matrices of period d , such that $r_m > 0$. We recall certain fundamental facts from the spectral theory of multi-diagonal periodic matrices [2], adopting to the 5-diagonal case.

For

$$(2.2) \quad j(w) = \begin{bmatrix} q_0 & \bar{p}_1 & r_2 & \dots & 0 & r_0/w & p_0/w \\ p_1 & q_1 & \bar{p}_2 & r_3 & \dots & 0 & r_1/w \\ r_2 & p_2 & q_2 & \bar{p}_3 & r_4 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & r_{d-3} & p_{d-3} & q_{d-3} & \bar{p}_{d-2} & r_{d-1} \\ r_0 w & 0 & \dots & r_{d-2} & p_{d-2} & q_{d-2} & \bar{p}_{d-1} \\ \bar{p}_0 w & r_1 w & 0 & \dots & r_{d-1} & p_{d-1} & q_{d-1} \end{bmatrix}$$

let

$$F(z, w) = \frac{\det\{j(w) - z \cdot I\}}{\prod_{j=0}^{d-1} r_j} = w^2 + 1/w^2 + A(z)w + A_*(z)1/w + B(z)$$

where A and B are polynomials, in particular, for even $d = 2k$

$$(2.3) \quad \begin{aligned} B(z) &= \frac{z^{2k}}{\prod_{j=0}^{2k-1} r_j} + \dots, \\ A_*(z) &:= \overline{A(\bar{z})} = \left(\frac{-1}{\prod_{j=0}^{k-1} r_{2j}} + \frac{-1}{\prod_{j=0}^{k-1} r_{2j+1}} \right) z^k + \dots \end{aligned}$$

Then the spectral curve corresponding to J is of the form

$$(2.4) \quad \mathcal{R} = \{P = (z, w) : F(z, w) = 0\}.$$

\mathcal{R} is endowed with an antiholomorphic involution $\tau P := (\bar{z}, 1/\bar{w})$ for which

$$\mathcal{R} \setminus \partial\mathcal{R}_+ = \mathcal{R}_+ \cup \mathcal{R}_-, \quad \mathcal{R}_+ = \{P = (z, w) \in \mathcal{R} : |w| < 1\},$$

see Fig. 3. Note that the spectrum of J (as the operator acting in ℓ^2) corresponds to the fixed line of the involution τ , $\tau P = P$,

$$z \in \sigma(J) \Leftrightarrow \exists w : P = (z, w) \in \partial\mathcal{R}_+.$$

In other words it is described by the condition $|w| = 1$.

Recall that the spectral surfaces related to Jacobi matrices are of the form

$$\tilde{\mathcal{R}} = \left\{ (w, z) : w + \frac{1}{w} = \tilde{A}(z) \right\},$$

where \tilde{A} is a real polynomial. In the similar decomposition $\tilde{\mathcal{R}} \setminus \partial\tilde{\mathcal{R}} = \tilde{\mathcal{R}}_+ \cup \tilde{\mathcal{R}}_-$ it possesses the following property: the number of boundary ovals, i.e., the number of intervals

$$\partial\tilde{\mathcal{R}} = \{z \in \mathbb{R} : |\tilde{A}(z)| \leq 2\}$$

is maximal for the given genus of the surface.

We say that the spectral curve \mathcal{R} , related to a 5-diagonal matrix, is of the simplest structure if it has *maximal possible number of components* of the boundary





FIGURE 3. Topology of the spectral curve

FIGURE 4. \mathcal{R}_+ as the two sheeted covering of z -plane

$\partial\mathcal{R}_+$ for the given genus. For example, in Fig. 3 the number of boundary components is 3, but its genus is 4 and the maximal possible number of components is 5. That is, the curve of this structure does not belong to the class.

In other words, let us represent \mathcal{R}_+ as two a sheeted covering of the z -plane, see Fig. 4. It is a hyperelliptic curve with a system of cuts $\partial\mathcal{R}_+$. We say that the spectral curve is of the simplest structure if this hyperelliptic curve has genus 0, i.e.:

$$\mathcal{R}_+ \simeq \bar{\mathbb{C}} \setminus E.$$

The corresponding equivalence can be written explicitly

$$(2.5) \quad z = \lambda + \frac{c^2}{\lambda - \lambda_0}, \quad \lambda \in \mathbb{C}, \quad c_1 = \lambda_0 - 2c, \quad c_2 = \lambda_0 + 2c,$$

where c_1, c_2 denote the only two possible critical values of z , in the case that both numbers are finite and if, say, $c_2 = \infty$ then

$$z = \lambda^2 - 2c, \quad c_1 = c^2 - 2c.$$

The set E , which corresponds to $\partial\mathcal{R}_+$, is a system of cuts in the complex plane \mathbb{C} with the property

$$(2.6) \quad z(\lambda) \in \mathbb{R} \text{ for } \lambda \in E.$$

It is essential to note that E is far from being an arbitrary system of cuts for which (2.6) holds. Recall that up to now the second function w was not involved into considerations. Meanwhile $w = w(\lambda)$ is a function in $\bar{\mathbb{C}} \setminus E$ with the following properties [2]:

- (i) w is single-valued and holomorphic,
- (ii) $|w| < 1$ in $\bar{\mathbb{C}} \setminus E$ and $|w| = 1$ on E ,
- (iii) zeros of w are $\{\lambda_0, \infty\} = z^{-1}(\infty) \not\subset E$ (of equal multiplicity).

For definiteness, here and below, we consider the case (2.5) (with two finite critical values). The properties (i)-(iii) imply that

$$(2.7) \quad \frac{1}{k} \log \frac{1}{|w(\lambda)|} = G_{\lambda_0}(\lambda) + G_{\infty}(\lambda),$$

where $G_{\lambda_0}(\lambda)$ is the Green function in the domain $\bar{\mathbb{C}} \setminus E$ with a logarithmic pole at λ_0 and k is the multiplicity of w in λ_0 and ∞ respectively.

Let us recall the concept of the complex Green function $b_{\lambda_0}(\lambda)$. It is an analytic multivalued function in $\bar{\mathbb{C}} \setminus E$ such that

$$(2.8) \quad \log \frac{1}{|b_{\lambda_0}(\lambda)|} = G_{\lambda_0}(\lambda).$$

Note that (2.8) determines b_{λ_0} up to a unimodular constant. In what follows we assume the normalizations $b_{\lambda_0}(\infty) > 0$ and $b_{\infty}(\lambda_0) > 0$.

Let $\pi_1(\bar{\mathbb{C}} \setminus E)$ be the fundamental group of this domain. Then b_{λ_0} generates the character $\mu_{\lambda_0} \in \pi_1(\bar{\mathbb{C}} \setminus E)^*$ on this group by

$$b_{\lambda_0} \circ \gamma = \mu_{\lambda_0}(\gamma) b_{\lambda_0}, \quad \gamma \in \pi_1(\bar{\mathbb{C}} \setminus E),$$

which indicates the multivalued structure of the complex Green function. Moreover, let us split E into connected components, $E = \cup_{j=0}^{\kappa} E_j$, and let γ_j 's be simple contours around E_j 's. Note that they form generators of the group $\pi_1(\bar{\mathbb{C}} \setminus E)$ subject to the condition

$$\gamma_0 \circ \cdots \circ \gamma_m = \text{trivial}.$$

Then, for a suitable choice of the direction of γ_j ,

$$(2.9) \quad \mu_{\lambda_0}(\gamma_j) = e^{2\pi i \omega_{\lambda_0}(E_j)},$$

where $\omega_{\lambda_0}(E_j)$ is the harmonic measure of E_j at λ_0 .

The factor $b_{\lambda_0} b_\infty$ removes the singularities of z in $\bar{\mathbb{C}} \setminus E$. Let

$$t_\infty = \frac{(b_\infty b_{\lambda_0} z)(\infty)}{|(b_\infty b_{\lambda_0} z)(\infty)|}, \quad t_{\lambda_0} = \frac{(b_\infty b_{\lambda_0} z)(\lambda_0)}{|(b_\infty b_{\lambda_0} z)(\lambda_0)|},$$

Define $\xi \in [0, 1)$ by the condition $t_\infty = e^{2\pi i \xi} t_{\lambda_0}$. Let $\mu = \mu_{\lambda_0} \mu_\infty$. Thus $\mu(\gamma_j) = e^{2\pi i \omega_j}$, where $\omega_j = \omega_{\lambda_0}(E_j) + \omega_\infty(E_j)$.

Theorem 2.1. *Let z be given by (2.5). Let $E = \cup_{j=0}^{\kappa} E_j \subset z^{-1}(\mathbb{R})$ be a system of cuts (2.6). Then $\mathcal{R} = \mathcal{R}_+ \cup \partial \mathcal{R}_+ \cup \mathcal{R}_-$, where $\mathcal{R}_+ \simeq \bar{\mathbb{C}} \setminus E$, is a spectral curve of a 5-diagonal periodic matrix if and only if the numbers ξ and ω_j (for all j) are rational. Moreover $w = t_\infty^{-k} (b_{\lambda_0} b_\infty)^k$, where k is a common denominator of these rational numbers.*

A proof is based on a fact of the general theory [2], that a given periodic J with the spectral curve \mathcal{R} possesses functional representation as the multiplication operator by z . We give such a representation following basically [4, 7].

For a fixed character α the multivalued analytic functions F , $F \circ \gamma = \alpha(\gamma)F$, such that $|F(\lambda)|^2$ has a harmonic majorant in $\bar{\mathbb{C}} \setminus E$, form the Hardy space $H^2(\alpha) \subset L^2_{\omega_\infty}$ with the norm given by the integral of the boundary values:

$$\|F\|^2 = \int_E |F(\lambda)|^2 \omega_\infty(d\lambda).$$

Note that the point-evaluation functional is bounded in this space and therefore in $H^2(\alpha)$ there is the reproducing kernel k_λ^α :

$$F(\lambda) = \langle F, k_\lambda^\alpha \rangle, \quad \lambda \in \bar{\mathbb{C}} \setminus E,$$

for all $F \in H^2(\alpha)$.

Lemma 2.2. *For a character $\alpha \in \pi_1(\bar{\mathbb{C}} \setminus E)^*$ let*

$$K_\lambda^\alpha = \frac{k_\lambda^\alpha}{\|k_\lambda^\alpha\|}$$

denote the normalized reproducing kernel at λ . Then, for an arbitrary system of unimodular constants t_m , the family

$$(2.10) \quad e_{2n} = t_{2n} b_{\lambda_0}^n b_\infty^n K_{\lambda_0}^{\alpha \mu_{\lambda_0}^{-n} \mu_\infty^{-n}}, \quad e_{2n+1} = t_{2n+1} b_{\lambda_0}^{n+1} b_\infty^n K_\infty^{\alpha \mu_{\lambda_0}^{-n-1} \mu_\infty^{-n}}$$

forms an orthonormal basis in $H^2(\alpha)$, $n \geq 0$. Moreover, extended to negative indexes it forms an orthonormal basis in $L^2_{\omega_\infty}$.

Proof. The system is orthonormal. Every function from $H^2(\alpha)$ orthogonal to it has a zero of infinite multiplicity in λ_0 (and ∞) and therefore vanishes identically. To prove the second claim one has to use the description of the orthogonal complement $L^2_{\omega_\infty} \ominus H^2(\alpha)$ by means of the Hardy space [7] and once again apply the same argument related to the corresponding H^2 -space and orthonormal basis of reproducing kernels in it. \square

Lemma 2.3. *The multiplication by z with respect to the basis (2.10) is a 5-diagonal self-adjoint matrix,*

$$(2.11) \quad ze_{m+2} = \bar{r}_{m+2}e_m + \bar{p}_{m+2}e_{m+1} + q_{m+2}e_{m+2} + p_{m+3}e_{m+3} + r_{m+4}e_{m+4}.$$

Moreover $r_m > 0$ if and only if

$$(2.12) \quad t_{2n-1} = t_\infty^{-n}t_{(-1)}, \quad t_{2n} = t_{\lambda_0}^{-n}t_{(0)}.$$

Proof. Since the factor $b_{\lambda_0}b_\infty$ removes the singularities of z in $\bar{\mathbb{C}} \setminus E$ the function $zb_{\lambda_0}b_\infty F$ belongs to $H^2(\alpha\mu)$ for every $F \in H^2(\alpha)$ and an arbitrary $\alpha \in \pi_1(\bar{\mathbb{C}} \setminus E)^*$. Thus the decomposition of ze_{m+2} starts with e_m ,

$$ze_{m+2} = \bar{r}_{m+2}e_m + \dots$$

Since $z(\lambda)$ is real on E the multiplication operator is self-adjoint; its matrix possesses the symmetry property and therefore it is 5-diagonal (2.11). Finally we put $\lambda = \lambda_0$ in (2.11) for even m

$$(zb_{\lambda_0}b_\infty)(\lambda_0)K_{\lambda_0}^{\alpha\mu^{-(n+2)}}(\lambda_0)t_{2n+2} = \bar{r}_{2n+2}K_{\lambda_0}^{\alpha\mu^{-n}}(\lambda_0)t_{2n}$$

and $\lambda = \infty$ for odd m

$$(zb_{\lambda_0}b_\infty)(\infty)K_\infty^{\alpha\mu^\infty\mu^{-(n+2)}}(\infty)t_{2n+1} = \bar{r}_{2n+1}K_\infty^{\alpha\mu^\infty\mu^{-n}}(\infty)t_{2n-1}.$$

Since $K_\lambda^\alpha(\lambda) > 0$ we get (2.12). \square

Proof of Theorem 2.1. Let J be a periodic self-adjoint 5-diagonal matrix and \mathcal{R} be its spectral surface such that $\mathcal{R}_+ \simeq \bar{\mathbb{C}} \setminus E$. Since $w(\lambda)$ is single-valued in the domain, (2.7) and (2.9) imply that $\omega_j = \omega_{\lambda_0}(E_j) + \omega_\infty(E_j)$ are rational. Further, due to (2.3) the function wz^k is regular in the domain, moreover $1/(wz^k)(\lambda_0)$ and $1/(wz^k)(\infty)$ are roots of the quadratic equation

$$\begin{aligned} C^2 + \left(\frac{-1}{\prod_{j=0}^{k-1} r_{2j}} + \frac{-1}{\prod_{j=0}^{k-1} r_{2j+1}} \right) C + \frac{1}{\prod_{j=0}^{2k-1} r_j} \\ = \left(C - \frac{1}{\prod_{j=0}^{k-1} r_{2j}} \right) \left(C - \frac{1}{\prod_{j=0}^{k-1} r_{2j}} \right) = 0. \end{aligned}$$

Thus $(wz^k)(\lambda_0) > 0$ and $(wz^k)(\infty) > 0$. Since $w = t(b_{\lambda_0}b_\infty)^k$, $t \in \mathbb{T}$, the ratio $(zb_{\lambda_0}b_\infty)^k(\infty)/(zb_{\lambda_0}b_\infty)^k(\lambda_0)$ is also positive. That is $e^{2\pi i k \xi} = 1$. And this finishes the necessity part of the theorem.

In the opposite direction, for the given system of cuts we define J according to Lemma 2.3. It remains to check that J is periodic. Let $w = t_\infty^{-k}(b_{\lambda_0}b_\infty)^k$. Since $\mu^k(\gamma) = 1$, for all $\gamma \in \pi_1(\bar{\mathbb{C}} \setminus E)$, it is single-valued. Note that w is normalized by the condition $(wz^k)(\infty) = |(b_{\lambda_0}b_\infty z)^k(\infty)| > 0$. We claim that

$$(2.13) \quad we_n = e_{n+2k}.$$

For odd n (2.13) holds automatically. For even n we should take into account that in addition $t_\infty^{-k}t_{\lambda_0}^k = e^{-2\pi i k \xi} = 1$. Thus (2.13) defines the shift operator. Since the multiplication operators by z and w commute, we have $JS^{2k} = S^{2k}J$. Therefore J is periodic. \square

Now, let us restrict ourselves to the *real* case, i.e., $c_2 = \bar{c}_1$ or both critical values are real. Without loss of generality $c_2 = \bar{c}_1 = 2i$ or $c_2 = -c_1 = 2$. Thus, according to (2.5),

$$(2.14) \quad z = \lambda - \frac{1}{\lambda}$$

in the first case, and

$$(2.15) \quad z = \lambda + \frac{1}{\lambda}$$

in the second one.

In the case (2.14), $z^{-1}(\mathbb{R}) = \mathbb{R}$, thus E is a system of intervals on the real axis. Since $z^{-1}(\infty) = \{0, \infty\}$, E is subject to the restriction $\{0, \infty\} \not\subset E$.

If z is of the form (2.15), then $z^{-1}(\mathbb{R}) = \mathbb{R} \cup \mathbb{T}$, that is, E is a union of real intervals and arcs of the unit circle, and again $\{0, \infty\} \not\subset E$. This case under the additional assumption $E \subset \mathbb{T}$ leads to periodic CMV matrices [6]. Indeed, in the current case the multiplication by λ is also well defined and represents a unitary matrix A such that

$$(2.16) \quad J = A + A^{-1} = A + A^*.$$

As it was mentioned this functional model is the same as that related to periodic and almost periodic CMV matrices, see e.g. [4]. Conversely, having a periodic CMV matrix A we obtain the periodic self-adjoint J of the class by (2.16).

Similarly, in the case (2.14) the multiplication by λ leads to the self-adjoint operator A such that

$$(2.17) \quad J = A - A^{-1},$$

where A^{-1} exists and corresponds to the multiplication by $1/\lambda$, i.e., to a periodic SMP matrix. Further details of the corresponding functional model are discussed in Section 4. Note that the case (2.15) under the additional assumption $E \subset \mathbb{R}$ leads to essentially the same class of self-adjoint operators.

3. PROOF OF THE MAIN THEOREM

Let $E = [b_0, a_0] \setminus \cup_{j=1}^{\kappa} (a_j, b_j)$ be a system of intervals on \mathbb{R} , recall $\{0, \infty\} \in \bar{\mathbb{C}} \setminus E$, $z = \lambda - 1/\lambda$. We apply Theorem 2.1 in the current case. As it is well known the Green function (say with respect to infinity) is represented by the hyperelliptic integral, see e.g. [8, 7],

$$G(\lambda, \infty) = \operatorname{Re} \int_{a_0}^{\lambda} \frac{\lambda^{\kappa} + \dots}{\sqrt{\prod_{j=0}^{\kappa} (\lambda - a_j)(\lambda - b_j)}} d\lambda.$$

Therefore for the sum of the Green functions we have

$$(3.1) \quad G(\lambda, \infty) + G(\lambda, 0) = \operatorname{Re} \int_{a_0}^{\lambda} \frac{M_{\kappa+1}(\lambda)}{\sqrt{\prod_{j=0}^{\kappa} (\lambda - a_j)(\lambda - b_j)}} \frac{d\lambda}{\lambda},$$

where $M_{\kappa+1}$ is a monic polynomial of degree $\kappa + 1$. Note that the residue of the corresponding differential at the origin is -1 .



3.1. Spectrum of SMP matrices for the Stieltjes class. Let us consider the simplest case $E \subset \mathbb{R}_+$ or $E \subset \mathbb{R}_-$ (we can say that the spectrum is on the upper (lower) sheet of \mathcal{R}_+). The strong Stieltjes moment problem is related to measures supported on the positive half-axis [3]. The shape of the sum $G(\lambda, \infty) + G(\lambda, 0)$ on $\mathbb{R} \setminus E$ is shown in Fig. 5. It implies immediately that all the zeros of the polynomial $M_{\kappa+1}$ in (3.1) are real in this case. Indeed, each gap (a_j, b_j) , $j \geq 1$, contains at least one critical point; there is a critical point between $-\infty$ and 0; the total number of critical points is $\kappa + 1$. Therefore

$$(3.2) \quad \tilde{\theta}(\lambda) = i \int_{a_0}^{\lambda} \frac{M_{\kappa+1}(\lambda)}{\sqrt{\prod_{j=0}^{\kappa} (\lambda - a_j)(\lambda - b_j)}} \frac{d\lambda}{\lambda}$$

is the Schwarz-Christoffel integral, which maps conformally the upper half-plane \mathbb{H} onto the (generalized) polygon in Fig. 6.

According to (2.7) $w(\lambda) = te^{ik\tilde{\theta}(\lambda)}$, $t \in \mathbb{T}$. Let $\tilde{\omega}_j$ be the coordinates of the base of the slits. Then $w(\lambda)$ is single valued in $\bar{\mathbb{C}} \setminus E$ if and only if $\tilde{\omega}_j k \in \pi\mathbb{Z}$ for all j . It remains to mention that due to the chosen normalization for b_{λ_0} and b_{∞} the product $b_{\lambda_0}(\lambda)b_{\infty}(\lambda)(\lambda - 1/\lambda)$ is positive at infinity and negative at the origin, that is $\xi = 1/2$. Thus we can parametrize the spectral sets of periodic SMP matrices in this case by sufficiently simple domains shown in Fig. 6 with rational $\tilde{\omega}_j$'s (quite similar to the Jacobi and CMV matrices cases).

3.2. Complex critical points and three real critical points in the same gap. The situation changes dramatically as soon as $0 \in (a_j, b_j)$, $j \geq 1$, that is, $E = E_- \cup E_+$, $E_{\pm} \subset \mathbb{R}_{\pm}$. Still all gaps, except for (a_j, b_j) , should contain a critical point, thus $M_{\kappa+1}$ has at least $\kappa - 1$ real critical points. However, the positions of two remaining critical points are not a priori fixed.

FIGURE 5. $G_0 + G_\infty$ on \mathbb{R} : the spectrum is on the upper sheetFIGURE 6. Image of the Abelian integral $\tilde{\theta}$

First, we consider the case when **two remaining critical points are complex** μ_0 and $\overline{\mu_0}$, $\text{Im } \mu_0 > 0$. Let us consider $\tilde{\theta}(\lambda)$ in the upper half-plane \mathbb{H} . Since locally $\tilde{\theta}(\lambda) = \tilde{\theta}(\mu_0) + C(\lambda - \mu_0)^2 + \dots$, there exist two orthogonal directions where $\text{Re } d\tilde{\theta} = 0$. Moreover for one of them $\text{Im } \tilde{\theta}$ has a local minimum at μ_0 and a local maximum for another one. We define the curve γ , $\mu_0 \in \gamma$, by the condition $\text{Re } d\tilde{\theta} = 0$, such that $\text{Im } \tilde{\theta}$ increases. Since there is no other critical point in \mathbb{H} this curve should terminate on the real axis. Note that $\text{Im } \tilde{\theta}(\lambda)$ decreases as λ approaches E . If so, in the gaps γ may approach either a critical point or 0 and ∞ . The first case is also not possible since the critical point is a local minimum for $\text{Im } \tilde{\theta} = G_{\lambda_0} + G_\infty$ along the real axis, thus it should be local maximum in the orthogonal direction γ . But along γ it increases. Thus, γ terminates at 0 and ∞ , see Fig. 7.

As the result we get $\mathbb{H} \setminus \gamma = \mathbb{H}_- \cup \mathbb{H}_+$ such that $E_\pm \subset \partial \mathbb{H}_\pm$. Let $\theta(\lambda) = k\tilde{\theta}(\lambda)$. Inspecting the boundary behavior of the given analytic function we obtain that





FIGURE 7. A complex critical point

FIGURE 8. $D_{\pm} = \theta(\mathbb{H}_{\pm})$ regions for a complex critical point

it maps conformally \mathbb{H}_{\pm} onto D_{\pm} shown in Fig. 8, where the point $-\omega_0 + ih_0^2$ corresponds to the critical point μ_0 .

Now we define

$$(3.3) \quad u(\lambda) = \sqrt{-i\theta(\lambda) - i\omega_0 - h_0^2} \quad \text{for } \lambda \in \mathbb{H}_+$$

Here we assume that $\text{Im } u(\lambda) > 0$. Similarly we define

$$(3.4) \quad u(\lambda) = -\sqrt{-i\theta(\lambda) - i\omega_0 - h_0^2} \quad \text{for } \lambda \in \mathbb{H}_-$$

and in this case $\text{Im } u(\lambda) < 0$. In this way we get the regions Π_{\pm} . Since



$$\theta = -(2\text{Re } u \text{Im } u + \omega_0) + i((\text{Re } u)^2 - (\text{Im } u)^2 + h_0^2),$$

these regions are bounded by hyperbolic curves (1.1)–(1.4), see Fig. 1. Gluing the images along the curve γ we obtain the conformal mapping of the upper half-plane \mathbb{H} onto the special comb domain $\Pi = \Pi_+ \cup \Pi_- \cup \mathbb{R}$.

Conversely, for the region Π described by these equations we define a conformal map $u : \mathbb{H} \rightarrow \Pi$, $u(0) = +\infty$, $u(\infty) = -\infty$, and set

$$(3.5) \quad w(\lambda) = e^{-(u^2(\lambda) + h_0^2 + i\omega_0)}, \quad z = \lambda - \frac{1}{\lambda}$$

Then, the set E corresponds to $|w| = 1$. Since the base of the slits for D_{\pm} are of the form $\pi\ell$, w extended in the lower half-plane is single-valued in $\mathbb{C} \setminus E$. Finally,

FIGURE 9. Three real critical points in the same gap, $\mathbb{H}_+ = \mathbb{H}_{0+} \cup \mathbb{H}_{\infty+}$ FIGURE 10. $D_{\pm} = \theta(\mathbb{H}_{\pm})$ three real critical points in the same gap

w is real for $\lambda \in \mathbb{R} \setminus E$, that is, ξ is rational. Based on Theorem 2.1 we conclude that this domain can be associated to a periodic SMP matrix.

Let us turn to the case of **three real critical points in the same gap**. In this case \mathbb{H} can be decomposed into three pieces. Let $\mu_1 < \mu_0 < \mu_2$ be critical points in the gap (a_i, b_i) . Note that with necessity μ_1 and μ_2 are points of local maximum and $\text{Im } \theta$ assumes a local minimum at μ_0 in this interval. Therefore there are directions γ_1, γ_2 orthogonal to the real axis at μ_1 and μ_2 respectively such that $\text{Im } \theta$ increases. Arguments like the above show that these curves, $\text{Re } d\theta = 0$, terminate at 0 and ∞ , see Fig. 9

In each of them, $\theta(\lambda)$ represents a conformal mapping, see Fig. 10. In this picture $-\omega_0 + ih_0^2$, $-\omega_0 + ih_1^2$, and $-\omega_0 + ih_2^2$ are images of the critical points μ_0, μ_1 , and μ_2 respectively and $\omega_0 = \pi \ell_0$. We make the change of variable (3.3), (3.4), having in mind that now \mathbb{H}_+ or \mathbb{H}_- consists of two components. We arrive at the parametrization of the spectral curve by the domains of the form Fig. 2 such that

$$h_-^{(\ell_0)} = \sqrt{h_2^2 - h_0^2}, \quad h_+^{(\ell_0)} = \sqrt{h_1^2 - h_0^2}$$

in (1.7).

As before, starting from a region Π , by (3.5) we arrive at the set E and the domain $\mathbb{C} \setminus E \simeq \mathcal{R}_+$ which corresponds to a periodic SMP matrix.

3.3. Other cases. In the previous subsection we considered critical values in two main generic positions. Now let us list the remaining special cases.

1. For two complex critical values, if $\omega_0 = \pi\ell_0$ then one of the cuts in (1.3) and one in (1.4) degenerates to the intervals on the imaginary axis. The length of such a cut can not be arbitrary long, thus h_{ℓ_0} and $h_{k+\ell_0}$ are subject for the conditions (1.5). As soon as one of these values approaches h_0 two complex critical values, from the upper and lower half-planes, approach the critical value in the corresponding gap. In the limit we have the critical value of multiplicity 3. The same special case can be obtained when two critical values μ_1 and μ_2 tend to μ_0 , correspondingly $h_{\pm}^{(\ell_0)} \rightarrow 0$, see Fig. 2.

2. The case of a critical point of multiplicity two and a simple critical point in a gap corresponds to $h_+^{(\ell_0)} = 0$, $h_-^{(\ell_0)} > 0$ or $h_-^{(\ell_0)} = 0$, $h_+^{(\ell_0)} > 0$.

3. Two critical points (or one critical point of multiplicity two) may appear in the interval which contains zero or infinity. The domain Π looks similar to that one shown in Fig. 2, but the degenerated hyperbola corresponds to the most left (or right) position, i.e., $\ell_0 = 0$ or $\ell_0 = m$.

4. It was assumed that $m \leq k$. If $m > k$ the domain Π in Fig. 1 remains the same, but we switch the normalization conditions to $u(0) = -\infty$ and $u(\infty) = +\infty$.

5. In the Stieltjes case, subsection 3.1, the spectral curve was described by a simpler domain, Fig. 6. By (3.3) it can be transformed to a Π region bounded from below by the real axis.

4. FUNCTIONAL MODEL FOR PERIODIC SMP MATRICES

Let $\bar{\mathbb{C}} \setminus E$ correspond to a periodic SMP matrix. We define

$$(4.1) \quad \mathcal{A}(\alpha) = \frac{K_0^\alpha(\infty)}{K_\infty^\alpha(\infty)}, \quad \mathcal{B}(\alpha) = \frac{K_0^\alpha(0)}{K_\infty^\alpha(0)}, \quad \alpha \in \pi_1(\bar{\mathbb{C}} \setminus E)^*.$$

For the reader's convenience we prove here a known lemma, see e.g. [4].

Lemma 4.1. *The following identities hold true*

$$(4.2) \quad \mathcal{C}(\alpha) := \sqrt{1 - |\mathcal{A}(\alpha)|^2} = b_\infty(0) \frac{K_0^{\alpha\mu_\infty^{-1}}(0)}{K_0^\alpha(0)}$$

and

$$(4.3) \quad \overline{\mathcal{A}(\alpha)} = \frac{K_\infty^\alpha(0)}{K_0^\alpha(0)}, \quad \mathcal{C}(\alpha) = b_0(\infty) \frac{K_\infty^{\alpha\mu_0^{-1}}(\infty)}{K_\infty^\alpha(\infty)}.$$

Proof. $\mathcal{A}(\alpha)$ is defined by the following orthogonal decompositions

$$\begin{aligned} K_0^\alpha &= \mathcal{A}(\alpha) K_\infty^\alpha + \sqrt{1 - |\mathcal{A}(\alpha)|^2} b_\infty K_0^{\alpha\mu_\infty^{-1}} \\ b_0 K_\infty^{\alpha\mu_0^{-1}} &= \sqrt{1 - |\mathcal{A}(\alpha)|^2} K_\infty^\alpha - \overline{\mathcal{A}(\alpha)} b_\infty K_0^{\alpha\mu_\infty^{-1}} \end{aligned}$$

Indeed, $\mathcal{A}(\alpha) = \frac{K_0^\alpha(\infty)}{K_\infty^\alpha(\infty)}$ and we get (4.2).

Since, in addition,

$$\begin{aligned} K_\infty^\alpha &= \overline{\mathcal{A}(\alpha)} K_0^\alpha + \sqrt{1 - |\mathcal{A}(\alpha)|^2} b_0 K_\infty^{\alpha\mu_0^{-1}} \\ b_\infty K_0^{\alpha\mu_0^{-1}} &= \sqrt{1 - |\mathcal{A}(\alpha)|^2} K_0^\alpha - \mathcal{A}(\alpha) b_0 K_\infty^{\alpha\mu_0^{-1}} \end{aligned}$$

we have (4.3). □

In what follows without loss of generality we assume that $t_{(-1)} = 1$ in (2.12). In the given case $\lambda_0 = 0$, so t_0 is the new notation for t_{λ_0} , and this is not the same as the initial $t_{(0)}$. Since

$$(b_\infty b_0 z)(\infty) = (b_\infty \lambda)(\infty) b_0(\infty), \quad (b_\infty b_0 z)(0) = -(b_0/\lambda)(0) b_\infty(0),$$

we have $t_\infty = \phi_\infty/|\phi_\infty|$ and $t_0 = -\phi_0/|\phi_0|$, where

$$\phi_\infty = \left(\frac{b_\infty \lambda}{b_0} \right) (\infty), \quad \phi_0 = \left(\frac{b_0}{b_\infty \lambda} \right) (0).$$

Also, recall that $t_\infty/t_0 = e^{2\pi i \xi}$.

Theorem 4.2. *The multiplication operator by λ with respect to the basis (2.10) is a periodic SMP matrix $A = A(\alpha, t_{(0)})$ with the following coefficients*

$$\begin{aligned} \bar{p}_{2n} &= \phi_\infty t_{(0)} e^{2\pi i \xi n} \mathcal{A}(\alpha \mu^{-n}) \mathcal{B}^{-1}(\alpha \mu^{-n}) \mathcal{C}(\alpha \mu_\infty \mu^{-n}) \mathcal{B}(\alpha \mu_\infty \mu^{-n}) \\ (4.4) \quad p_{2n+1} &= -\bar{\phi}_\infty t_{(0)} t_\infty e^{2\pi i \xi n} \mathcal{C}(\alpha \mu^{-n}) \mathcal{B}(\alpha \mu^{-n}) \mathcal{A}(\alpha \mu_\infty \mu^{-n}) \mathcal{B}^{-1}(\alpha \mu_\infty \mu^{-n}) \\ q_{2n} &= -\phi_\infty \mathcal{A}(\alpha \mu^{-n}) \mathcal{B}^{-1}(\alpha \mu^{-n}) \overline{\mathcal{A}(\alpha \mu_\infty \mu^{-n})} \mathcal{B}(\alpha \mu_\infty \mu^{-n}) \end{aligned}$$

and

$$\begin{aligned} (4.5) \quad \pi_{2n+1} &= t_{(0)} t_\infty e^{2\pi i \xi n} \bar{\phi}_0 \mathcal{C}(\alpha \mu^{-n}) \mathcal{B}(\alpha \mu^{-n}) \mathcal{A}(\alpha \mu_\infty \mu^{-(n+1)}) \mathcal{B}^{-1}(\alpha \mu_\infty \mu^{-(n+1)}) \\ \bar{\pi}_{2n+2} &= -t_{(0)} e^{2\pi i \xi (n+1)} \phi_0 \mathcal{A}(\alpha \mu^{-n}) \mathcal{B}^{-1}(\alpha \mu^{-n}) \mathcal{C}(\alpha \mu_\infty \mu^{-(n+1)}) \mathcal{B}(\alpha \mu_\infty \mu^{-(n+1)}) \\ \sigma_{2n+1} &= -\phi_0 \mathcal{A}(\alpha \mu^{-n}) \mathcal{B}^{-1}(\alpha \mu^{-n}) \overline{\mathcal{A}(\alpha \mu_\infty \mu^{-(n+1)})} \mathcal{B}(\alpha \mu_\infty \mu^{-(n+1)}) \end{aligned}$$

Proof. Note that $b_\infty \lambda k_0^\alpha \in H^2(\alpha \mu_\infty)$ and it is orthogonal to the subspace $b_0 b_\infty^2 H^2(\alpha \mu_0^{-1} \mu_\infty^{-2})$. Therefore, in fact, we have the three-terms recurrence relation,

$$\lambda e_{2n} = \bar{p}_{2n} e_{2n-1} + q_{2n} e_{2n} + p_{2n+1} e_{2n+1}.$$

Moreover

$$\bar{p}_{2n} = t_{(0)} e^{2\pi i \xi n} (\lambda b_\infty)(\infty) \frac{K_0^{\alpha \mu^{-n}}(\infty)}{K_\infty^{\alpha \mu_\infty \mu^{-n}}(\infty)},$$

$$\begin{aligned}
q_{2n} &= \langle \lambda K_0^{\alpha\mu^{-n}}, K_0^{\alpha\mu^{-n}} \rangle \\
&= \left\langle \lambda K_0^{\alpha\mu^{-n}} - \frac{(\lambda b_\infty)(\infty) K_\infty^{\alpha\mu^{-n}}}{b_\infty K_\infty^{\alpha\mu^{-n}}(\infty)} K_0^{\alpha\mu^{-n}}(\infty), K_0^{\alpha\mu^{-n}} \right\rangle \\
&= - \frac{(\lambda b_\infty)(\infty)}{b_\infty(0)} \frac{K_\infty^{\alpha\mu^{-n}}(0)}{K_\infty^{\alpha\mu^{-n}}(\infty)} \frac{K_0^{\alpha\mu^{-n}}(\infty)}{K_0^{\alpha\mu^{-n}}(0)},
\end{aligned}$$

and

$$\begin{aligned}
p_{2n+1} &= \left\langle \lambda t_{(0)} t_0^{-n} K_0^{\alpha\mu^{-n}}, t_\infty^{-n-1} b_0 K_\infty^{\alpha\mu_0^{-1}\mu^{-n}} \right\rangle \\
&= t_{(0)} e^{2\pi i \xi n} t_\infty \left\langle K_0^{\alpha\mu^{-n}}, \lambda b_0 K_\infty^{\alpha\mu_0^{-1}\mu^{-n}} - \frac{(\lambda b_0 b_\infty)(\infty) K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(\infty) K_\infty^{\alpha\mu^{-n}}}{b_\infty K_\infty^{\alpha\mu^{-n}}(\infty)} \right\rangle \\
&= -t_{(0)} e^{2\pi i \xi n} t_\infty \frac{(\lambda b_\infty)(\infty)}{b_\infty(0)} \frac{K_\infty^{\alpha\mu^{-n}}(0)}{K_\infty^{\alpha\mu^{-n}}(\infty)} \frac{b_0(\infty) K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(\infty)}{K_0^{\alpha\mu^{-n}}(0)}.
\end{aligned}$$

In its turn,

$$\frac{1}{\lambda} e_{2n+1} = \bar{\pi}_{2n+1} e_{2n} + \sigma_{2n+1} e_{2n+1} + \pi_{2n+2} e_{2n+2},$$

where

$$\bar{\pi}_{2n+1} = \bar{t}_{(0)} \bar{t}_\infty e^{-2\pi i \xi n} \left(\frac{b_0}{\lambda} \right) (0) \frac{K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(0)}{K_0^{\alpha\mu^{-n}}(0)},$$

$$\begin{aligned}
\sigma_{2n+1} &= \left\langle \frac{1}{\lambda} K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}, K_\infty^{\alpha\mu_0^{-1}\mu^{-n}} \right\rangle \\
&= \left\langle \frac{1}{\lambda} K_\infty^{\alpha\mu_0^{-1}\mu^{-n}} - \left(\frac{b_0}{\lambda} \right) (0) \frac{K_0^{\alpha\mu^{-n}}}{b_0 K_0^{\alpha\mu^{-n}}(0)} K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(0), K_\infty^{\alpha\mu_0^{-1}\mu^{-n}} \right\rangle \\
&= - \left(\frac{b_0}{\lambda} \right) (0) \frac{1}{b_0(\infty)} \frac{K_0^{\alpha\mu^{-n}}(\infty)}{K_0^{\alpha\mu^{-n}}(0)} \frac{K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(0)}{K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(\infty)},
\end{aligned}$$

and

$$\begin{aligned}
\pi_{2n+2} &= \left\langle \frac{1}{\lambda} t_\infty^{-n-1} K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}, t_{(0)} t_0^{-n-1} b_\infty K_0^{\alpha\mu^{-1}\mu^{-n}} \right\rangle \\
&= \bar{t}_{(0)} e^{-2\pi i \xi(n+1)} \left\langle K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}, \frac{b_\infty}{\lambda} K_0^{\alpha\mu^{-n-1}} - \frac{K_0^{\alpha\mu^{-n}} \left(\frac{b_0 b_\infty}{\lambda} K_0^{\alpha\mu^{-n-1}} \right) (0)}{b_0 K_0^{\alpha\mu^{-n}}(0)} \right\rangle \\
&= -\bar{t}_{(0)} e^{-2\pi i \xi(n+1)} \left(\frac{b_0 b_\infty}{\lambda} \right) (0) \frac{1}{b_0(\infty)} \frac{K_0^{\alpha\mu^{-n}}(\infty)}{K_0^{\alpha\mu^{-n}}(0)} \frac{K_0^{\alpha\mu^{-n-1}}(0)}{K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(\infty)}.
\end{aligned}$$

Now by making use of (4.1), (4.3), (4.2), we obtain (4.4)

$$\begin{aligned}
q_{2n} &= -\phi_\infty \overline{\mathcal{A}(\alpha\mu_\infty\mu^{-n})} \mathcal{B}(\alpha\mu_\infty\mu^{-n}) \mathcal{A}(\alpha\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu^{-n}), \\
\bar{p}_{2n} &= \phi_\infty t_{(0)} e^{2\pi i \xi n} b_\infty(0) \frac{K_0^{\alpha\mu^{-n}}(\infty)}{K_\infty^{\alpha\mu_\infty\mu^{-n}}(\infty)} \\
&= \phi_\infty t_{(0)} e^{2\pi i \xi n} \frac{K_0^{\alpha\mu^{-n}}(\infty)}{K_\infty^{\alpha\mu^{-n}}(\infty)} \frac{K_\infty^{\alpha\mu^{-n}}(\infty)}{K_\infty^{\alpha\mu_\infty\mu^{-n}}(\infty)} b_\infty(0) \\
&= \phi_\infty t_{(0)} e^{2\pi i \xi n} \mathcal{A}(\alpha\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu^{-n}) \mathcal{C}(\alpha\mu_\infty\mu^{-n}) \mathcal{B}(\alpha\mu_\infty\mu^{-n}), \\
\bar{p}_{2n+1} &= -\phi_\infty \bar{t}_{(0)} \bar{t}_\infty e^{-2\pi i \xi n} \overline{\mathcal{A}(\alpha\mu_\infty\mu^{-n})} \mathcal{B}(\alpha\mu_\infty\mu^{-n}) \\
&\quad \times \frac{b_0(\infty) K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(\infty)}{K_\infty^{\alpha\mu^{-n}}(\infty)} \frac{K_\infty^{\alpha\mu^{-n}}(\infty)}{K_0^{\alpha\mu^{-n}}(0)} \\
&= -\phi_\infty \bar{t}_{(0)} \bar{t}_\infty e^{-2\pi i \xi n} \overline{\mathcal{A}(\alpha\mu_\infty\mu^{-n})} \mathcal{B}(\alpha\mu_\infty\mu^{-n}) \mathcal{C}(\alpha\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu^{-n}),
\end{aligned}$$

as well as (4.5)

$$\begin{aligned}
\bar{\pi}_{2n+1} &= \bar{t}_{(0)} \bar{t}_\infty e^{-2\pi i \xi n} \phi_0 \frac{K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(0)}{K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(\infty)} b_0(\infty) \frac{K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(\infty)}{K_\infty^{\alpha\mu^{-n}}(\infty)} \frac{K_\infty^{\alpha\mu^{-n}}(\infty)}{K_0^{\alpha\mu^{-n}}(0)} \\
&= \bar{t}_{(0)} \bar{t}_\infty e^{-2\pi i \xi n} \phi_0 \overline{\mathcal{A}(\alpha\mu_\infty\mu^{-(1+n)})} \mathcal{B}(\alpha\mu_\infty\mu^{-(1+n)}) \mathcal{C}(\alpha\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu^{-n}), \\
\sigma_{2n+1} &= -\phi_0 \mathcal{A}(\alpha\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu^{-n}) \overline{\mathcal{A}(\alpha\mu_\infty\mu^{-(n+1)})} \mathcal{B}(\alpha\mu_\infty\mu^{-(n+1)}) \\
\bar{\pi}_{2n+2} &= -t_{(0)} e^{2\pi i \xi(n+1)} \phi_0 \mathcal{A}(\alpha\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu^{-n}) \times b_0(\infty) \frac{K_0^{\alpha\mu^{-(n+1)}}(0)}{K_0^{\alpha\mu_0^{-(n+1)}}(0)} \frac{K_0^{\alpha\mu_0^{-1}\mu^{-n}}(0)}{K_\infty^{\alpha\mu_0^{-1}\mu^{-n}}(\infty)} \\
&= -t_{(0)} e^{2\pi i \xi(n+1)} \phi_0 \mathcal{A}(\alpha\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu^{-n}) \mathcal{C}(\alpha\mu_\infty\mu^{-(n+1)}) \mathcal{B}(\alpha\mu_\infty\mu^{-(n+1)}). \quad \square
\end{aligned}$$

Remark 4.3. The structure of the reproducing kernels on the hyperelliptic Riemann surfaces is well known, see e.g. [7]. In particular, indeed $K_\infty^\alpha(0) = 0$, i.e., $\mathcal{A}(\alpha) = 0$, for some α . According to (4.4) and (4.5) it means that the corresponding $A = A(\alpha, t_{(0)})$ may degenerate, that is, q_{2n} or σ_{2n-1} vanishes for some n . Nevertheless all entries of A and A^{-1} have perfect sense. For example,

$$\begin{aligned}
r_{2n+1} &= \frac{\bar{p}_{2n} \bar{p}_{2n+1}}{q_{2n}} \\
&= |\phi_\infty| \mathcal{C}(\alpha\mu^{-n}) \mathcal{B}(\alpha\mu^{-n}) \mathcal{C}(\alpha\mu_\infty\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu_\infty\mu^{-n}), \\
-\rho_{2n} &= \frac{\bar{\pi}_{2n-1} \bar{\pi}_{2n}}{-\sigma_{2n-1}} \\
&= |\phi_0| \mathcal{C}(\alpha\mu^{-(n-1)}) \mathcal{B}(\alpha\mu^{-(n-1)}) \mathcal{C}(\alpha\mu_\infty\mu^{-n}) \mathcal{B}^{-1}(\alpha\mu_\infty\mu^{-n}),
\end{aligned}$$

where

$$\begin{aligned} Ae_{2n-1} &= r_{2n-1}e_{2n-3} + \bar{p}_{2n-1}e_{2n-2} + q_{2n-1}e_{2n-1} + p_{2n}e_{2n} + r_{2n+1}e_{2n+1}, \\ A^{-1}e_{2n} &= \rho_{2n}e_{2n-2} + \bar{\pi}_{2n}e_{2n-1} + \sigma_{2n}e_{2n} + \pi_{2n+1}e_{2n+1} + \rho_{2n+2}e_{2n+2}. \end{aligned}$$

REFERENCES

1. A. Eremenko and P. Yuditskii, *Comb functions*, Proceedings of the Conference on Orthogonal Polynomials, Special Functions, and Applications (2011), to appear.
2. P. van Moerbekke and D. Mumford, *The spectrum of difference operators and algebraic curves*, Acta Math. **143**(1-2), 1979, 93–154.
3. O. Njåstad, *Solutions of the strong Hamburger moment problem*. J. Math. Anal. Appl. 197 (1996), no. 1, 227–248.
4. F. Peherstorfer and P. Yuditskii, *Almost periodic Verblunsky coefficients and reproducing kernels on Riemann surfaces*, J. Approx. Theory **139**(1-2) (2006), 91–106.
5. B. Simon, *Orthogonal polynomials on the unit circle*. Part 1. Classical theory. American Mathematical Society Colloquium Publications, 54, Part 1. American Mathematical Society, Providence, RI, 2005. xxvi+466 pp.
6. B. Simon, *Orthogonal polynomials on the unit circle*. Part 2. Spectral theory. American Mathematical Society Colloquium Publications, 54, Part 2. American Mathematical Society, Providence, RI, 2005. pp. i–xxii and 467–1044.
7. M. Sodin and P. Yuditskii, *Almost periodic Jacobi matrices with homogeneous spectrum, infinite dimensional Jacobi inversion, and hardy spaces of character-automorphic functions*, J. Geom. Anal. **7** (1997), 387–435.
8. Yu. Tomchuk, *Orthogonal polynomials over a system of intervals on the real line*, Zap. Fiz.-Mat. Fak. i Kharkov. Mat. Obshch. (4) **29** (1963), 93–128. (Russian).

IONELA MOALE

RESEARCH INSTITUTE FOR SYMBOLIC COMPUTATION

JOHANNES KEPLER UNIVERSITY LINZ

ALTENBERGERSTR. 69, A-4040, LINZ, AUSTRIA

E-mail address: Ionela.Moale@risc.jku.at

PETER YUDITSKII

ABTEILUNG FÜR DYNAMISCHE SYSTEME UND APPROXIMATIONSTHEORIE

JOHANNES KEPLER UNIVERSITÄT LINZ

A-4040 LINZ, AUSTRIA

E-mail address: Petro.Yuditskiy@jku.at