

G_1 -STRUCTURES OF SECOND ORDER

DEMETRA DEMETROPOULOU-PSOMOPOULOU

Abstract

We introduce a generalization to the second order of the notion of the G_1 -structure, the so called generalized almost tangent structure. For this purpose, the concepts of the second order frame bundle $H^2(V_m)$, its structural group L_m^2 and its associated tangent bundle of second order $T^2(V_m)$ of a differentiable manifold V_m are described from the point of view that is used. Then, a G_1 -structure of second order -called G_1^2 -structure- is constructed on V_m by an endomorphism J acting on $T^2(V_m)$, satisfying the relation $J^2 = 0$ and some hypotheses on its rank. Its connection and characteristic cohomology class are defined.

Some of the G -structures of the first order are those defined by nilpotent operators of degree $r + 1$ ($r \geq 1$) that is, the G_r -structures, defined by J. Lehman-Lejeune ([15]) and studied by H.A. Eliopoulos ([11]).

The G_1 -structure of the first order, briefly G_1 -structure, is defined ([15]) on an m -dimensional differentiable manifold V_m of class C^∞ by means of an 1-form J , of constant rank p , with values in the tangent bundle, such that at each point $x \in V_m$,

$$(1) \quad J_x^2 = 0.$$

That is,

$\dim \text{Im } J_x = p \geq 1$, $\dim \ker J_x = q$, $m = p + q$ and q independent of the point x of V_m .

The G_1 -structure is also studied by [1]; it is called generalized almost tangent structure.

Our objective in the present paper is to find a prolongation of this structure, that is, there is defined a G -structure of second order on V_m , called the G_1 -structure of order 2, briefly a G_1^2 -structure, by means of

an 1-form of second order J , of constant rank $p + \binom{p+1}{2}$, with values in the tangent bundle of second order, satisfying at each point $x \in V_m$ the relation (1).

At first, a brief discussion of the notions of the frame bundle of second order $H^2(V_m)(V_m, L_m^2)$ and its associated ([13]) tangent bundle $T^2(V_m)(V_m, L_m^2, F^2)$ of order two, will be given from the standpoint used in the generalization of the real almost product structure to the second order ([6]).

Then, the G_1^2 -structure, its adapted basis, connection and characteristic cohomology class will be defined.

1. The fibre bundles $H^2(V_m)$ and $T^2(V_m)$

We recall from [6] the following:

Let V_m be an m -dimensional differentiable manifold of class C^∞ and $H^2(V_m) = \bigcup_{x \in V_m} H_x^2(V_m)$ the fibre bundle of all 2-frames of the manifold

V_m , where H_x^2 is ([7]) the set of all invertible 2-jets of R^m into V_m with source $0 \in R^m$ and target $x \in V_m$. This bundle is ([8]) a principal fibre bundle with basis V_m and structural group L_m^2 ; it is called the principal prolongation of order 2 of the manifold V_m .

The structural group L_m^2 is ([10]) the set $j^2 \circ f$ of all invertible 2-jets with source and target $0 \in R^m$ of a 2-mapping f at the point $0 \in R^m$. Hence, each $a \in L_m^2$ can be written in the form,

$$(1.1) \quad a = (a_{j_1}^i, a_{j_1 j_2}^i), \quad i, \quad j_1, \quad j_2 = 1, 2, \dots, m, \quad \det(a_{j_1}^i) \neq 0$$

and $a_{j_1 j_2}^i$ is symmetric with respect to j_1, j_2 .

Also,

$$\dim L_m^2 = m \binom{m+2}{2} - m = m^2 + m \binom{m+1}{2} = m^2 + m^2 \left(\frac{m+1}{2} \right).$$

If

$$\beta = (\beta_{k_1}^{j_1}, \beta_{k_1 k_2}^{j_1}) \in L_m^2,$$

then, from the composition of 2-jets ([8]), it follows that the product of the two elements a and β of L_m^2 ,

$$a\beta = c = (c_{k_1}^{j_1}, c_{k_1 k_2}^{j_1}),$$

can be defined by the relations

$$\begin{cases} c_{k_1}^{i_1} = a_{j_1}^{i_1} \beta_{k_1}^{j_1}, \\ c_{k_1 k_2}^{i_1} = a_{j_1}^{i_1} \beta_{k_1 k_2}^{j_1} + a_{j_1 j_2}^{i_1} \beta_{k_1}^{j_1} \beta_{k_2}^{j_2}. \end{cases}$$

Remark 1.1. We assume, now, that to the element $a \in L_m^2$, corresponds the matrix A of the form,

$$(1.2) \quad A = \begin{bmatrix} a_{j_1}^{i_1} & 0 \\ a_{j_1 j_2}^{i_1} & a_{j_1}^{i_1} a_{j_2}^{i_1} \end{bmatrix}.$$

If B is a matrix of the same form, corresponding to the element β of L_m^2 , then to $a\beta \in L_m^2$ corresponds the product of matrices AB ,

$$\begin{aligned} AB = C &= \begin{bmatrix} c_{k_1}^{i_1} & 0 \\ c_{k_1 k_2}^{i_1} & c_{k_1}^{i_1} c_{k_2}^{i_2} \end{bmatrix} = \begin{bmatrix} \beta_{k_1}^{j_1} & 0 \\ \beta_{k_1 k_2}^{j_1} & \beta_{k_1}^{j_1} \beta_{k_2}^{j_2} \end{bmatrix} \begin{bmatrix} a_{j_1}^{i_1} & 0 \\ a_{j_1 j_2}^{i_1} & a_{j_1}^{i_1} a_{j_2}^{i_2} \end{bmatrix} = \\ &= \begin{bmatrix} a_{j_1}^{i_1} \beta_{k_1}^{j_1} & 0 \\ a_{j_1}^{i_1} \beta_{k_1 k_2}^{j_1} + a_{j_1 j_2}^{i_1} \beta_{k_1}^{j_1} \beta_{k_2}^{j_2} & a_{j_1}^{i_1} a_{j_2}^{i_2} \beta_{k_1}^{j_1} \beta_{k_2}^{j_2} \end{bmatrix}, \end{aligned}$$

with the notation that first is written the matrix B and then A , but after the multiplication (each line of B with every column of A) first is written the elements of A and then of B .

It can be verified that to each element $a \in L_m^2$ corresponds a matrix A of the form (1.2) and conversely.

Hence, the group of matrices of the form A , subgroup of the group of matrices $Gl(N, R)$, $N = m + \frac{m(m+1)}{2}$, can be identified with the group L_m^2 .

The Lie algebra \underline{L}_m^2 of the Lie group L_m^2 is ([17]) defined by,

$$\underline{L}_m^2 = \{ \lambda / \lambda = (\lambda_{j_1}^i, \lambda_{j_1 j_2}^i), i, j_1, j_2 = 1, 2, \dots, m, \\ \lambda_{j_1}^i \in R^m \otimes R^{m^*}, \lambda_{j_1 j_2}^i \in R^m \otimes S^2(R^{m^*}) \},$$

where $S^2(R^{m^*})$ is the set of the 2-linear symmetric forms on R^m .

We consider, also, $T_{1,x}^2 * (V_m)$ the set of all 2-jets of the functions on V_m with source $x \in V_m$ and target 0. The set $T_1^{2*}(V_m) = \bigcup_{x \in V_m} T_{1,x}^{2*}(V_m)$ has ([8], [10]) the structure of a vector bundle with basis V_m structural group L_m^2 and fibre $L_{1,m}^2$; it is associated with $H^2(V_m)$.

$L_{1,m}^2$ is ([7], [10]) the set $j^2 \circ g$ of all 2-jets with source $0 \in R^m$ and target $0 \in R$ of a 2-mapping g at $0 \in R^m$. Therefore, each $y \in L_{1,m}^2$ can be written in the form,

$$(1.3) \quad y = (y_{i_1}, y_{i_1 i_2}), i_1, i_2 = 1, 2, \dots, m, y_{i_1 i_2} \\ \text{is symmetric with respect to } i_1, i_2.$$

Also,

$$\dim L_{1,m}^2 = \binom{m+2}{2} - 1 = m + \binom{m+1}{2} = m + \frac{m(m+1)}{2}.$$

Let $\{x^i\}_{i=1,2,\dots,m}$ be a system of local coordinates at $x \in V_m$ for a given chart. Then ([8]), the element $\omega \in T_1^2 * (V_m)$ can be expressed in the form,

$$(1.4) \quad \omega = (x^i, y_{i_1}, y_{i_1 i_2}), \quad i, i_1, i_2 = 1, 2, \dots, m, \quad y_{i_1 i_2}$$

is symmetric with respect to i_1, i_2 .

If

$$(x^{j'}, y_{j'_1}, y_{j'_1 j'_2}),$$

is the expression of ω in a new coordinate system $\{x^{j'}\}_{j'=1,2,\dots,m}$ at $x \in V_m$, then ([10]) the transformation law for the local coordinates of ω is given by the equations,

$$(1.5) \quad \begin{cases} x^i = \varphi^i(x^{j'}), \\ y_{j'_1} = y_{i_1} a_{j'_1}^{i_1}, \\ y_{j'_1 j'_2} = y_{i_1} a_{j'_1 j'_2}^{i_1} + y_{i_1 i_2} a_{j'_1}^{i_1} a_{j'_2}^{i_2}, \end{cases}$$

where $a_{j'_1}^i = \frac{\partial \varphi^i}{\partial x^{j'_1}}$, $a_{j'_1 j'_2}^i = \frac{\partial^2 \varphi^i}{\partial x^{j'_1} \partial x^{j'_2}}$, so that $(a_{j'_1}^i, a_{j'_1 j'_2}^i) \in L_m^2$.

The dual vector bundle $T^2(V_m)$ of the vector bundle $T_1^2 * (V_m)$, has ([4]) the basis V_m , the structural group L_m^2 and the fibre $F^2 = (L_{1,m}^2)^*$. Hence ([2]) $T^2(V_m)(V_m, L_m^2, F^2)$ is the vector bundle of all tangent vectors of order 2 and $v \in T_x^2$ is a tangent vector of order 2 (or 2-tangent vector) at the point $x \in V_m$.

Remark 1.2. However, there are other, different, notions of the tangent bundle of higher order ([16], [19]) using another point of view.

From [6] follows that $T^2(V_m)(V_m, L_m^2, F^2)$ is also associated with the principal fibre bundle $H^2(V_m)(V_m, L_m^2)$, which can be identified with the space of bases of the vector spaces T_x^2 at $x \in V_m$.

Let now,

$$(1.6) \quad e = (e_{i_1}, e_{i_1 i_2}), \quad i_1, i_2 = 1, 2, \dots, m, \quad e_{i_1 i_2} \text{ symmetric in the indices } i_1, i_2,$$

be the natural basis of T_x^2 defined by the local chart $\{x^i\}_{i=1,2,\dots,m}$ at the point $x \in V_m$. Then, every $v \in T_x^2$ can be expressed uniquely in the form,

$$(1.7) \quad v = \sum_{i_1=1}^m v^{i_1} e_{i_1} + \sum_{i \leq i_1 \leq i_2 \leq m} v^{i_1 i_2} e_{i_1 i_2},$$

where $(v^{i_1}, v^{i_1 i_2})$ are some constants and $v^{i_1 i_2}$ is symmetric in the indices i_1, i_2 .

For another local chart $\{x^{j'}\}_{j=1,2,\dots,m}$ at $x \in V_m$ the corresponding transformation law for the local coordinates of $v \in T_x^2$ will be,

$$(1.8) \quad \begin{cases} v^{j'_1} = a_{i_1}^{j'_1} v^{i_1} + a_{i_1 i_2}^{j'_1} v^{i_1 i_2}, \\ v^{j'_1 j'_2} = a_{i_1}^{j'_1} a_{i_2}^{j'_2} v^{i_1 i_2}, \end{cases}$$

where $a = (a_{i_1}^{j'_1}, a_{i_1 i_2}^{j'_1}) \in L_m^2$.

For convenience in calculation, from now on, we will keep using matrices.

Thus, the relation (1.8) for the element v of T_x^2 in the overlap of two local charts (U, x^i) and $(V, x^{j'})$, can be written in the form,

$$(1.9) \quad \begin{bmatrix} v^{j'_1} & v^{j'_1 j'_2} \end{bmatrix} = \begin{bmatrix} v^{i_1} & v^{i_1 i_2} \end{bmatrix} \begin{bmatrix} a_{i_1}^{j'_1} & 0 \\ a_{i_1 i_2}^{j'_1} & a_{i_1}^{j'_1} a_{i_2}^{j'_2} \end{bmatrix},$$

or briefly,

$$V_x^v = A_u^v(x) V_x^u,$$

where

$$(1.10) \quad \begin{cases} A_u^v(x) = \begin{bmatrix} a_{i_1}^{j'_1} & 0 \\ a_{i_1 i_2}^{j'_1} & a_{i_1}^{j'_1} a_{i_2}^{j'_2} \end{bmatrix}, V_x^u = [v^{i_1} \ v^{i_1 i_2}], \\ V_x^v = [v^{j'_1} \ v^{j'_1 j'_2}]. \end{cases}$$

The element $a = (a_{i_1}^{j'_1}, a_{i_1 i_2}^{j'_1})$ of L_m^2 is identified with the matrix A defined by (1.10).

Similarly, according to the relation (1.5), the transformation law for each element ω of T_x^{2*} (that is of $T_{1,x}^2 * (V_m)$) by means of matrices, can be written in the form,

$$(1.11) \quad \begin{bmatrix} \omega_{j'_1} \\ \omega_{j'_1 j'_2} \end{bmatrix} = \begin{bmatrix} a_{i_1}^{i_1} & 0 \\ a_{i_1 i_2}^{i_1} & a_{i_1}^{i_1} a_{i_2}^{i_2} \end{bmatrix} \begin{bmatrix} \omega_{i_1} \\ \omega_{i_1 i_2} \end{bmatrix},$$

where $(a_{j'_1}^{i_1}, a_{j'_1 j'_2}^{i_1}) \in L_m^2$ is the inverse element of $a = (a_{i_1}^{j'_1}, a_{i_1 i_2}^{j'_1}) \in L_m^2$. Briefly,

$$\omega_v^x = \omega_u^x A_v^u(x), \quad A_v^u(x) = [A_u^v(x)]^{-1}.$$

2. G_1^2 -structures

Definition 2.1. A G_1 -structure of second order, briefly G_1^2 -structure, can be defined on an m -dimensional differentiable manifold V_m of class C^∞ , by means of an 1-form of second order J , with values in the second order tangent bundle $T^2(V_m)$, such that at each point $x \in V_m$,

$$(2.1) \quad J_x \text{ is of rank } p + \binom{p+1}{2} \text{ everywhere in } V_m$$

and

$$(2.2) \quad J_x^2 = 0.$$

From the condition (2.2) follows that $J_x(T_x^2)$ is composed of the eigenvectors of J . On the other hand $J_x(T_x^2) \subset \ker J_x$.

If S_x is the complementary space of $\ker J_x$ with respect to $T_x^2(V_m)$ then,

$$T_x^2(V_m) = \ker J_x \oplus S_x$$

and J induces an isomorphism between S_x and $J(S_x)$.

Thus,

$$\dim T_x^2(V_m) = m + \binom{m+1}{2} > 2 \left[p + \binom{p+1}{2} \right] \text{ and } m \geq 2p.$$

We have,

$$\dim S_x = p + \binom{p+1}{2}$$

and

$$\dim \ker J_x = q + \binom{q+1}{2} + pq, \quad p \leq q, \quad m = p + q$$

and the dimension of $\ker J_x$ is independent of the point $x \in V_m$.

Obviously,

$$\dim T_x^2 = p + \binom{p+1}{2} + q + \binom{q+1}{2} + pq.$$

Also,

$$(2.3) \quad \dim \ker J_x =$$

$$= p + (q-p) + \binom{p+1}{2} + p(q-p) + p(m-q) + \binom{q-p+1}{2} + (q-p)(m-q).$$

Let $\{e_{i_1}, e_{i_1 i_2}\}$, $i_1, i_2 = 1, 2, \dots, m$ be a basis of the second order tangent space T_x^2 . Using the indices,

$$\begin{aligned}\alpha(1) &= 1, 2, \dots, p, \alpha(2) = p+1, \dots, q, \\ A(1) &= 1, 2, \dots, q \text{ (that is, } A(1) = (\alpha(1), \alpha(2))\text{)} \text{ and} \\ A(2) &= q+1, \dots, q+p = m \text{ (that is, } A(2) = q+\alpha(1)\text{)},\end{aligned}$$

it follows, that the second order tangent vectors

$$(2.4) \quad \{e_{A_1(1)}, e_{A_1(1)A_2(1)}, e_{\alpha_1(1)A_2(1)}\}$$

and

$$(2.5) \quad \{e_{A_1(2)}, e_{A_1(2)A_2(2)}\}$$

define a basis on $\ker J_x$ and S_x respectively.

Also, by (2.3) the basis (2.4) can be written in the form,

$$(2.6) \quad \{(e_{\alpha_1(1)}, e_{\alpha_1(2)}), \\ (e_{\alpha_1(1)\alpha_2(1)}, e_{\alpha_1(1)\alpha_2(2)}, e_{\alpha_1(1)A_2(2)}, e_{\alpha_1(2)\alpha_2(2)}, e_{\alpha_1(2)A_2(2)})\}.$$

Definition 2.2. The basis $\{e_{i_1}, e_{i_1 i_2}\}$ of T_x^2 will be called adapted to the G_1^2 -structure with respect to x or simply G_1^2 -adapted basis if

$$(2.7) \quad \begin{cases} J_x e_{A_1(2)} = e_{\alpha_1(1)}, \\ J_x e_{A_1(2)A_2(2)} = e_{\alpha_1(1)\alpha_2(1)}. \end{cases}$$

To the operator J_x corresponds the element F of the tensor product $T_x^2 \otimes (T_x^2)^*$,

$$(2.8) \quad \begin{cases} F = (F_{i_1}^{j_1}, F_{i_1}^{j_1 j_2}, F_{i_1 i_2}^{j_1}, F_{i_1 i_2}^{j_1 j_2}) \text{ } i_1, i_2, j_1, j_2 = 1, 2, \dots, m, \\ F_{i_1}^{j_1 j_2} \text{ symmetric with respect to } j_1, j_2, F_{i_1 i_2}^{j_1} \text{ symmetric} \\ \text{with respect to } i_1, i_2 \text{ and } F_{i_1 i_2}^{j_1 j_2} \text{ symmetric in indices } i_1, \\ i_2 \text{ and } j_1, j_2. \end{cases}$$

Then, J is given by

$$(2.9) \quad \begin{cases} (J_x v)^{j_1} = F_{i_1}^{j_1} v^{i_1} + F_{i_1 i_2}^{j_1} v^{i_1 i_2}, \\ (J_x v)^{j_1 j_2} = F_{i_1}^{j_1 j_2} v^{i_1} + F_{i_1 i_2}^{j_1 j_2} v^{i_1 i_2}, \end{cases}$$

where $v = (v^{i_1}, v^{i_1 i_2})$ is a 2-tangent vector at $x \in V_m$.

The tensor F of the relation (2.8) can be represented by the following matrix,

$$(2.10) \quad F = \begin{bmatrix} F_{i_1}^{j_1} & F_{i_1}^{j_1 j_2} \\ F_{i_1 i_2}^{j_1} & F_{i_1 i_2}^{j_1 j_2} \end{bmatrix}.$$

From (2.2) and (2.9) one verifies easily the following equations,

$$(2.11) \quad \begin{cases} F_{i_1}^{j_1} F_{j_1}^{k_1} + F_{i_1}^{j_1 j_2} F_{j_1 j_2}^{k_1} = 0, \\ F_{i_1}^{j_1} F_{j_1}^{k_1 k_2} + F_{i_1}^{j_1 j_2} F_{j_1 j_2}^{k_1 k_2} = 0, \\ F_{i_1 i_2}^{j_1} F_{j_1}^{k_1} + F_{i_1 i_2}^{j_1 j_2} F_{j_1 j_2}^{k_1} = 0, \\ F_{i_1 i_2}^{j_1} F_{j_1}^{k_1 k_2} + F_{i_1 i_2}^{j_1 j_2} F_{j_1 j_2}^{k_1 k_2} = 0. \end{cases}$$

Using, now, an G_1^2 -adapted basis on T_x^2 and the relations:

$$\begin{aligned} J_x e_{\alpha_1(1)} &= 0, & J_x e_{\alpha_1(2)} &= 0, & J_x e_{A_1(2)} &= e_{\alpha_1(1)}, \\ J_x e_{\alpha_1(1)\alpha_2(1)} &= 0, & J_x e_{\alpha_1(1)\alpha_2(2)} &= 0, & J_x e_{\alpha_1(1)A_2(2)} &= 0, \\ J_x e_{\alpha_1(2)\alpha_2(2)} &= 0, & J_x e_{\alpha_1(2)A_2(2)} &= 0, & J_x e_{A_1(2)A_2(2)} &= e_{\alpha_1(1)\alpha_2(1)}, \end{aligned}$$

it follows that the tensor F associated to the operator J can be represented by the matrix:

$$(2.12) \quad F = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ \delta_{\alpha_1(1)}^{\beta_1(1)} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Thus we have,

Proposition 2.1. *There is always a G_1^2 -adapted basis of $T_x^2(V_m)$ at each point $x \in V_m$, in which F has the constant components given by the matrix (2.12).*

Let $\{e_{j'_1}, e_{j'_1 j'_2}\}$ be another G_1^2 -adapted basis, then the transformation law between the G_1^2 -adapted bases may be written,

$$\begin{aligned} e_{i_1} &= e_{j'_1} l_{i_1}^{j'_1}, \\ e_{i_1 i_2} &= e_{j'_1} l_{i_1 i_2}^{j'_1} + e_{j'_1 j'_2} l_{i_1}^{j'_1} l_{i_2}^{j'_2}, \end{aligned}$$

where the element $l = (l_{i_1}^{j'_1}, l_{i_1 i_2}^{j'_1}) \in L_m^2$. Explicitly,

$$(2.13) \quad \left\{ \begin{array}{l} l_{i_1}^{j'_1} = \left(l_{\alpha_1(1)}^{\beta'_1(1)}, l_{\alpha_1(2)}^{\beta'_1(1)}, l_{A_1(2)}^{\beta'_1(1)}, l_{\alpha_1(2)}^{\beta'_1(2)}, l_{A_1(2)}^{\beta'_1(2)} \right) \in G_1 \\ l_{i_1 i_2}^{j'_1} = \left(l_{\alpha_1(1)\alpha_2(1)}^{\beta'_1(1)}, l_{\alpha_1(1)\alpha_2(2)}^{\beta'_1(1)}, l_{\alpha_1(1)A_2(2)}^{\beta'_1(1)}, \right. \\ \quad l_{\alpha_1(2)\alpha_2(2)}^{\beta'_1(1)}, l_{\alpha_1(2)A_2(2)}^{\beta'_1(1)}, l_{A_1(2)A_2(2)}^{\beta'_1(1)}, \\ \quad l_{\alpha_1(1)\alpha_2(2)}^{\beta'_1(2)}, l_{\alpha_1(1)A_2(2)}^{\beta'_1(2)}, l_{\alpha_1(2)\alpha_2(2)}^{\beta'_1(2)}, \\ \quad \left. l_{\alpha_1(2)A_2(2)}^{\beta'_1(2)}, l_{A_1(2)A_2(2)}^{\beta'_1(2)} \right), \end{array} \right.$$

where G_1 is ([15]) the structural group of the G_1 -structure of the first order. It is consisting of the matrices of the form

$$\begin{bmatrix} A & 0 & 0 \\ B & \Gamma & 0 \\ \Delta & E & A \end{bmatrix},$$

with $A \in L_p$, $B \in \text{End}(R^p, R^{q-p})$, $\Gamma \in L_{q-p}$, $\Delta \in \text{End}(R^p, R^{m-q})$ and $E \in \text{End}(R^{q-p}, R^{m-q})$.

Using matrices, l can be written by the matrix

$$(2.14) \quad L = \begin{bmatrix} l_{i_1}^{j'_1} & 0 \\ l_{i_1 i_2}^{j'_1} & l_{i_1}^{j'_1} l_{i_2}^{j'_2} \end{bmatrix},$$

where

$$(2.15) \quad l_{i_1}^{j'_1} = \begin{bmatrix} l_{\alpha_1(1)}^{\beta'_1(1)} & 0 & 0 \\ l_{\alpha_1(2)}^{\beta'_1(1)} & l_{\alpha_1(2)}^{\beta'_1(2)} & 0 \\ l_{A_1(2)}^{\beta'_1(1)} & l_{A_1(2)}^{\beta'_1(2)} & l_{\alpha_1(1)}^{\beta'_1(1)} \end{bmatrix} \in G_1,$$

$$(2.16) \quad l_{i_1 i_2}^{j'_1} = \begin{bmatrix} l_{\alpha_1(1)\alpha_2(1)}^{\beta'_1(1)} & 0 & 0 \\ l_{\alpha_1(1)\alpha_2(2)}^{\beta'_1(1)} & l_{\alpha_1(1)\alpha_2(2)}^{\beta'_1(2)} & 0 \\ l_{\alpha_1(1)A_2(2)}^{\beta'_1(1)} & l_{\alpha_1(1)A_2(2)}^{\beta'_1(2)} & 0 \\ l_{\alpha_1(2)\alpha_2(2)}^{\beta'_1(1)} & l_{\alpha_1(2)\alpha_2(2)}^{\beta'_1(2)} & 0 \\ l_{\alpha_1(2)A_2(2)}^{\beta'_1(1)} & l_{\alpha_1(2)A_2(2)}^{\beta'_1(2)} & 0 \\ l_{A_1(2)A_2(2)}^{\beta'_1(1)} & l_{A_1(2)A_2(2)}^{\beta'_1(2)} & l_{\alpha_1(1)\alpha_2(2)}^{\beta'_1(1)} \end{bmatrix}.$$

Let G_1^2 be the subgroup of L_m^2 , consisting of all elements of the form (2.13) with corresponding matrix of the form (2.14) with (2.15) and (2.16). It can be verified that,

Proposition 2.2. *The group G_1^2 can be characterized as the subgroup of L_m^2 defined by all elements of L_m^2 which commute with F .*

Let $E_1^2(V_m)$ be the set of all the adapted bases at the different points of V_m and p the canonical mapping,

$$p : E_1^2(V_m) \longrightarrow V_m,$$

which associates with an adapted basis at x the point x itself. $E_1^2(V_m)$ is equipped with a structure of principal fibre bundle of basis V_m and structural group G_1^2 .

Conversely, we assume that the differentiable manifold V_m admits a G_1^2 -structure, where G_1^2 is the group of matrices of the form (2.14) with (2.15) and (2.16). Then, it can be defined on V_m , a tensor field F of type $(1, 1)$ and of rank $p + \binom{p+1}{2}$. F has (2.12) as components with respect to the adapted basis and satisfies the condition (2.2).

Thus, we have,

Theorem 2.1. *A necessary and sufficient condition for a differentiable manifold V_m to admit a G_1^2 -structure is that the structural group of the second order frame bundle $H^2(V_m)$ be reduced to the group G_1^2 .*

3. G_1^2 -Connections

Definition 3.1. Any infinitesimal connection ([18], [5]) defined on the principal bundle $E_1^2(V_m, G_m^2)$ is called a G_1^2 -connection.

We consider a covering of V_m by open neighborhoods endowed with local cross sections of $E_1^2(V_m)$. Any G_1^2 -connection may be defined in each neighborhood U by a local form π with values in the Lie algebra \mathcal{G}_1^2 of the group G_1^2 .

Hence, a G_1^2 -connection is represented by the element of the Lie algebra \mathcal{G}_1^2 ,

$$(3.1) \quad \pi_u = (\pi_{j_1}^{i_1}, \pi_{j_1 j_2}^{i_1}), \quad i_1, j_1, j_2 = 1, 2, \dots, m,$$

where the linear differential forms on U , $(\pi_{j_1}^{i_1}) \in R^m \otimes R^{m*}$ and $(\pi_{j_1 j_2}^{i_1}) \in R^m \otimes S^2(R^{m*})$ satisfy the relations,

$$(3.2) \quad \left\{ \begin{array}{l} \pi_{\alpha_1(1)}^{\beta_1(2)} = \pi_{\alpha_1(1)}^{B_1(2)} = \pi_{\alpha_1(2)}^{\beta_1(2)} = 0, \quad \pi_{\alpha_1(1)}^{\beta_1(1)} = \pi_{A_1(2)}^{B_1(2)}, \\ \pi_{\alpha_1(1)\alpha_2(1)}^{\beta_1(2)} = \pi_{\alpha_1(1)\alpha_2(1)}^{B_1(2)} = \pi_{\alpha_1(1)\alpha_2(2)}^{B_1(2)} = \pi_{\alpha_1(1)A_2(2)}^{B_1(2)} = \\ = \pi_{\alpha_1(2)\alpha_2(2)}^{B_1(2)} = \pi_{\alpha_1(2)A_2(2)}^{B_1(2)} = 0, \quad \pi_{\alpha_1(1)\alpha_2(1)}^{\beta_1(1)} = \pi_{A_1(2)A_2(2)}^{B_1(2)}. \end{array} \right.$$

It can be verified that,

Proposition 3.1. *With respect to a G₁²-connection, the absolute differential of the tensor F is zero.*

E₁²(V_m) may be considered as a sub-bundle of the fibre bundle H₁²(V_m) of 2-frames that is of bases of vector spaces {T_x²}_{x ∈ V_m} ([6]).

A G₁²-connection defines canonically a connection of order 2 ([9], [14]) on V_m with which it may be identified.

Conversely, let us consider a connection of order 2 and a covering of V_m by open neighborhoods equipped with local cross sections of E₁²(V_m). This connection may be defined on each neighborhood by a local form ω with values in the Lie algebra of L_m²,

$$(3.3) \quad \begin{aligned} \omega = (\omega_{j_1}^{i_1}, \omega_{j_1 j_2}^{i_1}), \quad i_1, j_1, j_2 = 1, 2, \dots, m, \quad \omega_{j_1}^{i_1} \in R^m \otimes R^{m*}, \\ \omega_{j_1 j_2}^{i_1} \in R^m \otimes S^2(R^{m*}) \text{ and } (\omega_{j_1}^{i_1}), (\omega_{j_1 j_2}^{i_1}) \\ \text{are local linear differential forms.} \end{aligned}$$

In order that the given connection may be identified with a G₁²-connection it is necessary and sufficient that the form (3.3) belongs in the Lie algebra of the structural group G₁² of E₁²(V_m). That is, comparing with (3.2),

Proposition 3.2. *In order that a connection of order 2 may be identified with a G₁²-connection, it is necessary and sufficient that the absolute differential of the tensor F is zero with respect to this connection.*

Given a G₁²-connection Y, the curvature form of this connection is the tensor 2-form, of adjoint type.

$$(3.4) \quad \Omega = \nabla \pi = d\pi + \pi \wedge \pi,$$

which is defined on E₁²(V_m) with values to G₁².

If we consider a covering of V_m by neighborhoods equipped with local cross sections of E₁²(V_m), then, Ω may be defined in each neighborhood U by a local form with values in the Lie algebra G₁²

$$(3.5) \quad \Omega_u = (\Omega_{j_1}^{i_1}, \Omega_{j_1 j_2}^{i_1}), \quad i, j_1, j_2 = 1, 2, \dots, m,$$

where $\Omega_{j_1}^{i_1} \in R^m \otimes R^{m*}$, $\Omega_{j_1 j_2}^{i_1} \in R^m \otimes S^2(R^{m*})$ and $(\Omega_{j_1}^{i_1}), (\Omega_{j_1 j_2}^{i_1})$ are linear differential forms on U.

It may be seen from (3.4).

$$(3.6) \quad \begin{cases} \Omega_{j_1}^{i_1} = d\pi_{j_1}^{i_1} + \pi_{k_1}^{i_1} \wedge \pi_{j_1}^{k_1}, \\ \Omega_{j_1 j_2}^{i_1} = d\pi_{j_1 j_2}^{i_1} + \pi_{k_1}^{i_1} \wedge \pi_{j_1 j_2}^{k_1} + \pi_{k_1 k_2}^{i_1} \wedge \pi_{j_1}^{k_1} \pi_{j_2}^{k_2}, \end{cases}$$

In particular, it can be verified that,

$$\begin{aligned}\Omega_{\alpha(1)}^{\alpha(1)} &= d\pi_{\alpha(1)}^{\alpha(1)}, \Omega_{\alpha(2)}^{\alpha(2)} = d\pi_{\alpha(2)}^{\alpha(2)}, \Omega_{A(2)}^{A(2)} = d\pi_{A(2)}^{A(2)}, \\ \Omega_{\alpha(1)\beta(1)}^{\alpha(1)} &= d\pi_{\alpha(1)\beta(1)}^{\alpha(1)}, \Omega_{\alpha(1)\beta(2)}^{\alpha(1)} = d\pi_{\alpha(1)\beta(2)}^{\alpha(1)}, \Omega_{\alpha(1)B(2)}^{\alpha(1)} = d\pi_{\alpha(1)B(2)}^{\alpha(1)}, \\ \Omega_{\alpha(2)\beta(2)}^{\alpha(2)} &= d\pi_{\alpha(2)\beta(2)}^{\alpha(2)}, \Omega_{\alpha(2)B(2)}^{\alpha(2)} = d\pi_{\alpha(2)B(2)}^{\alpha(2)}, \Omega_{A(2)B(2)}^{A(2)} = d\pi_{A(2)B(2)}^{A(2)}.\end{aligned}$$

Then,

$$\begin{aligned}\Psi = (\Omega_{\alpha(1)}^{\alpha(1)} = \Omega_{A(2)}^{A(2)}, \Omega_{\alpha(2)}^{\alpha(2)}, \Omega_{\alpha(1)\beta(1)}^{\alpha(1)} = \Omega_{A(2)B(2)}^{A(2)}, \Omega_{\alpha(1)\beta(2)}^{\alpha(1)} \\ \Omega_{\alpha(1)B(2)}^{\alpha(1)}, \Omega_{\alpha(2)\beta(2)}^{\alpha(2)}, \Omega_{\alpha(2)B(2)}^{\alpha(2)})\end{aligned}$$

is a closed 2-form on $E_1^2(V_m)$.

Definition 3.2. We call Ψ the characteristic form of the G_1^2 -connection Y .

Proposition 3.3. *The characteristic 2-forms of all the G_1^2 -connections have the same cohomology class of degree 2 (characteristic cohomology class of the G_1^2 -structure).*

4. *G*-structures of second order defined by linear operators satisfying algebraic relations

Using the way discussed previously (sections 1, 2) a generalization to the second order of the real almost product structure is given already in [6].

On the other hand, the definition (2.1) for a differentiable manifold V_{2m} , with rank $J = m + \binom{m+1}{2}$ gives a generalization of the almost tangent structure to the second order.

In this case, matrix (2.12) reduces to the form,

$$(4.1) \quad F = \begin{bmatrix} \begin{bmatrix} 0 & 0 \\ \delta_{\alpha_1}^{\beta_1} & 0 \end{bmatrix} & \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} & \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} & \begin{bmatrix} 0 & 0 & 0 \\ \delta_{\alpha_1}^{\beta_1} \delta_{\alpha_2}^{\beta_2} & 0 & 0 \end{bmatrix} \end{bmatrix},$$

and the matrix (2.14) to the form,

$$L = \begin{bmatrix} \begin{bmatrix} l_{\alpha_1}^{\beta_1} & 0 \\ l_{\alpha_1}^{\beta_1} & l_{\alpha_1}^{\beta_1} \end{bmatrix} & \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ \begin{bmatrix} l_{\alpha_1}^{\beta_1} l_{\alpha_2}^{\beta_2} & 0 \\ l_{\alpha_1}^{\beta_1} & 0 \end{bmatrix} & \begin{bmatrix} l_{\alpha_1}^{\beta_1} l_{\alpha_2}^{\beta_2} & 0 & 0 \\ l_{\alpha_1}^{\beta_1} l_{\alpha_2}^{\beta_2} & l_{\alpha_1}^{\beta_1} l_{\alpha_2}^{\beta_2} & 0 \\ l_{\alpha_1}^{\beta_1} l_{\alpha_2}^{\beta_2} & l_{\alpha_1}^{\beta_1} l_{\alpha_2}^{\beta_2} & l_{\alpha_1}^{\beta_1} l_{\alpha_2}^{\beta_2} \end{bmatrix} \end{bmatrix}$$

with the indices $\alpha = \alpha(1) = 1, 2, \dots, m$, $A = A(2) = m+1, \dots, 2m = m+\alpha$, and

$$\begin{bmatrix} l_{\alpha_1}^{\beta_1'} & 0 \\ l_{A_1}^{\beta_1'} & l_{\alpha_1}^{\beta_1'} \end{bmatrix} \in G_{mm}^m,$$

where G_{mm}^m is ([3], [12]) the structural group of the almost tangent structure with $l_{\alpha_1}^{\beta_1'} \in L_m$, $l_{A_1}^{\beta_1'} \in \text{End}(R^m, R^m)$.

Thus, G -structures on V_m of the first order defined by linear operators and satisfying some algebraic relations can be generalized to G -structure of the second order, defined by endomorphism,

$$J : T^2(V_m) \longrightarrow T^2(V_m)$$

and satisfying the same algebraic relations.

References

1. A. CARFAGNA D'ANDREA, R. DI FEBO MARINELLI, Su una generalizzazione delle structure quasi tangenti,, *Rend. Mat.* **2**, II série **VII** (1982), 257-267.
2. B. CENKI, On the G -structures of higher order, *Casopis Pěst. Mat.* **90** (1965), 26-32.
3. R.S. CLARK AND BRUCKHEIMER, Sur les structures presque tangentes, *Comptes Rendus* **251** (1960), 627-629.
4. C. COBDILLON, "Géométrie différentielle et mécanique analytique," Herman Paris, 1969.
5. L.A. CORDERO, C.T.J. DODSON, M. DE LEÓN, "Differential Geometry of frame bundles," Kluwer Academic Publ., Dordrecht, 1989.
6. D. DEMETROPOULOU-PSOMOPOULOU, A prolongation of the real almost product structure of a differentiable manifold, *Demonstratio Mathematica* **20**, 3-4 (1987), 423-439.
7. C. EHRESMANN, Les prolongements d'une variété différentiable. I Calcul des jets, prolongements principal, *Comptes Rendus* **233** (1951), 598-600.
8. C. EHRESMANN, Les prolongements d'une variété différentiable, Congr. Un. Matema. It. Taormina, 1951, 317-325.
9. C. EHRESMANN, Les prolongements d'une variété différentiable, V. Convariants différentiels et prolongements d'une structure infinitesimale, *Comptes Rendus* **234** (1952), 1424-1425.

10. C. EHRESMANN, Introduction à la théorie des structures infinitésimales et des pseudogroupes de Lie, *Coll. Intern. du C.N.R.S.*, Geom. Diff. Strasbourg (1953), 97–100.
11. H.A. ELIOPOULOS, On the connections and the holonomy group of certain G -structures, *Ann. di Math. pur. et appl. (IV)* **78** (1968), 1–12.
12. H.A. ELIOPOULOS, Structures presque tangents sur les variétés différentiables, *C.R. Acad. Sc. Paris* **255** (1962), 1563–1565.
13. J. GANCARZEWICZ, Liftings of functions and vector fields to natural bundles, *Dissertationes Mathematicae* **12** (1983).
14. S. KOBAYASHI, K. NOMIZU, “*Foundations of differential geometry*,” vol. I, Interscience Publ., New York, 1963.
15. J. LEHMANN-LEJEUNE, Sur l’intégrabilité de certaines G -structures, *C.R. Acad. Sc. Paris* **258** (1964), 5326–5329.
16. M. DE LEÓN, P.R. RODRIGUES, “*Generalized classical mechanics and field theory*,” North-Holland Mathematical Studies **112**, North-Holland, Amsterdam, 1985.
17. P. LIBERMANN, Pseudogroupes infinitésimaux. Pseudogroupes infinitésimaux de Lie., *C.R. de l’Acad. des Sc. de Paris* **246** (1958), 531–534.
18. A. LICHNEROWICZ, “*Théorie globale des connexiones et des groupes d’holonomie*,” Ed. Cremonese, Roma, 1955.
19. K. YANO, S. ISHIHARA, “*Tangent and cotangent bundles*,” Pure and Appl. Math. Ser. **16**, Marcel Dekker, New York, 1973.

Department of Mathematics
 Faculty of Sciences
 University of Thessaloniki
 54006 Thessaloniki
 GREECE

Primera versió rebuda el 20 de Novembre de 1990,
 darrera versió rebuda el 19 d’Abril de 1991