

# HEAT KERNEL LOWER GAUSSIAN ESTIMATES IN THE DOUBLING SETTING WITHOUT POINCARÉ INEQUALITY

SALAHADDINE BOUTAYEB

## Abstract

In the setting of a manifold with doubling property satisfying a Gaussian upper estimate of the heat kernel, one gives a characterization of the lower Gaussian estimate in terms of certain Hölder inequalities.

## CONTENTS

Introduction	457
1. Assumptions	461
2. Preliminaries	462
3. Hölder continuity of the heat kernel	462
4. Characterization of the lower gaussian estimate by some Hölder inequalities	466
4.1. First characterization	466
4.2. Second characterization	470
4.3. Further results	475
References	477

## Introduction

Let  $M$  be a connected non-compact Riemannian manifold without boundary, and let  $\Delta$  denote the Laplace-Beltrami operator on  $M$ . Consider the heat equation

$$\frac{\partial u}{\partial t} - \Delta u = 0,$$

where  $u = u(x, t)$ ,  $x \in M$ ,  $t > 0$ .

2000 *Mathematics Subject Classification*. Primary: 58J35; Secondary: 47D07, 46E35.

*Key words*. Heat kernel, Hölder inequalities.

The heat kernel  $p_t(x, y)$  is by definition the smallest positive fundamental solution to the heat equation on  $]0, +\infty[ \times M$ . In particular, the Cauchy problem

$$\begin{cases} \frac{\partial u}{\partial t} = \Delta u \\ u(0, \cdot) = f(\cdot), \end{cases}$$

where  $f$  is a bounded continuous function, is solved by

$$u(t, x) = \int_M p_t(x, y) f(y) dy.$$

In the Euclidean space  $\mathbb{R}^n$ , the heat kernel is given by the following well known formula:

$$p_t(x, y) = \frac{1}{(4\pi t)^{n/2}} \exp\left(-\frac{|x - y|^2}{4t}\right).$$

Denote by  $B(x, r)$  the ball of center  $x \in M$  and radius  $r > 0$  with respect to the Riemannian distance  $d$ , and by  $V(x, r)$  its Riemannian volume. If  $M$  is geodesically complete and has non-negative Ricci curvature, Li and Yau (see [14]) found that the heat kernel satisfies the two-sided Gaussian estimate:

$$\frac{c}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{ct}\right) \leq p_t(x, y) \leq \frac{C}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{Ct}\right),$$

$\forall t > 0, x, y \in M.$

One says that  $M$  satisfies the doubling property if

$$(D) \quad V(x, 2r) \leq CV(x, r), \quad \forall x \in M, r > 0.$$

A consequence of (D) is that there exists  $\nu > 0$  such that

$$(D_\nu) \quad \frac{V(x, s)}{V(x, r)} \leq C \left(\frac{s}{r}\right)^\nu, \quad \forall x \in M, s \geq r > 0.$$

Note that, if  $\text{Ric} M \geq 0$ , then  $M$  Satisfies the doubling property. We say that  $M$  admits the relative Faber-Krahn inequality if for any ball  $B(x, r) \subset M$  and any precompact open set  $\Omega \subset B(x, r)$

$$(FK) \quad \lambda_1(\Omega) \geq \frac{b}{r^2} \left(\frac{V(x, r)}{\mu(\Omega)}\right)^\nu,$$

where  $\lambda_1(\Omega)$  is the first Dirichlet eigenvalue for  $-\Delta$ ,  $b$  and  $\nu$  some positive constants.

On a geodesically complete manifold (see [9]), Grigor'yan proved that  $(FK)$  is equivalent to the Gaussian upper estimate

$$(UE) \quad p_t(x, y) \leq \frac{C}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{Ct}\right), \quad t > 0, x, y \in M,$$

in conjunction with the doubling property  $(D_\nu)$ .

Saloff-Coste (see [18], [19]) found that the two-sided Gaussian estimate of the heat kernel

$$\frac{c}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{ct}\right) \leq p_t(x, y) \leq \frac{C}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{Ct}\right),$$

$$t > 0, x, y \in M,$$

is equivalent to  $(D)$  and the 2-Poincaré inequalities

$$(P_2) \quad \int_{B(x, r)} |f(y) - f_r(x)|^2 d\mu(y) \leq Cr^2 \int_{B(x, Cr)} |\nabla f(y)|^2 d\mu(y),$$

for all  $f \in \mathcal{C}_0^\infty(M)$ ,  $x \in M$ ,  $r > 0$ , where  $f_r(x) = \frac{1}{V(x, r)} \int_{B(x, r)} f(y) d\mu(y)$ , see also [8].

Assuming that the Gaussian upper estimate holds, Coulhon and Ouhabaz gave some simple inequalities (other than Poincaré) that are necessary/sufficient to complete the two-sided Gaussian estimate.

In the case where the volume growth is polynomial, that is

$$V(x, r) \simeq r^D, \quad D > 0,$$

Coulhon (see [5]) proved that, if  $(UE)$  holds, then

$$(LE) \quad \frac{c}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{ct}\right) \leq p_t(x, y), \quad t > 0, x, y \in M,$$

is equivalent to the Sobolev type inequality

$$(S^D) \quad \frac{|f(x) - f(y)|}{d(x, y)^{\alpha - (D/p)}} \leq C \|\Delta^{\alpha/2} f\|_p, \quad \forall f \in \mathcal{C}_0^\infty(M), x, y \in M,$$

for  $p > 1$  and some  $\alpha > \frac{D}{p}$ .

In this polynomial setting and assuming  $(UE)$ , Ouhabaz (see [16]) had given previously a characterization of  $(LE)$  in terms of a weaker Gagliardo-Nirenberg type inequality:

$$(G^D) \quad \frac{|f(x) - f(y)|}{[d(x, y)]^{\alpha - (D/p)}} \leq C \|f\|_p^{1-\theta} \|\Delta^{\alpha/2\theta} f\|_p^\theta, \quad x, y \in M,$$

for  $p > 1$  and some  $\alpha > \frac{D}{p}$ ,  $\theta \in ]0, 1[$ .

In the more general case when the doubling condition  $(D_\nu)$  is satisfied, Coulhon proved that, if  $(UE)$  holds, then for any  $p > 1$  and  $\alpha > \frac{\nu}{p}$ :

$$|f(x) - f(y)| \leq C \frac{d^\alpha(x, y)}{V^{1/p}(x, d(x, y))} \|\Delta^{\alpha/2} f\|_p,$$

implies  $(LE)$ .

The aim of the present paper is to give similar necessary and sufficient conditions for the lower Gaussian estimate  $(LE)$  in the general setting when the volume is only doubling. More precisely, the necessary/sufficient conditions will be given by the following main results:

**Theorem.** Assume that  $(D_\nu)$  and  $(UE)$  holds. Then

$$(LE) \quad \frac{c}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{ct}\right) \leq p_t(x, y), \quad t > 0, x, y \in M,$$

implies that for all  $p$  large enough there are  $\alpha, \alpha' > \frac{\nu}{p}$  such that

$$(S) \quad \begin{cases} |f(x) - f(y)| \leq \frac{C}{V^{1/p}(x, d)} \max \left\{ d^\alpha \|\Delta^{\alpha/2} f\|_p, d^{\alpha'} \|\Delta^{\alpha'/2} f\|_p \right\} \\ \forall f \in \mathcal{D}, x, y \in M. \end{cases}$$

Conversely,  $(S)$  for any  $p > 1$  and  $\alpha, \alpha' > \frac{\nu}{p}$ , implies  $(LE)$ , where  $d$  denote to  $d(x, y)$ .

**Theorem.** Assume that  $(D_\nu)$  and  $(UE)$  holds. Then

$$(LE) \quad \frac{c}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{ct}\right) \leq p_t(x, y), \quad t > 0, x, y \in M,$$

implies for all  $p$  large enough, there are  $\alpha, \alpha' > \frac{\nu}{p}$  and  $\theta, \theta' \in ]0, 1[$  such that  $\alpha - \alpha' = \frac{\nu}{p}$ ,  $\frac{\alpha}{\theta} = \frac{\alpha'}{\theta'}$  and

$$(G) \quad \begin{cases} |f(x) - f(y)| \leq \frac{C}{V^{1/p}(x, d)} \\ \quad \times \max \left\{ d^\alpha \|f\|_p^{1-\theta} \|\Delta^{\alpha/2\theta} f\|_p^\theta, d^{\alpha'} \|f\|_p^{1-\theta'} \|\Delta^{\alpha'/2\theta'} f\|_p^{\theta'} \right\} \\ \forall f \in \mathcal{D}, x, y \in M. \end{cases}$$

Conversely,  $(G)$  for any  $p > 1$ ,  $\alpha, \alpha' > \frac{\nu}{p}$  and  $\theta, \theta' \in ]0, 1[$ , implies  $(LE)$ , where  $d$  denote to  $d(x, y)$ .

There are several classes of fractal spaces which satisfy the more general form of the heat kernel estimate:

$$\frac{c}{V(x, t^{1/w})} E_w(c, x, y, t) \leq p_t(x, y) \leq \frac{C}{V(x, t^{1/w})} E_w(C, x, y, t),$$

for  $t > 0$ ,  $x, y \in M$ , where

$$E_w(\lambda, x, y, t) = \exp \left( - \left( \frac{d^w(x, y)}{\lambda t} \right)^{1/(w-1)} \right), \quad \forall t, \lambda > 0, x, y \in M,$$

and  $w$  is a so-called escape time or random walk dimension (see [1], [2], [3], see also [10], [11], [12]). We are also able to write our characterizations of lower heat kernel estimate in this general form.

We choose to write this paper in a more general setting, when  $M$  is a metric measure space endowed with a symmetric Markov semigroup.

## 1. Assumptions

In the sequel, we shall place ourselves in a setting similar to the one in [5], [10], [13]: Let  $(M, d, \mu)$  be a metric measure space endowed with a symmetric Markov semigroup  $e^{-tA}$  on  $L^2(M, \mu)$  with a measurable kernel  $p_t$ , that is

$$e^{-tA}f(x) = \int_M p_t(x, y)f(y) d\mu(y), \quad t > 0, f \in L^2(M, \mu), \mu\text{-a.e. } x \in M.$$

The powers  $A^\alpha$ ,  $\alpha > 0$ , of the operator  $A$  are densely defined on  $L^p(M, \mu)$ ,  $1 \leq p < +\infty$ , we denote by  $\mathcal{D}_p(A^\alpha)$  their domain. Background and more information about this functional setting can be found in [17] and references therein. For simplicity, we always write  $\mathcal{D}$  for the space  $\mathcal{D}_p(A^\alpha)$  required by the context. We assume that  $\|e^{-tA}f\|_p \rightarrow 0$  as  $t \rightarrow +\infty$ , for all  $f \in \mathcal{D}$  and  $1 \leq p < +\infty$ .

We shall assume that for all  $x \in M$  and  $r > 0$ ,  $0 < V(x, r) < +\infty$  and that the metric space  $(M, d)$  satisfies the chain condition: there exists  $C > 0$  such that, for all  $x, y \in M$ , for any  $n \in \mathbb{N}^*$ , there exists a sequence  $\{x_i\}_{i=0}^n$  of points in  $M$  such that  $x_0 = x$ ,  $x_n = y$  and

$$d(x_i, x_{i+1}) \leq C \frac{d(x, y)}{n}, \quad \forall i = 0, \dots, n-1.$$

We say that such a sequence is a chain connecting  $x$  and  $y$ .

Note that these assumptions are well satisfied in the setting of the introduction when  $M$  is a connected non-compact complete Riemannian manifold. They will be standing assumptions in this paper and we will refer to them by writing: let  $(M, d, \mu, A)$  be as above. We recall that a symmetric Markov semigroup on  $L^2(M, \mu)$  is analytic on  $L^p(M, \mu)$  (see [21]).

## 2. Preliminaries

### Notation:

Consider a parameter  $w \geq 2$ ,

$$(UE_w) \quad p_t(x, y) \leq \frac{C}{V(x, t^{1/w})} E_w(C, x, y, t), \quad \forall t > 0, x, y \in M.$$

$$(LE_w) \quad \frac{c}{V(x, t^{1/w})} E_w(c, x, y, t) \leq p_t(x, y), \quad \forall t > 0, x, y \in M.$$

$$(DLE_w) \quad \frac{c}{V(x, t^{1/w})} \leq p_t(x, x), \quad \forall t > 0, x \in M.$$

$$(LY_w) \quad \frac{c}{V(x, t^{1/w})} E_w(c, x, y, t) \leq p_t(x, y) \leq \frac{C}{V(x, t^{1/w})} E_w(C, x, y, t),$$

for all  $t > 0$  and  $x, y \in M$ .

Note that  $(UE_2) = (UE)$  and similarly for the others  $w$ -estimations. For all  $p > 1$ , we denote by  $p'$  the conjugate exponent of  $p$ ,  $\frac{1}{p} + \frac{1}{p'} = 1$ . Letters  $c, C, C'$  are normally used to denote unimportant positive constants, whose values may change at each occurrence. In the sequel, for the sake of simplicity, we sometimes denote  $d(x, y)$  by  $d$ .

We recall the following proposition. for the proof, in the case  $w = 2$ , see [4, Lemma 1, p. 224], [7, §6] or [20, Theorem 4.2.8], [13, §3.3]; for the general case  $w \geq 2$ , see [16, §4] or [5, §3].

**Proposition 1.** *Assume that  $(M, d, \mu)$  satisfies the doubling property.*

*If  $(UE_w)$  holds, then  $(DLE_w)$  holds.*

The following lemma follows from [10, Corollary 3.5], [13, Lemma 5.1], see also [5, p. 801].

**Lemma 2.** *Assume that  $(M, d, \mu)$  satisfies the doubling property. If  $(UE_w)$  holds and there exist  $a, c > 0$  such that*

$$\frac{c}{V(x, t^{1/w})} \leq p_t(x, y), \text{ for all } t > 0, x, y \in M \text{ with } d(x, y) \leq at^{1/w},$$

*then  $(LE_w)$  holds.*

## 3. Hölder continuity of the heat kernel

At first, let us recall the following proposition, which follows from [19, Proposition 3.2] in the case  $w = 2$  and [13, §5.3] for the general case.

**Proposition 3.** Assume that  $(M, d, \mu)$  satisfies the doubling property. If  $(LY_w)$  holds, then there exists  $\eta \in ]0, 1[$  and  $C > 0$  such that

$$|p_t(x, z) - p_t(y, z)| \leq \frac{C}{V(z, t^{1/w})} \left( \frac{d(x, y)}{t^{1/w}} \right)^\eta,$$

for all  $t > 0$  and  $\mu$ -a.e.  $x, y, z \in M$ .

In the next result, we give an estimation with a Gaussian factor.

**Proposition 4.** Assume that  $(M, d, \mu)$  satisfies the doubling condition  $(D_\nu)$ . If  $(LY_w)$  holds, then there exists  $\eta \in ]0, 1[$  and  $C > 0$  such that for all  $p > 1$ :

$$|p_t(x, z) - p_t(y, z)| \leq C \frac{(E_w(C, x, z, t) + E_w(C, y, z, t))^{1/p}}{V^{1/p}(x, d)V^{1/p'}(z, t^{1/w})} \begin{cases} \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu}{p}} & \text{if } d \geq t^{1/w} \\ \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'}} & \text{if not,} \end{cases}$$

for all  $t > 0$  and  $\mu$ -a.e.  $x, y, z \in M$ , where  $d$  denote to  $d(x, y)$ .

*Proof:* Let  $x, y, z \in M$ ,  $p > 1$  and  $p'$  its conjugate exponent, one may write

$$|p_t(x, z) - p_t(y, z)| \leq |p_t(x, z) - p_t(y, z)|^{1/p'} \cdot (p_t(x, z) + p_t(y, z))^{1/p}.$$

According to Proposition 3 and  $(UE_w)$ , it follows

$$(1) \quad |p_t(x, z) - p_t(y, z)| \leq \frac{C}{V^{1/p'}(z, t^{1/w})} \left( \frac{d(x, y)}{t^{1/w}} \right)^{\frac{\eta}{p'}} \times \left( \frac{1}{V(x, t^{1/w})} E_w(C, x, z, t) + \frac{1}{V(y, t^{1/w})} E_w(C, y, z, t) \right)^{1/p}.$$

Denote to  $d(x, y)$  by  $d$ . Since  $V(x, d) \leq V(y, 2d)$  and  $V(y, 2d) \leq CV(y, d)$ , then  $V(x, d) \leq CV(y, d)$ .

Assume that  $d \geq t^{1/w}$ . From  $(D_\nu)$ , one has

$$\frac{1}{V(x, t^{1/w})} \leq \frac{C}{V(x, d)} \left( \frac{d}{t^{1/w}} \right)^\nu$$

and

$$\frac{1}{V(y, t^{1/w})} \leq \frac{C}{V(y, d)} \left( \frac{d}{t^{1/w}} \right)^\nu \leq \frac{C'}{V(x, d)} \left( \frac{d}{t^{1/w}} \right)^\nu.$$

Then, (1) yields

$$|p_t(x, z) - p_t(y, z)| \leq \frac{C}{V^{1/p}(x, d)} \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\kappa}{p}} \frac{1}{V^{1/p'}(z, t^{1/w})} \\ \times [E_w(C, x, z, t) + E_w(C, y, z, t)]^{1/p}.$$

If  $d \leq t^{1/w}$ . One has also

$$\frac{1}{V(y, t^{1/w})} \leq \frac{1}{V(y, d)} \leq \frac{C}{V(x, d)}.$$

Analogously, by (1) it follows that

$$|p_t(x, z) - p_t(y, z)| \leq \frac{C}{V^{1/p}(x, d)} \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'}} \frac{1}{V^{1/p'}(z, t^{1/w})} \\ \times [E_w(C, x, z, t) + E_w(C, y, z, t)]^{1/p}. \quad \square$$

Now, we derive an oscillation estimate for the semigroup  $(e^{-tA})_{t \geq 0}$ .

**Proposition 5.** *Assume that  $(M, d, \mu)$  satisfies the doubling condition  $(D_\nu)$ . If  $(LY_w)$  holds, then for all  $p > 1$  there exists  $C > 0$  such that*

$$|e^{-tA}f(x) - e^{-tA}f(y)| \\ \leq \frac{C\|f\|_p}{V^{1/p}(x, d(x, y))} \begin{cases} \left( \frac{d(x, y)}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\kappa}{p}} & \text{if } d(x, y) \geq t^{1/w} \\ \left( \frac{d(x, y)}{t^{1/w}} \right)^{\frac{\eta}{p'}} & \text{if not,} \end{cases}$$

for all  $t > 0$ ,  $f \in \mathcal{D}$  and  $\mu$ -a.e.  $x, y \in M$ .

*Proof:* From the definition of the heat kernel, one can write

$$|e^{-tA}f(x) - e^{-tA}f(y)| \leq \int_M |(p_t(x, z) - p_t(y, z))f(z)| d\mu(z),$$

for all  $t > 0$ ,  $f \in \mathcal{D}$ ,  $x, y \in M$ .



Assume that  $d \geq t^{1/w}$ . From Proposition 4, it follows

$$\begin{aligned} \int_M |(p_t(x, z) - p_t(y, z)) f(z)| d\mu(z) &\leq \frac{C}{V^{1/p}(x, d)} \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu}{p}} \\ &\quad \times \int_M \frac{|f(z)|}{V^{1/p'}(z, t^{1/w})} (E_w(C, x, z, t) + E_w(C, y, z, t))^{1/p} d\mu(z), \end{aligned}$$

and by using Hölder inequality, we obtain

$$\begin{aligned} (2) \quad \int_M |(p_t(x, z) - p_t(y, z)) f(z)| d\mu(z) &\leq \frac{C}{V^{1/p}(x, d)} \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu}{p}} \|f\|_p \\ &\quad \times \left( \int_M \frac{1}{V(z, t^{1/w})} (E_w(C, x, z, t) + E_w(C, y, z, t))^{p'/p} d\mu(z) \right)^{1/p'}. \end{aligned}$$

Let us prove that for all  $\gamma > 0$ , there is  $C > 0$ , such that for all  $t > 0$ .

$$(3) \quad \int_M \frac{1}{V(z, t^{1/w})} (E_w(C, x, z, t) + E_w(C, y, z, t))^\gamma d\mu(z) \leq C.$$

Indeed, since  $(a + b)^\gamma \leq C_\gamma(a^\gamma + b^\gamma)$ , for all  $a \geq 0$  and  $b \geq 0$ , then

$$\begin{aligned} &(E_w(C, x, z, t) + E_w(C, y, z, t))^\gamma \\ &\leq C (E_w^\gamma(C, x, z, t) + E_w^\gamma(C, y, z, t)) \\ &\leq C (E_w(C\gamma^{1-w}, x, z, t) + E_w(C\gamma^{1-w}, y, z, t)). \end{aligned}$$

Let  $c$  be the constant in  $(LE_w)$ . Set  $a = \frac{c}{C\gamma^{1-w}}$  and  $s = \frac{t}{a}$ , then

$$\begin{aligned} E_w(C\gamma^{1-w}, x, z, t) &= E_w(c/a, x, z, as) \\ &= E_w(c, x, z, s). \end{aligned}$$

Therefore

$$\begin{aligned} &\int_M \frac{1}{V(z, t^{1/w})} (E_w(C, x, z, t) + E_w(C, y, z, t))^\gamma d\mu(z) \\ &\leq \int_M \frac{C}{V(z, (as)^{1/w})} (E_w(c, x, z, s) + E_w(c, y, z, s)) d\mu(z). \end{aligned}$$

By  $(D_\nu)$ , one has  $V(z, \sqrt{as}) \simeq V(z, \sqrt{s})$ ,  $s > 0$ . On the other hand, according to  $(LE_w)$  and since  $(e^{-tA})_{t \geq 0}$  is Markovian, one has

$$\int_M \frac{1}{V(z, \sqrt{s})} E_w(c, x, z, t) d\mu(z) \leq \int_M p_s(x, z) d\mu(z) \leq 1,$$

for all  $s > 0$  and  $x \in M$ . Therefore, we deduce (3).

By choosing  $\gamma = p'/p$ , (2) yields

$$|e^{-tA}f(x) - e^{-tA}f(y)| \leq \frac{C}{V^{1/p}(x,d)} \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu}{p}} \|f\|_p.$$

If  $d \leq t^{1/w}$ . Using Proposition 4 and arguing as before, it follows

$$|e^{-tA}f(x) - e^{-tA}f(y)| \leq \frac{C}{V^{1/p}(x,d)} \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'}} \|f\|_p. \quad \square$$

#### 4. Characterization of the lower gaussian estimate by some Hölder inequalities

##### 4.1. First characterization.

**Theorem 6.** *Let  $(M, d, \mu, A)$  as above satisfy the doubling condition  $(D_\nu)$ . Let  $w \geq 2$ . Assume that  $(UE_w)$  holds. Then*

$$(LE_w) \quad \frac{c}{V(x, t^{1/w})} E_w(c, x, y, t) \leq p_t(x, y), \quad \forall t > 0, \forall x, y \in M,$$

*implies that for all  $p$  large enough there are  $\alpha, \alpha' > \frac{\nu}{p}$  such that*

$$(S_w) \quad \begin{cases} |f(x) - f(y)| \leq \frac{C}{V^{1/p}(x,d)} \max \left\{ d^\alpha \|A^{\alpha/w} f\|_p, d^{\alpha'} \|A^{\alpha'/w} f\|_p \right\} \\ \forall f \in \mathcal{D}, x, y \in M. \end{cases}$$

*Conversely,  $(S_w)$  for any  $p > 1$  and  $\alpha, \alpha' > \frac{\nu}{p}$ , implies  $(LE_w)$ , where  $d$  denote to  $d(x, y)$ .*

*Proof:*

( $\Rightarrow$ ) For all  $f \in \mathcal{D}$ ,  $t > 0$ , we know that  $\frac{\partial^k}{\partial t^k} e^{-tA} f = A^k e^{-tA} f$  for all  $k \in \mathbb{N}$ . Then

$$f = \sum_{i=0}^{k-1} \frac{t^i}{i!} A^i e^{-tA} f + \frac{1}{(k-1)!} \int_0^t s^{k-1} A^k e^{-sA} f ds, \quad \forall k \geq 1.$$

Since  $(e^{-tA})_{t>0}$  is analytic, then for all  $p > 1$

$$\|A^i e^{-tA} f\|_p = \|A^i e^{-(t/2)A} e^{-(t/2)A} f\|_p \leq C t^{-i} \|e^{-(t/2)A} f\|_p.$$

On the other hand,  $e^{-tA} f \rightarrow 0$ ,  $\forall f \in \mathcal{D}$  when  $t \rightarrow +\infty$ , then for all  $k \geq 1$ :

$$f = \frac{1}{(k-1)!} \int_0^{+\infty} t^{k-1} A^k e^{-tA} f dt, \quad \forall f \in \mathcal{D}.$$

Hence

$$|f(x) - f(y)| \leq C_k \int_0^{+\infty} t^{k-1} |A^k e^{-tA} f(x) - A^k e^{-tA} f(y)| ds,$$

for all  $k \geq 1$ , and  $f \in \mathcal{D}$ .

By rewriting

$$\begin{aligned} & |A^k e^{-tA} f(x) - A^k e^{-tA} f(y)| \\ &= |e^{-(t/2)A} A^k e^{-(t/2)A} f(x) - e^{-(t/2)A} A^k e^{-(t/2)A} f(y)| \end{aligned}$$

and applying Proposition 5 to  $A^k e^{-(t/2)A} f$ , it follows

$$\begin{aligned} & |e^{-tA} f(x) - e^{-tA} f(y)| \\ & \leq \frac{C \|A^k e^{-(t/2)A} f\|_p}{V^{1/p}(x, d(x, y))} \begin{cases} \left( \frac{d(x, y)}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu}{p}} & \text{if } d(x, y) \geq t^{1/w} \\ \left( \frac{d(x, y)}{t^{1/w}} \right)^{\frac{\eta}{p'}} & \text{if not,} \end{cases} \end{aligned}$$

for all  $t > 0$  and  $\mu$ -a.e.  $x, y \in M$ . Then

$$\begin{aligned} |f(x) - f(y)| & \leq \frac{C}{V^{1/p}(x, d)} \left( d^{\frac{\eta}{p'} + \frac{\nu}{p}} \int_0^{d^w} t^{k - \frac{1}{w}(\frac{\eta}{p'} + \frac{\nu}{p}) - 1} \|A^k e^{-(t/2)A} f\|_p dt \right. \\ & \quad \left. + d^{\frac{\eta}{p'}} \int_{d^w}^{+\infty} t^{k - \frac{\eta}{wp'} - 1} \|A^k e^{-(t/2)A} f\|_p dt \right). \end{aligned}$$

Now, we choose  $k > \frac{1}{w}(\frac{\eta}{p'} + \frac{\nu}{p})$  and let  $\delta \in ]0, \frac{p}{p'}\eta[$ .

One can write

$$\|A^k e^{-(t/2)A} f\|_p = \|A^{k - \frac{1}{w}(\frac{\eta}{p'} + \frac{\nu + \delta}{p})} e^{-(t/2)A} (A^{\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu + \delta}{p})} f)\|_p,$$

then by analyticity, one has

$$\|A^k e^{-(t/2)A} f\|_p \leq C_1 t^{\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu + \delta}{p}) - k} \|A^{\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu + \delta}{p})} f\|_p.$$

Similarly, one also has

$$\|A^k e^{-(t/2)A} f\|_p \leq C_2 t^{\frac{1}{w}(\frac{\eta}{p'} - \frac{\delta}{p}) - k} \|A^{\frac{1}{w}(\frac{\eta}{p'} - \frac{\delta}{p})} f\|_p.$$

Then

$$|f(x) - f(y)| \leq \frac{C}{V^{1/p}(x, d)} \left( d^{\frac{\eta}{p'} + \frac{\nu}{p}} \int_0^{d^w} t^{-\frac{\delta}{wp} - 1} \|A^{\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu + \delta}{p})} f\|_p dt \right. \\ \left. + d^{\frac{\eta}{p'}} \int_{d^w}^{+\infty} t^{-\frac{\delta}{wp} - 1} \|A^{\frac{1}{w}(\frac{\eta}{p'} - \frac{\delta}{p})} f\|_p dt \right).$$

Therefore

$$|f(x) - f(y)| \leq \frac{C}{V^{1/p}(x, d)} (d^\alpha \|A^{\alpha/w} f\|_p + d^{\alpha'} \|A^{\alpha'/w} f\|_p) \\ \leq \frac{C}{V^{1/p}(x, d)} \max\{d^\alpha \|A^{\alpha/w} f\|_p, d^{\alpha'} \|A^{\alpha'/w} f\|_p\},$$

where  $\alpha = \frac{\eta}{p'} + \frac{\nu + \delta}{p} > \frac{\nu}{p}$ ,  $\alpha' = \frac{\eta}{p'} - \frac{\delta}{p} > 0$  and  $\delta \in ]0, \frac{p}{p'}\eta[$ .

For  $p > \frac{\nu + \eta}{\eta}$ , there is  $\delta_0 > 0$  such that  $p > \frac{\nu + \eta + \delta}{\eta}$ ,  $\forall \delta \in ]0, \delta_0[$ .

Therefore for  $\delta \in ]0, \delta_0[$ , one has  $\eta > \frac{\nu + \eta + \delta}{p}$ , then  $\eta(1 - \frac{1}{p}) > \frac{\nu + \delta}{p}$ , hence  $\alpha' = \frac{\eta}{p'} - \frac{\delta}{p} > \frac{\nu}{p}$ .

( $\Leftarrow$ ) Let  $p > 1$ , assume that for some  $\alpha$  and  $\alpha' > \frac{\nu}{p}$ :

$$(S_w) \quad \begin{cases} |f(x) - f(y)| \leq \frac{C}{V^{1/p}(x, d)} \max\{d^\alpha \|A^{\alpha/w} f\|_p, d^{\alpha'} \|A^{\alpha'/w} f\|_p\} \\ \forall f \in \mathcal{D}, x, y \in M. \end{cases}$$

Let  $z \in M$ , by analyticity of  $(e^{-tA})_{t>0}$ ,  $p_t(\cdot, z)$  belong to  $\mathcal{D}_p(A^{\alpha/w})$ . Then by choosing  $f = p_t(\cdot, z)$  in  $(S_w)$ , one has for all  $x, y \in M$ :

$$|p_t(x, z) - p_t(y, z)| \\ \leq \frac{C}{V^{1/p}(x, d)} \max\{d^\alpha \|A^{\alpha/w} p_t(\cdot, z)\|_p, d^{\alpha'} \|A^{\alpha'/w} p_t(\cdot, z)\|_p\}.$$

By analyticity, we obtain

$$\|A^{\alpha/w} p_t(\cdot, z)\|_p = \|A^{\alpha/w} e^{-(t/2)A} p_{t/2}(\cdot, z)\|_p \leq C t^{-\alpha/w} \|p_{t/2}(\cdot, z)\|_p.$$

Since  $(e^{-tA})_{t>0}$  is symmetric Markovian, then  $\|p_{t/2}(\cdot, z)\|_1 \leq 1$ ,  $\forall t > 0$ .

Then from  $(UE_w)$  and Hölder inequality, it follows

$$\|p_t(\cdot, z)\|_p \leq \frac{C}{[V(z, t^{1/w})]^{1 - \frac{1}{p}}}, \quad \forall t > 0,$$

and by  $(D_\nu)$ , one has

$$\|p_{t/2}(\cdot, z)\|_p \leq \frac{C}{[V(z, t^{1/w})]^{1-\frac{1}{p}}},$$

then

$$\|A^{\alpha/w} p_t(\cdot, z)\|_p \leq C \frac{t^{-\alpha/w}}{[V(z, t^{1/w})]^{1-\frac{1}{p}}}.$$

Therefore

$$\begin{aligned} & |p_t(x, z) - p_t(y, z)| \\ & \leq \frac{C}{V^{1/p}(x, d)} \max \left\{ \left( \frac{d}{t^{1/w}} \right)^\alpha \frac{1}{[V(z, t^{1/w})]^{1-\frac{1}{p}}}, \left( \frac{d}{t^{1/w}} \right)^{\alpha'} \frac{1}{[V(z, t^{1/w})]^{1-\frac{1}{p}}} \right\} \\ & \leq \frac{C}{V(z, t^{1/w})} \left( \frac{V(z, t^{1/w})}{V(x, d)} \right)^{1/p} \max \left\{ \left( \frac{d}{t^{1/w}} \right)^\alpha, \left( \frac{d}{t^{1/w}} \right)^{\alpha'} \right\}. \end{aligned}$$

By Proposition 1,  $(UE_w)$  and  $(D_\nu)$  yield

$$(DLE_w) \quad \frac{c}{V(z, t^{1/w})} \leq p_t(z, z), \quad \forall z \in M.$$

Thus

$$|p_t(x, z) - p_t(y, z)| \leq C \left( \frac{V(z, t^{1/w})}{V(x, d)} \right)^{1/p} \max \left\{ \left( \frac{d}{t^{1/w}} \right)^\alpha, \left( \frac{d}{t^{1/w}} \right)^{\alpha'} \right\} p_t(z, z).$$

For  $z = x$ , one has

$$|p_t(x, x) - p_t(y, x)| \leq C \left( \frac{V(x, t^{1/w})}{V(x, d)} \right)^{1/p} \max \left\{ \left( \frac{d}{t^{1/w}} \right)^\alpha, \left( \frac{d}{t^{1/w}} \right)^{\alpha'} \right\} p_t(x, x).$$

If  $d \leq t^{1/w}$ ,  $(D_\nu)$  yields

$$|p_t(x, x) - p_t(y, x)| \leq C \max \left\{ \left( \frac{d}{t^{1/w}} \right)^{\alpha-\frac{\nu}{p}}, \left( \frac{d}{t^{1/w}} \right)^{\alpha'-\frac{\nu}{p}} \right\} p_t(x, x).$$

Since  $\alpha$  and  $\alpha' > \frac{\nu}{p}$ , then for a small enough  $d \leq at^{1/w}$

$$|p_t(x, x) - p_t(y, x)| \leq \frac{1}{2} p_t(x, x),$$

consequently

$$p_t(x, y) \geq \frac{1}{2} p_t(x, x) \geq \frac{C}{V(x, t^{1/w})},$$

$$\forall x, y \in M, t > 0, \text{ such that } d(x, y) \leq at^{1/w}.$$

Therefore, by Lemma 2, one concludes that  $(LE_w)$  is satisfied.  $\square$

#### 4.2. Second characterization.

**Theorem 7.** *Let  $(M, d, \mu, A)$  as above satisfy the doubling condition  $(D_\nu)$ . Let  $w \geq 2$ . Assume that  $(UE_w)$  holds. Then*

$$(LE_w) \quad \frac{c}{V(x, t^{1/w})} E_w(c, x, y, t) \leq p_t(x, y), \quad \forall t > 0, x, y \in M,$$

*implies for all  $p$  large enough, there are  $\alpha, \alpha' > \frac{\nu}{p}$  and  $\theta, \theta' \in ]0, 1[$  such that  $\alpha - \alpha' = \frac{\nu}{p}$ ,  $\frac{\alpha}{\theta} = \frac{\alpha'}{\theta'}$  and*

$$(G_w) \quad \begin{cases} |f(x) - f(y)| \leq \frac{C}{V^{1/p}(x, d)} \\ \quad \times \max \left\{ d^\alpha \|f\|_p^{1-\theta} \|A^{\alpha/w\theta} f\|_p^\theta, d^{\alpha'} \|f\|_p^{1-\theta'} \|A^{\alpha'/w\theta'} f\|_p^{\theta'} \right\} \\ \forall f \in \mathcal{D}, x, y \in M. \end{cases}$$

*Conversely,  $(G_w)$  for any  $p > 1$ ,  $\alpha, \alpha' > \frac{\nu}{p}$  and  $\theta, \theta' \in ]0, 1[$ , implies  $(LE_w)$ , where  $d$  denote to  $d(x, y)$ .*

*Proof:*

$(\Rightarrow)$  Let  $\beta > \frac{\eta}{p'} + \frac{\nu}{p}$ . Firstly assume that  $\beta < w$ . For all  $f \in \mathcal{D}$ , we know that

$$f = e^{-tA} f + \int_0^t A e^{-sA} f ds, \quad \forall t > 0,$$

then

$$(4) \quad |f(x) - f(y)| \leq |e^{-tA} f(x) - e^{-tA} f(y)| + \int_0^t |A e^{-sA} f(x) - A e^{-sA} f(y)| ds$$

and

$$\begin{aligned} & \int_0^t |A e^{-sA} f(x) - A e^{-sA} f(y)| ds \\ &= \int_0^t |e^{-(s/2)A} A e^{-(s/2)A} f(x) - e^{-(s/2)A} A e^{-(s/2)A} f(y)| ds. \end{aligned}$$

Assume that  $d \geq t^{1/w}$ . From Proposition 5, we obtain

$$\begin{aligned} & \int_0^t |e^{-(s/2)A} A e^{-(s/2)A} f(x) - e^{-(s/2)A} A e^{-(s/2)A} f(y)| ds \\ & \leq \frac{C}{V^{1/p}(x, d)} d^{\frac{\eta}{p'} + \frac{\nu}{p}} \int_0^t s^{-\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu}{p})} \|A e^{-(s/2)A} f\|_p ds. \end{aligned}$$

Since  $(e^{-tA})_{t>0}$  is analytic, then  $\|A e^{-(s/2)A} f\|_p \leq C s^{\frac{\beta}{w}-1} \|A^{\beta/w} f\|_p$ . Hence

$$\begin{aligned} & \int_0^t s^{-\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu}{p})} \|A e^{-(s/2)A} f\|_p ds \leq C \|A^{\beta/w} f\|_p \int_0^t s^{-\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu}{p}) + \frac{\beta}{w} - 1} ds \\ & = C t^{-\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu}{p}) + \frac{\beta}{w}} \|A^{\beta/w} f\|_p. \end{aligned}$$

Thus

$$\begin{aligned} (5) \quad & \int_0^t |A e^{-sA} f(x) - A e^{-sA} f(y)| ds \\ & \leq \frac{C}{V^{1/p}(x, d)} d^{\frac{\eta}{p'} + \frac{\nu}{p}} t^{-\frac{1}{w}(\frac{\eta}{p'} + \frac{\nu}{p}) + \frac{\beta}{w}} \|A^{\beta/w} f\|_p. \end{aligned}$$

If  $d \leq t^{1/w}$ .

$$\begin{aligned} & \int_0^t |A e^{-sA} f(x) - A e^{-sA} f(y)| ds = \int_0^{d^w} |A e^{-sA} f(x) - A e^{-sA} f(y)| ds \\ & \quad + \int_{d^w}^t |e^{-(s/2)A} A e^{-(s/2)A} f(x) - e^{-(s/2)A} A e^{-(s/2)A} f(y)| ds. \end{aligned}$$

From (5) and Proposition 5, we obtain

$$\begin{aligned} & \int_0^t |A e^{-sA} f(x) - A e^{-sA} f(y)| ds \\ & \leq \frac{C}{V^{1/p}(x, d)} d^\beta \|A^{\beta/w} f\|_p + \frac{C}{V^{1/p}(x, d)} d^{\frac{\eta}{p'}} \int_{d^w}^t s^{-\frac{\eta}{wp'}} \|A e^{-(s/2)A} f\|_p ds. \end{aligned}$$

Since  $(e^{-tA})_{t>0}$  is analytic, then

$$\begin{aligned} & \int_{d^w}^t s^{-\frac{\eta}{wp'}} \|A e^{-(s/2)A} f\|_p ds \leq C \|A^{\beta/w} f\|_p \int_{d^w}^t s^{-\frac{\eta}{wp'} + \frac{\beta}{w} - 1} ds \\ & = C (t^{-\frac{\eta}{wp'} + \frac{\beta}{w}} - d^{-\frac{\eta}{p'} + \beta}) \|A^{\beta/w} f\|_p. \end{aligned}$$

Therefore

$$\begin{aligned}
 & \int_0^t |Ae^{-sA}f(x) - Ae^{-sA}f(y)| ds \\
 & \leq \frac{C}{V^{1/p}(x,d)} d^\beta \|A^{\beta/w}f\|_p + \frac{C}{V^{1/p}(x,d)} (d^{\frac{\eta}{p'}} t^{-\frac{\eta}{wp'} + \frac{\beta}{w}} - d^\beta) \|A^{\beta/w}f\|_p \\
 (6) \quad & \leq \frac{C}{V^{1/p}(x,d)} d^{\frac{\eta}{p'}} t^{-\frac{\eta}{wp'} + \frac{\beta}{w}} \|A^{\beta/w}f\|_p.
 \end{aligned}$$

Hence by (4), Proposition 5, (5) and (6), one has for all  $t > 0$

$$|f(x) - f(y)| \leq \frac{C}{V^{1/p}(x,d)} \times (*),$$

where

$$\begin{aligned}
 (*) = \max \left\{ \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu}{p}} (\|f\|_p + t^{\beta/w} \|A^{\beta/w}f\|_p), \right. \\
 \left. \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'}} (\|f\|_p + t^{\beta/w} \|A^{\beta/w}f\|_p) \right\}.
 \end{aligned}$$

By choosing  $t = (\frac{\|f\|_p}{\|A^{\beta/w}f\|_p})^{w/\beta}$ , one obtains

$$\begin{aligned}
 |f(x) - f(y)| & \leq \frac{C}{V^{1/p}(x,d)} \\
 & \times \max \left\{ d^{\frac{\eta}{p'} + \frac{\nu}{p}} \|f\|_p^{1-\theta} \|A^{\beta/w}f\|_p^\theta, d^{\frac{\eta}{p'}} \|f\|_p^{1-\theta'} \|A^{\beta/w}f\|_p^{\theta'} \right\},
 \end{aligned}$$

where  $\theta = \frac{1}{\beta}(\frac{\eta}{p'} + \frac{\nu}{p})$  and  $\theta' = \frac{\eta}{\beta p'}$ .

Now, if  $\beta \geq w$ . Fix  $k > \frac{\beta}{w}$ , we know that

$$f = \sum_{i=0}^{k-1} \frac{t^i}{i!} A^i e^{-tA} f + \frac{1}{(k-1)!} \int_0^t s^{k-1} A^k e^{-sA} f ds, \quad \forall t > 0.$$

By Proposition 5 and the analyticity, it yields, for all  $i = 0, \dots, k-1$ :

$$\begin{aligned}
 |A^i e^{-tA}f(x) - A^i e^{-tA}f(y)| & \leq \frac{C}{V^{1/p}(x,d(x,y))} \|A^i e^{-(t/2)A}f\|_p \times (**) \\
 & \leq \frac{C t^{-i}}{V^{1/p}(x,d(x,y))} \|f\|_p \times (**)
 \end{aligned}$$



for all  $t > 0$ ,  $f \in \mathcal{D}$  and  $\mu$ -a.e.  $x, y \in M$ . where

$$(**) = \begin{cases} \left( \frac{d(x, y)}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu}{p}} & \text{if } d(x, y) \geq t^{1/w} \\ \left( \frac{d(x, y)}{t^{1/w}} \right)^{\frac{\eta}{p'}} & \text{if not.} \end{cases}$$

On the other hand, one can write

$$\begin{aligned} & \int_0^t s^{k-1} (A^k e^{-sA} f(x) - A^k e^{-sA} f(y)) ds \\ &= \int_0^t s^{k-1} (e^{-(s/2)A} A^k e^{-(s/2)A} f(x) - e^{-(s/2)A} A^k e^{-(s/2)A} f(y)) ds, \end{aligned}$$

and by analyticity, we have

$$\|A^k e^{-(s/2)A} f\|_p \leq C s^{\frac{\beta}{w}-k} \|A^{\beta/w} f\|_p.$$

Then, by using Proposition 5 and arguing similarly as the first case, we also obtain

$$\begin{aligned} |f(x) - f(y)| &\leq \frac{C}{V^{1/p}(x, d)} \\ &\times \max \left\{ d^{\frac{\eta}{p'} + \frac{\nu}{p}} \|f\|_p^{1-\theta} \|A^{\beta/w} f\|_p^\theta, d^{\frac{\eta}{p'}} \|f\|_p^{1-\theta'} \|A^{\beta/w} f\|_p^{\theta'} \right\}, \end{aligned}$$

where  $\theta = \frac{1}{\beta}(\frac{\eta}{p'} + \frac{\nu}{p})$  and  $\theta' = \frac{\eta}{\beta p'}$ .

Set  $\alpha = \frac{\eta}{p'} + \frac{\nu}{p}$  and  $\alpha' = \frac{\eta}{p'}$ , then  $\beta = \frac{\alpha}{\alpha'} = \frac{\theta'}{\alpha}$ ,  $\alpha - \alpha' = \frac{\nu}{p}$ ,  $\alpha > \frac{\nu}{p}$ ,  $\alpha' > 0$  and

$$\begin{aligned} |f(x) - f(y)| &\leq \frac{C}{V^{1/p}(x, d)} \\ &\times \max \left\{ d^\alpha \|f\|_p^{1-\theta} \|A^{\alpha/w} f\|_p^\theta, d^{\alpha'} \|f\|_p^{1-\theta'} \|A^{\alpha'/w} f\|_p^{\theta'} \right\}. \end{aligned}$$

For  $p > \frac{\nu+\eta}{\eta}$ , one has  $\eta > \frac{\nu+\eta}{p}$ , then  $\eta(1 - \frac{1}{p}) > \frac{\nu}{p}$ , hence  $\alpha' = \frac{\eta}{p'} > \frac{\nu}{p}$ .

( $\Leftarrow$ ) Let  $p > 1$ , assume that for some  $\alpha, \alpha' > \frac{\nu}{p}$  and  $\theta, \theta' \in ]0, 1[$ :

$$\begin{aligned} |f(x) - f(y)| &\leq \frac{C}{V^{1/p}(x, d)} \\ &\times \max \left\{ d^\alpha \|f\|_p^{1-\theta} \|A^{\alpha/w} f\|_p^\theta, d^{\alpha'} \|f\|_p^{1-\theta'} \|A^{\alpha'/w} f\|_p^{\theta'} \right\}, \end{aligned}$$

for all  $f \in \mathcal{D}$  and  $x, y \in M$ .

Let  $z \in M$ , by analyticity of  $(e^{-tA})_{t>0}$ ,  $p_t(\cdot, z)$  belong to  $\mathcal{D}_p(A^{\alpha/w})$ . Then by choosing  $f = p_t(\cdot, z)$  in the previous inequality, one has for all  $x, y \in M$ :

$$\begin{aligned} |p_t(x, z) - p_t(y, z)| &\leq \frac{C}{V^{1/p}(x, d)} \\ &\times \max \left\{ d^\alpha \|p_t(\cdot, z)\|_p^{1-\theta} \|A^{\alpha/w\theta} p_t(\cdot, z)\|_p^\theta, \right. \\ &\quad \left. d^{\alpha'} \|p_t(\cdot, z)\|_p^{1-\theta'} \|A^{\alpha'/w\theta'} p_t(\cdot, z)\|_p^{\theta'} \right\}. \end{aligned}$$

By analyticity,

$$\|A^{\alpha/w\theta} p_t(\cdot, z)\|_p = \|A^{\alpha/w\theta} e^{-(t/2)A} p_{t/2}(\cdot, z)\|_p \leq C t^{-\alpha/w\theta} \|p_{t/2}(\cdot, z)\|_p.$$

Since  $(e^{-tA})_{t>0}$  is symmetric Markovian, then  $\|p_{t/2}(\cdot, z)\|_1 \leq 1$ . Hence  $(UE_w)$  yields

$$\|p_t(\cdot, z)\|_p \leq \frac{C}{[V(z, t^{1/w})]^{1-\frac{1}{p}}}, \quad \forall t > 0,$$

and by  $(D_\nu)$ , one has

$$\|p_{t/2}(\cdot, z)\|_p \leq \frac{C}{[V(z, t^{1/w})]^{1-\frac{1}{p}}},$$

then

$$\|p_t(\cdot, z)\|_p^{1-\theta} \|A^{\alpha/w\theta} p_t(\cdot, z)\|_p^\theta \leq C \frac{t^{\alpha/w}}{[V(z, t^{1/w})]^{1-\frac{1}{p}}}.$$

Therefore

$$\begin{aligned} &|p_t(x, z) - p_t(y, z)| \\ &\leq \frac{C}{V^{1/p}(x, d)} \max \left\{ \left( \frac{d}{t^{1/w}} \right)^\alpha \frac{1}{[V(z, t^{1/w})]^{1-\frac{1}{p}}}, \left( \frac{d}{t^{1/w}} \right)^{\alpha'} \frac{1}{[V(z, t^{1/w})]^{1-\frac{1}{p}}} \right\} \\ &\leq \frac{C}{V(z, t^{1/w})} \left( \frac{V(z, t^{1/w})}{V(x, d)} \right)^{1/p} \max \left\{ \left( \frac{d}{t^{1/w}} \right)^\alpha, \left( \frac{d}{t^{1/w}} \right)^{\alpha'} \right\}. \end{aligned}$$

Then, similarly as in the proof of the converse in Theorem 6, we obtain the desired result.  $\square$

A consequence of Theorems 6 and 7 is the following.

**Corollary 8.** *Let  $(M, d, \mu, A)$  as above satisfy the doubling condition  $(D_\nu)$ . Assume that  $(UE_w)$  holds. Then*

$$(S_w) \text{ for any } p > 1 \text{ and } \alpha, \alpha' > \frac{\nu}{p},$$

*implies, for all  $p$  large enough, that there are  $\alpha, \alpha' > \frac{\nu}{p}$  and  $\theta, \theta' \in ]0, 1[$  such that  $\alpha - \alpha' = \frac{\nu}{p}$ ,  $\frac{\alpha}{\theta} = \frac{\alpha'}{\theta'}$  and  $(G_w)$  holds.*

*Conversely,  $(G_w)$  for any  $p > 1$  and  $\alpha, \alpha' > \frac{\nu}{p}$ ,  $\theta, \theta' \in ]0, 1[$ , implies, for all  $p$  large enough, that there are  $\alpha, \alpha' > \frac{\nu}{p}$  such that  $(S_w)$  holds.*

**Remark 9.** We can also deduce  $(G_w)$  from  $(S_w)$  by using the momentum inequality: for a nonnegative operator in a Banach space

$$\|A^{\alpha/w} f\|_p \leq C \|f\|_p^{1-\theta} \|A^{\alpha/w\theta} f\|_p^\theta,$$

(see [15]). The previous corollary, gives us more information about the relation between  $\alpha, \alpha', \theta$  and  $\theta'$ .

### 4.3. Further results.

Let  $(M, d, \mu, A)$  be as before and  $w \geq 2$ . From the doubling property, one can also write, there are  $\nu > 0$  and  $\nu' \geq 0$  such that

$$(D_{\nu, \nu'}) \quad C' \left( \frac{s}{r} \right)^{\nu'} \leq \frac{V(x, s)}{V(x, r)} \leq C \left( \frac{s}{r} \right)^\nu, \quad \forall x \in M, s \geq r > 0.$$

Note that, when  $M$  has an infinite measure, we can take  $\nu' > 0$ , (see [8]).

By using  $(D_{\nu, \nu'})$  instead of  $(D_\nu)$  and arguing as before, almost all the previous results can be written for all  $p > 1$  with  $\nu$  and  $\nu'$ . In other words, one has just take  $\nu' = 0$  (which is always possible) to find the previous proofs.

More precisely, instead of Propositions 4 and 5, we obtain: If  $(LY_w)$  holds, then there exists  $\eta \in ]0, 1[$  such that for all  $p > 1$ :

$$\begin{aligned} & |p_t(x, z) - p_t(y, z)| \\ & \leq C \frac{(E_w(C, x, z, t) + E_w(C, y, z, t))^{1/p}}{V^{1/p}(x, d) V^{1/p'}(z, t^{1/w})} \begin{cases} \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu}{p}} & \text{if } d \geq t^{1/w} \\ \left( \frac{d}{t^{1/w}} \right)^{\frac{\eta}{p'} + \frac{\nu'}{p}} & \text{if not,} \end{cases} \end{aligned}$$

for all  $t > 0$  and  $\mu$ -a.e.  $x, y, z \in M$ .

And

$$(7) \quad |e^{-tA}f(x) - e^{-tA}f(y)| \leq \frac{C\|f\|_p}{V^{1/p}(x,d)} \begin{cases} \left(\frac{d}{t^{1/w}}\right)^{\frac{\eta}{p'} + \frac{\nu}{p}}, & \text{if } d \geq t^{1/w} \\ \left(\frac{d}{t^{1/w}}\right)^{\frac{\eta}{p'} + \frac{\nu'}{p}} & \text{if not,} \end{cases}$$

for all  $t > 0$ ,  $f \in \mathcal{D}$  and  $\mu$ -a.e.  $x, y \in M$ , where  $d$  denote to  $d(x, y)$ .

Consequently, we obtain the following proposition.

**Proposition 10.** *Let  $(M, d, \mu, A)$  as before satisfy the doubling condition  $(D_{\nu, \nu'})$ , assume that the  $(UE_w)$  holds, then for all  $p > 1$ ,*

- (i)  $(LE_w) \Rightarrow (S_w)$  for some  $\alpha > \frac{\nu}{p}$  and  $\alpha' > \frac{\nu'}{p}$ .
- (ii)  $(LE_w) \Rightarrow (G_w)$  for some  $\alpha > \frac{\nu}{p}$ ,  $\alpha' > \frac{\nu'}{p}$   
such that  $\alpha - \alpha' = \frac{\nu - \nu'}{p}$  and  $\frac{\alpha}{\theta} = \frac{\alpha'}{\theta'}$ .

*Proof:* (i) By using (7) instead of Proposition 5 in the proof of Theorem 6, we obtain for all  $p > 1$ ,  $(S_w)$  holds for  $\alpha = \frac{\eta}{p'} + \frac{\nu + \delta}{p}$  and  $\alpha' = \frac{\eta}{p'} + \frac{\nu - \delta}{p}$  where  $\delta \in ]0, \frac{\nu}{p'}\eta[$ , whence  $\alpha > \frac{\nu}{p}$  and  $\alpha' > \frac{\nu'}{p}$ .

(ii) Similarly as in the proof of Theorem 7 and by using (7) instead of Proposition 5, we obtain that for all  $p > 1$ , there are  $\alpha = \frac{\eta}{p'} + \frac{\nu}{p}$ ,  $\alpha' = \frac{\eta}{p'} + \frac{\nu'}{p}$ ,  $\theta, \theta' \in ]0, 1[$  such that  $\frac{\alpha}{\theta} = \frac{\alpha'}{\theta'}$  and  $(G_w)$  holds. It is clear that  $\alpha - \alpha' = \frac{\nu - \nu'}{p}$ .  $\square$

In the case when the volume growth is polynomial (i.e.  $V(x, r) \simeq r^D$ ,  $D > 0$ ), we reobtain the following result from Ouhabaz (see [16]).

**Corollary 11.** *Assume that  $V(x, r) \simeq r^D$ , if  $(UE_w)$  holds, then for all  $p > 1$*

$$(LE_w) \quad \Leftrightarrow \quad \frac{|f(x) - f(y)|}{d(x, y)^{\alpha - \frac{D}{p}}} \leq C\|f\|_p^\theta \|A^{\alpha/w\theta} f\|_p^{1-\theta},$$

for some  $\alpha > \frac{D}{p}$ ,  $\theta \in ]0, 1[$ .

*Proof:* If  $V(x, r) \simeq r^D$ , then one can write  $(D_{\nu, \nu'})$  with  $\nu = \nu' = D$ . Therefore

( $\Leftarrow$ ) it suffices to take  $\alpha = \alpha'$  and use Theorem 7.

( $\Rightarrow$ ) by applying (ii) of the previous proposition,  $(LE_w)$  implies  $(G_w)$  with  $\alpha = \alpha' > \frac{D}{p}$  and  $\theta = \theta'$ .  $\square$

**Acknowledgements.** I would like to thank Professor Thierry Coulhon, who brought the subject to my attention, for interesting and stimulating discussions. My thanks go also to Professor El-Maati Ouhabaz, for interesting conversations and suggestions during our meeting at the colloquium “Analyse Fonctionnelle et Harmonique, 19–22 Novembre 07”, held at the CIRM “Centre International des Rencontres Mathématiques”. I would also like to thank the referees for useful comments.

### References

- [1] M. T. BARLOW, Which values of the volume growth and escape time exponent are possible for a graph?, *Rev. Mat. Iberoamericana* **20(1)** (2004), 1–31.
- [2] M. T. BARLOW AND R. F. BASS, Random walks on graphical Sierpinski carpets, in: “*Random walks and discrete potential theory*” (Cortona, 1997), Sympos. Math. XXXIX, Cambridge Univ. Press, Cambridge, 1999, pp. 26–55.
- [3] M. T. BARLOW AND R. F. BASS, Divergence form operators on fractal-like domains, *J. Funct. Anal.* **175(1)** (2000), 214–247.
- [4] I. BENJAMINI, I. CHAVEL, AND E. A. FELDMAN, Heat kernel lower bounds on Riemannian manifolds using the old ideas of Nash, *Proc. London Math. Soc. (3)* **72(1)** (1996), 215–240.
- [5] T. COULHON, Off-diagonal heat kernel lower bounds without Poincaré, *J. London Math. Soc. (2)* **68(3)** (2003), 795–816.
- [6] T. COULHON, Ultracontractivity and Nash type inequalities, *J. Funct. Anal.* **141(2)** (1996), 510–539.
- [7] T. COULHON AND A. GRIGOR’YAN, Random walks on graphs with regular volume growth, *Geom. Funct. Anal.* **8(4)** (1998), 656–701.
- [8] A. GRIGOR’YAN, The heat equation on noncompact Riemannian manifolds, (Russian), *Mat. Sb.* **182(1)** (1991), 55–87; translation in: *Math. USSR-Sb.* **72(1)** (1992), 47–77.
- [9] A. GRIGOR’YAN, Heat kernel upper bounds on a complete non-compact manifold, *Rev. Mat. Iberoamericana* **10(2)** (1994), 395–452.

- [10] A. GRIGOR'YAN, J. HU, AND K.-S. LAU, Heat kernels on metric measure spaces and an application to semilinear elliptic equations, *Trans. Amer. Math. Soc.* **355**(5) (2003), 2065–2095 (electronic).
- [11] A. GRIGOR'YAN AND A. TELCS, Sub-Gaussian estimates of heat kernels on infinite graphs, *Duke Math. J.* **109**(3) (2001), 451–510.
- [12] B. HAMBLBY AND T. KUMAGAI, Heat kernel estimates for symmetric random walks on a class of fractal graphs and stability under rough isometries, in: “*Fractal geometry and applications: a jubilee of Benoît Mandelbrot, Part 2*”, Proc. Sympos. Pure Math. **72**, Amer. Math. Soc., Providence, RI, 2004, pp. 233–259.
- [13] W. HEBISCH AND L. SALOFF-COSTE, On the relation between elliptic and parabolic Harnack inequalities, *Ann. Inst. Fourier (Grenoble)* **51**(5) (2001), 1437–1481.
- [14] P. LI AND S.-T. YAU, On the parabolic kernel of the Schrödinger operator, *Acta Math.* **156**(3–4) (1986), 153–201.
- [15] S. MARTINES, M. SANS, AND M. D. MARTINES, Some inequalities for fractional integrals and derivatives, (Russian), *Dokl. Akad. Nauk SSSR* **315**(5) (1990), 1049–1051; translation in: *Soviet Math. Dokl.* **42**(3) (1991), 876–879.
- [16] E.-M. OUHABAZ, Heat kernels of multiplicative perturbations: Hölder estimates and Gaussian lower bounds, *Indiana Univ. Math. J.* **47**(4) (1998), 1481–1495.
- [17] E.-M. OUHABAZ, “*Analysis of heat equations on domains*”, London Mathematical Society Monographs Series **31**, Princeton University Press, Princeton, NJ, 2005.
- [18] L. SALOFF-COSTE, A note on Poincaré, Sobolev, and Harnack inequalities, *Internat. Math. Res. Notices* **2** (1992), 27–38.
- [19] L. SALOFF-COSTE, Parabolic Harnack inequality for divergence-form second-order differential operators, Potential theory and degenerate partial differential operators (Parma), *Potential Anal.* **4**(4) (1995), 429–467.
- [20] L. SALOFF-COSTE, “*Aspects of Sobolev-type inequalities*”, London Mathematical Society Lecture Note Series **289**, Cambridge University Press, Cambridge, 2002.
- [21] E. M. STEIN, “*Topics in harmonic analysis related to the Littlewood-Paley theory*”, Annals of Mathematics Studies **63**, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1970.

Département de Mathématiques  
Université de Cergy-Pontoise  
Site de Saint Martin, 2 avenue Adolphe-Chauvin  
95302 Cergy-Pontoise cedex  
France  
*E-mail address:* `salahaddine.boutayeb@u-cergy.fr`

Primera versió rebuda el 29 d'abril de 2008,  
darrera versió rebuda el 18 de setembre de 2008.