

INEQUALITIES FOR POISSON INTEGRALS WITH SLOWLY GROWING DIMENSIONAL CONSTANTS

LOUKAS GRAFAKOS*, ENRICO LAENG**, AND CARLO MORPURGO***

Abstract

Let P_t be the Poisson kernel. We study the following L^p inequality for the Poisson integral $Pf(x, t) = (P_t * f)(x)$ with respect to a Carleson measure μ :

$$\|Pf\|_{L^p(\mathbb{R}_+^{n+1}, d\mu)} \leq c_{p,n} \kappa(\mu)^{\frac{1}{p}} \|f\|_{L^p(\mathbb{R}^n, dx)},$$

where $1 < p < \infty$ and $\kappa(\mu)$ is the Carleson norm of μ . It was shown by Verbitsky [V] that for $p > 2$ the constant $c_{p,n}$ can be taken to be independent of the dimension n . We show that $c_{2,n} = O((\log n)^{\frac{1}{2}})$ and that $c_{p,n} = O(n^{\frac{1}{p} - \frac{1}{2}})$ for $1 < p < 2$ as $n \rightarrow \infty$. We observe that standard proofs of this inequality rely on doubling properties of cubes and lead to a value of $c_{p,n}$ that grows exponentially with n .

1. Introduction

The object of study in this article is the following Carleson measure inequality [C1], [C2], valid for $1 < p < \infty$

$$(1.1) \quad \|Pf\|_{L^p(\mathbb{R}_+^{n+1}, d\mu)} \leq c_{p,n} \kappa(\mu)^{\frac{1}{p}} \|f\|_{L^p(\mathbb{R}^n)},$$

where μ is a Carleson measure on \mathbb{R}_+^{n+1} with norm $\kappa(\mu)$ and P is the Poisson integral of a function f on \mathbb{R}^n .

We begin by recalling these notions and establishing notation. We denote by $\bar{B}(x_0, r)$ the closed ball in \mathbb{R}^n with radius r centered at x_0 . The Carleson tent $T[\bar{B}(x_0, r)]$ over the ball $\bar{B}(x_0, r)$ is defined as the

2000 *Mathematics Subject Classification*. Primary: 42B25; Secondary: 31B05.

Key words. Carleson measures, harmonic functions, dimension free estimates.

*Supported in part by the NSF Grant DMS-0400387.

**Supported in part by the Italian Fondi Ministeriali MIUR.

***Supported by the NSF Grant DMS-0200574.

set of all points $(x, t) \in \mathbb{R}^n \times \mathbb{R}^+$ such that $|x - x_0|^2 + t^2 < r^2$. Denote by $|K|$ the Lebesgue measure of a set K in \mathbb{R}^n . A Borel measure μ on \mathbb{R}_+^{n+1} is called a Carleson measure if its Carleson norm $\kappa(\mu) = \sup \left\{ \frac{\mu(T[\overline{B}(x_0, r)])}{|\overline{B}(x_0, r)|} : x_0 \in \mathbb{R}^n, r > 0 \right\}$ is a finite number.

We define the Poisson integral Pf , i.e., the harmonic extension of f to \mathbb{R}_+^{n+1} , as the convolution

$$(1.2) \quad Pf(x, a) = P_a * f(x) = \int_{\mathbb{R}^n} P_a(x - y)f(y) dy,$$

where the Poisson kernel is defined, for all $a > 0$, $x \in \mathbb{R}^n$ by

$$(1.3) \quad P_a(x) = \frac{\gamma_n a}{(a^2 + |x|^2)^{\frac{n+1}{2}}}, \quad \text{with } \gamma_n = \frac{\Gamma\left(\frac{n+1}{2}\right)}{\pi^{\frac{n+1}{2}}}.$$

The Lebesgue measure of the closed unit ball $\overline{B}(0, 1)$ and the surface measure of its boundary are denoted by

$$(1.4) \quad \Omega_n = |\overline{B}(0, 1)| = \frac{\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2} + 1\right)}, \quad \omega_{n-1} = |\partial\overline{B}(0, 1)| = \frac{2\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2}\right)} = n\Omega_n.$$

Verbitsky [V] gave an elegant proof of (1.1) that yields a constant $c_{p,n}$ independent of n whenever $p > 2$. The starting point of Verbitsky's argument is the use of interpolation and duality to derive (1.1) from the equivalent inequalities

$$(1.5) \quad \|Pf\|_{L^{2,\infty}(\mathbb{R}_+^{n+1}, d\mu)} \leq c' \kappa(\mu)^{\frac{1}{2}} \|f\|_{L^2(\mathbb{R}^n)},$$

$$(1.6) \quad \|P^*(g, \mu)\|_{L^2(\mathbb{R}^n)} \leq c' \kappa(\mu)^{\frac{1}{2}} \|g\|_{L^{2,1}(\mathbb{R}_+^{n+1}, d\mu)},$$

for some absolute constant c' independent of n , where P^* is the Balayage operator defined for functions g on \mathbb{R}_+^{n+1} by

$$P^*(g, \mu)(x) = \int_{\mathbb{R}_+^{n+1}} P_b(x - y)g(y, b) d\mu(y, b).$$

Using the semigroup property for the Poisson kernel,

$$(1.7) \quad \int_{\mathbb{R}^n} P_a(\tau - x)P_b(\tau - y) d\tau = P_{a+b}(x - y),$$

Verbitsky obtained (1.6) with c' independent of n for characteristic functions of subsets of \mathbb{R}_+^{n+1} and this is enough to establish (1.6) for general μ -measurable functions g (see [SW]).

The following theorem is the main result of this article, which grew out of our attempts to extend Verbitsky's theorem to the case $1 < p \leq 2$:

Theorem 1.1. *For $1 < p < 2$ the following Carleson measure inequality holds:*

$$(1.8) \quad \|Pf\|_{L^p(\mathbb{R}_+^{n+1}, d\mu)} \leq C_p n^{\frac{1}{p}-\frac{1}{2}} \kappa(\mu)^{\frac{1}{p}} \|f\|_{L^p(\mathbb{R}^n)},$$

with C_p independent of n . For $p = 2$ the following holds:

$$(1.9) \quad \|Pf\|_{L^2(\mathbb{R}_+^{n+1}, d\mu)} \leq C(\log n)^{\frac{1}{2}} \kappa(\mu)^{\frac{1}{2}} \|f\|_{L^2(\mathbb{R}^n)},$$

with C independent of n .

We observe that the estimate above is sharper than those obtained from the other known proofs of the Carleson measure inequality (1.1) (see, e.g., [A], [C2], [G], [H], [N], [S]); these proofs yield constants that grow exponentially in n .

The authors would like to thank Igor Verbitsky for pointing out this problem to them and for sharing some of his ideas with them.

2. An integral formula involving Poisson kernels

In order to extend the arguments in [V] we are lead to consider the equivalent inequalities for $k = 3, 4, 5, \dots$

$$\|Pf\|_{L^{\frac{k}{k-1}, \infty}(\mathbb{R}_+^{n+1}, d\mu)} \leq \bar{c}(k, n) \kappa(\mu)^{\frac{k-1}{k}} \|f\|_{L^{\frac{k}{k-1}}(\mathbb{R}^n)},$$

$$\|P^*(g, \mu)\|_{L^k(\mathbb{R}^n)} \leq \bar{c}(k, n) \kappa(\mu)^{\frac{k-1}{k}} \|g\|_{L^{k,1}(\mathbb{R}_+^{n+1}, d\mu)},$$

$$\|P^*(\chi_E, \mu)\|_{L^k(\mathbb{R}^n)} \leq c(k, n) \kappa(\mu)^{\frac{k-1}{k}} \mu(E)^{\frac{1}{k}}, \quad \forall E \subset \mathbb{R}_+^{n+1}.$$

For $k = 2$ these inequalities hold with $c(2, n) \leq c_{2,n} \leq c_2$ independent of n , due to [V], (see (1.5) above). Our goal is to obtain good estimates for the constants $c(k, n)$ for all $k = 3, 4, \dots$ and then use the Marcinkiewicz interpolation theorem to deduce (1.1).

Since

$$\|P^*(\chi_E, \mu)\|_{L^k(\mathbb{R}^n)}^k = \int_{E^k} \int_{\mathbb{R}^n} \prod_{j=1}^k P_{a_j}(x_j - \tau) d\tau d\mu(x_1, a_1) \dots d\mu(x_k, a_k)$$

our first task is to derive a workable formula for

$$\int_{\mathbb{R}^n} \prod_{j=1}^k P_{a_j}(x_j - \tau) d\tau,$$

which, in the case $k = 2$, is computed explicitly via the semigroup property. In this section we prove the following:

Proposition 2.1. *Let $P_a(x)$ be the Poisson kernel, as in definition (1.3). Then, for any $a_1, \dots, a_k \in (0, \infty)$ and for any $x_1, \dots, x_k \in \mathbb{R}^n$, we have*

$$(2.1) \quad \int_{\mathbb{R}^n} \prod_{j=1}^k P_{a_j}(\tau - x_j) d\tau \\ = \frac{a_1 a_2 \dots a_k}{\pi^{(k-1)\frac{n}{2} + \frac{k}{2}}} \int_0^\infty \dots \int_0^\infty \frac{(u_1 u_2 \dots u_k)^{\frac{n-1}{2}}}{(u_1 + u_2 + \dots + u_k)^{\frac{n}{2}}} \\ \times \text{Exp} \left[- \sum_{j=1}^k a_j^2 u_j - \frac{\sum_{1 \leq i < j \leq k} u_i u_j |x_i - x_j|^2}{u_1 + u_2 + \dots + u_k} \right] du_1 \dots du_k.$$

In particular, when $k = 3$, the formula above can be written as

$$(2.2) \quad \int_{\mathbb{R}^n} P_a(\tau - x) P_b(\tau - y) P_c(\tau - z) d\tau \\ = \frac{abc}{\pi^{n+\frac{3}{2}}} \int_0^\infty \int_0^\infty \int_0^\infty \frac{(uvw)^{\frac{n-1}{2}}}{(u+v+w)^{\frac{n}{2}}} \\ \times \text{Exp} \left[- \left\{ a^2 u + b^2 v + c^2 w \right. \right. \\ \left. \left. + \frac{uv|x-y|^2 + uw|x-z|^2 + vw|y-z|^2}{u+v+w} \right\} \right] du dv dw,$$

and when $k = 2$, we have

$$(2.3) \quad \int_{\mathbb{R}^n} P_a(\tau - x) P_b(\tau - y) d\tau \\ = \frac{ab}{\pi^{\frac{n}{2}+1}} \int_0^{+\infty} \int_0^{+\infty} \frac{(uv)^{\frac{n-1}{2}}}{(u+v)^{\frac{n}{2}}} e^{-\{a^2 u + b^2 v + \frac{uv}{u+v}|x-y|^2\}} du dv.$$

Proof: We start from the ‘‘subordination’’ formula

$$P_a(x) = \int_0^{+\infty} \beta_a(u) e^{-u|x|^2} du$$

where

$$\beta_a(u) = \frac{a}{\pi^{\frac{n+1}{2}}} e^{-a^2 u} u^{\frac{n-1}{2}}$$

which can be easily deduced using the definition of the Gamma function.

The integral on the left hand side in (2.1) can now be rewritten as

$$\begin{aligned}
(2.4) \quad & \int_{\mathbb{R}^n} \prod_{j=1}^k P_{a_j}(\tau - x_j) d\tau \\
&= \int_{\mathbb{R}^n} \prod_{j=1}^k \left(\int_0^{+\infty} \beta_{a_j}(u_j) e^{-u_j |\tau - x_j|^2} du_j \right) d\tau \\
&= \frac{a_1 \cdots a_k}{\pi^{\frac{(n+1)k}{2}}} \int_{\mathbb{R}^n} \int_{[0, \infty)^k} (u_1 \cdots u_k)^{\frac{n-1}{2}} e^{-\sum_{j=1}^k (a_j^2 u_j + u_j |\tau - x_j|^2)} du_1 \cdots du_k d\tau \\
&= \frac{a_1 \cdots a_k}{\pi^{\frac{(n+1)k}{2}}} \int_{[0, \infty)^k} (u_1 \cdots u_k)^{\frac{n-1}{2}} e^{-\sum_{j=1}^k a_j^2 u_j} \\
&\quad \times \left[\int_{\mathbb{R}^n} e^{-\sum_{j=1}^k u_j |\tau - x_j|^2} d\tau \right] du_1 \cdots du_k.
\end{aligned}$$

Let us now rewrite the inner integral inside square brackets using Cartesian coordinates, namely $\tau = (\tau_1, \dots, \tau_n)$ and $x_j = (x_{j1}, \dots, x_{jn})$. We obtain

$$\begin{aligned}
& \int_{\mathbb{R}^n} e^{-\sum_{j=1}^k u_j |\tau - x_j|^2} d\tau \int_{\mathbb{R}^n} e^{-\{u_1(\tau_1 - x_{11})^2 + u_2(\tau_1 - x_{21})^2 + \cdots + u_k(\tau_1 - x_{k1})^2\}} \dots \\
& \dots e^{-\{u_1(\tau_n - x_{1n})^2 + u_2(\tau_n - x_{2n})^2 + \cdots + u_k(\tau_n - x_{kn})^2\}} d\tau_1 d\tau_2 \dots d\tau_n = \prod_{r=1}^n I_r
\end{aligned}$$

where, for each index $r = 1, 2, \dots, n$, we have defined

$$(2.5) \quad I_r = \int_{-\infty}^{+\infty} e^{-(A\tau_r^2 - 2B\tau_r + C)} d\tau_r$$

with

$$\begin{aligned}
A &= u_1 + u_2 + \cdots + u_k \\
B &= B_r = u_1 x_{1r} + u_2 x_{2r} + \cdots + u_k x_{kr} \\
C &= C_r = u_1 x_{1r}^2 + u_2 x_{2r}^2 + \cdots + u_k x_{kr}^2.
\end{aligned}$$

We claim that, for each choice of the coordinate index $r = 1, \dots, n$ we have

$$B^2 - AC = B_r^2 - AC_r = - \sum_{i < j} u_i u_j (x_{ir} - x_{jr})^2$$

where the sum is taken over all possible pairs of distinct indices i and j both running from 1 to k (there are $k(k-1)/2$ of such pairs). In fact this claim is easily checked by observing that in $B^2 - AC$ all the square terms of the kind $u_i^2 x_{ir}^2$ cancel out, while the remaining mixed terms can be collected in groups of three, each group giving $-u_i u_j (x_{ir}^2 - 2x_{ir} x_{jr} + x_{jr}^2)$.

Now, completing the square in (2.5) we obtain

$$I_r = e^{\frac{B^2 - AC}{A}} \int_{-\infty}^{+\infty} e^{-A(\tau_r - \frac{B}{A})^2} d\tau_r = \sqrt{\frac{\pi}{u_1 + \dots + u_k}} e^{-\frac{\sum_{i < j} u_i u_j (x_{ir} - x_{jr})^2}{u_1 + \dots + u_k}}$$

which implies that

$$\int_{\mathbb{R}^n} e^{-\sum_{j=1}^k u_j |\tau - x_j|^2} d\tau = \left(\frac{\pi}{u_1 + \dots + u_k} \right)^{\frac{n}{2}} e^{-\frac{\sum_{i < j} u_i u_j |x_i - x_j|^2}{u_1 + \dots + u_k}}.$$

Using this identity in the square brackets in (2.4) and simplifying, we obtain (2.1). \square

Remark. The reader may wonder if it is possible to obtain a formula for the left hand side in (2.1) that does not involve any integrals, something analogous to (1.7) for $k \geq 3$. When $n = 1$ using residues (plus some involved algebraic manipulations) we were able to show that

$$\begin{aligned} & \int_{\mathbb{R}} P_a(\tau - x) P_b(\tau - y) P_c(\tau - z) d\tau \\ &= \frac{ab}{(a+c)(b+c)} P_{a+c}(x-z) P_{b+c}(y-z) \\ &+ \frac{ac}{(a+b)(c+b)} P_{b+c}(y-z) P_{a+b}(x-y) \\ &+ \frac{bc}{(b+a)(c+a)} P_{a+c}(x-z) P_{a+b}(x-y) \\ &+ 4\pi \frac{abc(a+b+c)}{(a+b)(a+c)(b+c)} P_{a+b}(x-y) P_{a+c}(x-z) P_{b+c}(y-z). \end{aligned}$$

Unfortunately, for $n > 1$, an integral-free formula of this sort is difficult to obtain. It seems that the size and complexity of the formula grows quickly with n , and furthermore, there is no obvious “leading term” to be used in our estimates as $n \rightarrow \infty$. On the other hand we will show that (2.1), after some manipulations, suffices for the purposes of the proof of Theorem 1.1.

3. Proof of Theorem 1.1: the case $\frac{3}{2} < p < 2$

For clarity of exposition we first give a detailed proof of Theorem 1.1 in the case $3/2 < p < 2$. In the next section we indicate how the same technique can be adapted to the case $p \in \left(\frac{k}{k-1}, 2\right)$, any $k = 4, 5, \dots$.

Let us start by showing that the Balayage operator P^* satisfies the following estimate for all Carleson measures μ , for all μ -measurable subsets E of \mathbb{R}_+^{n+1} , and for all $n = 1, 2, \dots$,

$$(3.1) \quad \|P^*(\chi_E, \mu)\|_{L^3(\mathbb{R}^n)}^3 \leq c'' n^{\frac{1}{2}} \kappa(\mu)^2 \mu(E),$$

for some absolute constant c'' . Once (3.1) is established, using duality and [SW] we obtain for some other absolute constant c'''

$$(3.2) \quad \|Pf\|_{L^{\frac{3}{2}, \infty}(\mathbb{R}_+^{n+1}, d\mu)} \leq c''' n^{\frac{1}{6}} \kappa(\mu)^{\frac{2}{3}} \|f\|_{L^{\frac{3}{2}}(\mathbb{R}^n)}$$

and thus, by the Marcinkiewicz interpolation and (1.5), Theorem 1.1 follows in the case $3/2 < p < 2$.

We have

$$(3.3) \quad \|P^*(\chi_E, \mu)\|_{L^3(\mathbb{R}^n)}^3 \\ = \int_{E^3} \int_{\mathbb{R}^n} P_a(\tau - x) P_b(\tau - y) P_c(\tau - z) d\tau d\mu(x, a) d\mu(y, b) d\mu(z, c).$$

We will use the following modification of formula (2.2):

Lemma 3.1. *The integral in (2.2) can also be written as*

$$\int_{\mathbb{R}^n} P_a(\tau - x) P_b(\tau - y) P_c(\tau - z) d\tau \\ = \frac{\Gamma\left(\frac{n+3}{2}\right)}{\pi^{n+\frac{3}{2}}} abc \int_0^1 \int_0^1 \frac{(1-t)^n t^{\frac{n-1}{2}} (s(1-s))^{\frac{n-1}{2}}}{\{a^2 t + (1-t)B^2 + t(1-t)|x-q|^2\}^{n+\frac{3}{2}}} dt ds$$

where

$$(3.4) \quad B^2 = B^2(s, y, z) = b^2 s + c^2(1-s) + s(1-s)|y-z|^2$$

$$(3.5) \quad q = q(s, y, z) = sy + (1-s)z.$$

We will also need the following estimates:

Lemma 3.2. *The following inequalities hold for $\alpha > 0$, $\beta > 0$, $\gamma > 0$ such that $\beta > \alpha$ and $\gamma < \beta + 2$ and for any $D \in \mathbb{R}$.*

$$\begin{aligned} & \int_0^1 t^\alpha (1-t)^\beta \{a^2 t + (1-t)B^2 + t(1-t)D^2\}^{-\gamma} dt \\ & \leq \frac{1}{aB} \int_0^1 t^\alpha (1-t)^\beta \{a^2 t + (1-t)B^2 + t(1-t)D^2\}^{-\gamma+1} dt \\ & \leq \frac{1}{aB} \int_0^1 t^\alpha (1-t)^{\beta-\gamma+1} \{B^2 + t(a^2 + D^2)\}^{-\gamma+1} dt. \end{aligned}$$

The proofs of these lemmas are postponed until the end of this section.

Returning to the proof of (3.1), we split (3.3) in the six regions $a \leq b \leq c$, $a \leq c \leq b$, $b \leq c \leq a$, $b \leq a \leq c$, $c \leq b \leq a$, and $c \leq a \leq b$; by symmetry we need only consider the region over which $a \leq b \leq c$. Following [V] we write

$$(3.6) \quad \begin{aligned} & \|P^*(\chi_E, \mu)\|_{L^3(\mathbb{R}^n)}^3 (6\mu(E))^{-1} \\ & \leq \sup_{(z,c) \in E} \int_{\mathbb{R}^n \times (0,c]} \left[\int_{\mathbb{R}^n \times (0,b]} \int_{\mathbb{R}^n} P_a(\tau - x) \right. \\ & \quad \left. \times P_b(\tau - y) P_c(\tau - z) d\tau d\mu(x, a) \right] d\mu(y, b). \end{aligned}$$

Observe that $a \leq b \leq c$ implies $a \leq B$. Applying Lemma 3.2 for each fixed s , (y, b) , and (z, c) , with B and q as in (3.4) and (3.5), $D = |x - q|$,

$\alpha = (n-1)/2$, $\beta = n$ and $\gamma = n + 3/2$, we have

$$\begin{aligned}
(3.7) \quad & \int_{\mathbb{R}^n \times (0,b]} \frac{\Gamma(n+\frac{3}{2})}{\pi^{n+\frac{3}{2}}} abc \int_0^1 \frac{t^{\frac{n-1}{2}}(1-t)^n}{(a^2t+(1-t)B^2+t(1-t)|x-q|^2)^{n+\frac{3}{2}}} dt d\mu(x, a) \\
& \leq \int_{\mathbb{R}^n \times (0,b]} \frac{bc\Gamma(n+\frac{3}{2})}{B\pi^{n+\frac{3}{2}}} \int_0^1 \frac{t^{\frac{n-1}{2}}(1-t)^{-1/2}}{(B^2+t(a^2+|x-q|^2))^{n+\frac{1}{2}}} dt d\mu(x, a) \\
& \leq \frac{bc(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{B\pi^{n+\frac{3}{2}}} \int_0^1 \int_{\mathbb{R}_+^{n+1}} \int_{B^2+t(a^2+|x-q|^2)}^\infty \frac{t^{\frac{n-1}{2}}(1-t)^{-\frac{1}{2}}}{r^{n+\frac{3}{2}}} dt dr d\mu(x, a) \\
& \leq \frac{bc(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{B\pi^{n+\frac{3}{2}}} \int_0^1 \int_{B^2}^\infty \mu \left(T \left[\overline{B} \left(q, \left(\frac{r-B^2}{t} \right)^{\frac{1}{2}} \right) \right] \right) \\
& \quad \times \frac{t^{\frac{n-1}{2}}(1-t)^{-\frac{1}{2}}}{r^{n+\frac{3}{2}}} dt dr d\mu(x, a) \\
& \leq \Omega_n \kappa(\mu) \frac{bc(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{B\pi^{n+\frac{3}{2}}} \int_0^1 \int_{B^2}^\infty (r-B^2)^{\frac{n}{2}} \\
& \quad \times \frac{t^{-\frac{1}{2}}(1-t)^{-\frac{1}{2}}}{r^{n+\frac{3}{2}}} dt dr d\mu(x, a) \\
& = \kappa(\mu) \frac{bc(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{B^{n+2}\pi^{\frac{n+3}{2}}\Gamma(\frac{n}{2}+1)} \int_0^1 \int_1^\infty \frac{(r-1)^{\frac{n}{2}}}{r^{n+\frac{3}{2}}} t^{-\frac{1}{2}}(1-t)^{-\frac{1}{2}} dt dr d\mu(x, a) \\
& = \kappa(\mu) \frac{bc(n+\frac{1}{2})\Gamma(\frac{n+1}{2})}{B^{n+2}\pi^{\frac{n+1}{2}}} \\
& = \kappa(\mu) \frac{bc(n+\frac{1}{2})\Gamma(\frac{n+1}{2})}{\pi^{\frac{n+1}{2}}(b^2s+c^2(1-s)+s(1-s)|y-z|^2)^{\frac{n}{2}+1}}.
\end{aligned}$$

Next, observe that the following identity holds:

$$\begin{aligned}
(3.8) \quad & \int_0^1 \frac{bc(s(1-s))^{\frac{n-1}{2}}}{(b^2s+c^2(1-s)+s(1-s)|y-z|^2)^{\frac{n}{2}+1}} ds \\
& = \frac{\gamma_n \pi^{\frac{n}{2}+1}}{\Gamma(\frac{n}{2}+1)} \frac{(b+c)}{[(b+c)^2+|y-z|^2]^{\frac{n+1}{2}}}.
\end{aligned}$$

In fact, by the semigroup formula for the Poisson kernel and (2.3) we have

$$P_{b+c}(y-z) = \frac{bc}{\pi^{\frac{n}{2}+1}} \int_0^{+\infty} \int_0^{+\infty} \frac{(uv)^{\frac{n-1}{2}}}{(u+v)^{\frac{n}{2}}} e^{-(b^2u+c^2v+\frac{uv}{u+v}|y-z|^2)} du dv.$$

On the other hand, after the change of variables $v = \lambda u$ the above quantity becomes

$$\begin{aligned} \frac{bc}{\pi^{\frac{n}{2}+1}} \int_0^{+\infty} \frac{\lambda^{\frac{n-1}{2}}}{(1+\lambda)^{\frac{n}{2}}} d\lambda \int_0^{+\infty} u^{\frac{n}{2}} e^{-Wu} du \\ = \frac{bc\Gamma\left(\frac{n}{2}+1\right)}{\pi^{\frac{n}{2}+1}} \int_0^{+\infty} \frac{\lambda^{\frac{n-1}{2}}}{(1+\lambda)^{\frac{n}{2}} W^{\frac{n}{2}+1}} d\lambda \end{aligned}$$

where $W = b^2 + c^2\lambda + \frac{\lambda}{1+\lambda}|y-z|^2$, and the further change of variables $s = 1/(\lambda+1)$ (and therefore $\lambda = (1-s)/s$) gives

$$P_{b+c}(y-z) = \frac{\Gamma\left(\frac{n}{2}+1\right)}{\pi^{\frac{n}{2}+1}} \int_0^1 \frac{bc(s(1-s))^{\frac{n-1}{2}}}{(b^2s+c^2(1-s)+s(1-s)|y-z|^2)^{\frac{n}{2}+1}} ds.$$

We now apply Lemma 3.1 in the inner most integral in (3.6) and then use estimate (3.7) integrated with respect to $(s(1-s))^{\frac{n-1}{2}} ds d\mu(y, b)$ on $[0, 1] \times \mathbb{R}^n \times (0, c]$ and identity (3.8) to obtain

(3.9)

$$\begin{aligned} \int_{\mathbb{R}^n \times (0, c]} \left[\int_{\mathbb{R}^n \times (0, b]} \int_{\mathbb{R}^n} P_a(\tau-x) P_b(\tau-y) P_c(\tau-z) d\tau d\mu(x, a) \right] d\mu(y, b) \\ \leq \kappa(\mu) \frac{(n+\frac{1}{2})\Gamma\left(\frac{n+1}{2}\right)}{\pi^{\frac{n+1}{2}}} \int_{\mathbb{R}_+^{n+1}} \frac{\gamma_n \pi^{\frac{n}{2}+1} (b+c)}{\Gamma\left(\frac{n}{2}+1\right) [(b+c)^2 + |y-z|^2]^{\frac{n+1}{2}}} d\mu(y, b) \\ \leq \kappa(\mu) \frac{2\pi^{1/2} (n+\frac{1}{2})\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n}{2}+1\right)} \int_{\mathbb{R}_+^{n+1}} \gamma_n \frac{c}{[(b+c)^2 + |y-z|^2]^{\frac{n+1}{2}}} d\mu(y, b). \end{aligned}$$

By (13) in [V]

$$\sup_{(z, c) \in \mathbb{R}_+^{n+1}} \int_{\mathbb{R}_+^{n+1}} \gamma_n \frac{c}{[(b+c)^2 + |y-z|^2]^{\frac{n+1}{2}}} d\mu(y, b) \leq C\kappa(\mu)$$

for some absolute constant C independent of n . Therefore (3.9) is bounded above by some absolute constant independent of n times

$$\kappa(\mu)^2 \frac{(n+\frac{1}{2})\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n}{2}+1\right)} \sim \kappa(\mu)^2 \sqrt{n}$$

as $n \rightarrow \infty$, by the well known expansion $\Gamma(z)/\Gamma(z+\alpha) \sim z^{-\alpha}$ for $z \rightarrow \infty$. This last bound for (3.9), together with (3.6), proves (3.1) and hence Theorem 1.1 in the case $3/2 < p < 2$.

Proof of Lemma 3.1: We start with the following n -dimensional identity

$$(3.10) \quad \frac{|y-x|^2}{\alpha} + \frac{|z-x|^2}{\beta} = \frac{|y-z|^2}{\alpha+\beta} + \frac{\alpha+\beta}{\alpha\beta} \left| x - \frac{\beta}{\alpha+\beta}y - \frac{\alpha}{\alpha+\beta}z \right|^2$$

which is valid for all x, y and z in \mathbb{R}^n and for all $\alpha, \beta > 0$ and can be verified by a straightforward calculation.

We now apply (3.10), with $\alpha = \frac{u+v+w}{uv}$ and $\beta = \frac{u+v+w}{uw}$, to the first two terms of

$$(3.11) \quad \frac{uv|x-y|^2 + uw|x-z|^2 + vw|y-z|^2}{u+v+w}$$

which appear in the exponential factor in (2.2). Adding up the third term of (3.11), simplifying, and plugging back into (2.2) we obtain that

$$\begin{aligned} & \int_{\mathbb{R}^n} P_a(\tau-x)P_b(\tau-y)P_c(\tau-z) d\tau \\ &= \frac{abc}{\pi^{n+\frac{3}{2}}} \int_0^{+\infty} \int_0^{+\infty} \int_0^{+\infty} \frac{(uvw)^{\frac{n-1}{2}}}{(u+v+w)^{\frac{n}{2}}} \\ & \quad \times \text{Exp} \left[- \left\{ a^2u + b^2v + c^2w + \frac{vw}{v+w}|y-z|^2 \right. \right. \\ & \quad \left. \left. + \frac{u(v+w)}{u+v+w} \left| x - \frac{v}{v+w}y + \frac{w}{v+w}z \right|^2 \right\} \right] du dv dw. \end{aligned}$$

The change of variables $v = \lambda u$ and $w = \mu u$ transforms the above integral into

$$(3.12) \quad \frac{abc}{\pi^{n+\frac{3}{2}}} \int_0^{+\infty} \int_0^{+\infty} \frac{(\lambda\mu)^{\frac{n-1}{2}}}{(1+\lambda+\mu)^{\frac{n}{2}}} d\lambda d\mu \int_0^{+\infty} u^{n+\frac{1}{2}} e^{-Wu} du,$$

where

$$W = a^2 + b^2\lambda + c^2\mu + \frac{\lambda\mu}{\lambda+\mu}|y-z|^2 + \frac{\lambda+\mu}{1+\lambda+\mu} \left| x - \frac{\lambda}{\lambda+\mu}y + \frac{\mu}{\lambda+\mu}z \right|^2.$$

Expressing the inner integral in (3.12) as a Gamma function we obtain

$$\frac{abc\Gamma\left(n+\frac{3}{2}\right)}{\pi^{n+\frac{3}{2}}} \int_0^{+\infty} \int_0^{+\infty} \frac{(\lambda\mu)^{\frac{n-1}{2}}}{(1+\lambda+\mu)^{\frac{n}{2}} W^{n+\frac{3}{2}}} d\lambda d\mu.$$

Now let us set $t = \frac{1}{1 + \lambda + \mu}$ and $s = \frac{\lambda}{\lambda + \mu}$. This is an invertible transformation of the first quadrant of the (λ, μ) plane onto the square $(0, 1) \times (0, 1)$ of the (t, s) plane. We have $\lambda = \frac{s(1-t)}{t}$ and $\mu = \frac{(1-t)(1-s)}{t}$, while the Jacobian of the transformation is $\frac{1-t}{t^3}$. Collecting and simplifying the various powers of t , $(1-t)$, s and $(1-s)$ that appear we deduce that $\int_{\mathbb{R}^n} P_a(\tau-x)P_b(\tau-y)P_c(\tau-z) d\tau$ is in fact equal to the expression in the statement of Lemma 3.1. \square

Proof of Lemma 3.2: To prove the first inequality, we scale the parameters involved so that $B = 1$ and thus $0 < a \leq 1$. We need to show that

$$\begin{aligned} 0 &\leq \int_0^1 \frac{t^\alpha(1-t)^\beta}{\{a^2t+(1-t)+t(1-t)D^2\}^{\gamma-1}} \left(\frac{1}{a} - \frac{1}{a^2t+(1-t)+t(1-t)D^2} \right) dt \\ &= \frac{1}{a} \int_0^1 \frac{t^\alpha(1-t)^\beta}{\{a^2t+(1-t)+t(1-t)D^2\}^\gamma} (a^2t+1-t-a+t(1-t)D^2) dt. \end{aligned}$$

But $a^2t+1-t-a+t(1-t)D^2 \geq a^2t+1-t-a = (1-a)(1-t(1+a))$ and therefore we need to show that

$$\int_0^1 \frac{t^\alpha(1-t)^\beta}{\{a^2t+(1-t)+t(1-t)D^2\}^\gamma} (1-(1+a)t) dt \geq 0.$$

As $a^2t+(1-t) \leq 1$ and $1-(1+a)t \geq 1-2t$, it will be sufficient to prove that

$$\int_0^1 \frac{t^\alpha(1-t)^\beta}{\{1+t(1-t)D^2\}^\gamma} (1-2t) dt \geq 0.$$

We split this integral in the parts from 0 to $\frac{1}{2}$ and from $\frac{1}{2}$ to 1. Switching variables $t \rightarrow 1-t$ in the second of these integrals we obtain

$$\int_0^{\frac{1}{2}} \frac{(t(1-t))^\alpha}{\{1+t(1-t)D^2\}^\gamma} ((1-t)^{\beta-\alpha} - t^{\beta-\alpha}) (1-2t) dt \geq 0$$

and the first inequality in the lemma is proved. Note that here we have used the hypothesis $\beta > \alpha$.

To prove the second inequality in Lemma 3.2 we start with the elementary fact

$$\begin{aligned} a^2t + (1-t)B^2 + t(1-t)D^2 &= B^2 - t(B^2 - a^2 - D^2) - t^2D^2 \\ &\geq B^2 - t(B^2 - a^2 - D^2) - t^2(a^2 + D^2) = (1-t)[B^2 + t(a^2 + D^2)] \end{aligned}$$

which yields

$$\{a^2t + (1-t)B^2 + t(1-t)D^2\}^{-\gamma+1} \leq (1-t)^{-\gamma+1} \{B^2 + t(a^2 + D^2)\}^{-\gamma+1}.$$

Multiplying both sides by $(1-t)^\beta$ and taking the integral from 0 to 1 with respect to the weight t^α we obtain the second claimed inequality in the lemma. Note that the exponent $\beta - \gamma + 1$ can be negative, but $\beta - \gamma + 1 > -1$ by our assumptions, so that integrability in $t = 1$ is guaranteed. Lemma 3.2 is proved. \square

4. Proof of Theorem 1.1: the general case

In this section we outline how to derive the estimate

$$(4.1) \quad \|P^*(\chi_E, \mu)\|_{L^k(\mathbb{R}^n)}^k \leq C^k (k!)^{\frac{3}{2}} n^{\frac{k-2}{2}} \kappa(\mu)^{k-1} \mu(E), \quad k = 4, 5, \dots$$

for some absolute constant $C > 0$, independent of n and k and for all μ -measurable subsets E of \mathbb{R}_+^{n+1} .

As a consequence of (4.1) we obtain

$$(4.2) \quad \|Pf\|_{L^{\frac{k}{k-1}, \infty}(\mathbb{R}_+^{n+1}, d\mu)} \leq C k^{\frac{3}{2}} n^{\frac{1}{2} - \frac{1}{k}} \kappa(\mu)^{\frac{k-1}{k}} \|f\|_{L^{\frac{k}{k-1}}(\mathbb{R}^n)}$$

for all $k = 3, 4, \dots$. These estimates extend (3.1) and (3.2), respectively. Once (4.2) is known, we deduce (1.8) as follows: for a given $p \in (1, 2)$ we find a positive integer $k_0 \geq 4$ such that $k_0/(k_0 - 1) < (p + 1)/2$. We use the Marcinkiewicz interpolation theorem to interpolate between estimate (1.5) (i.e. $p = 2$) and estimate (4.2) with $p = k_0/(k_0 - 1)$. Using a good value for the interpolation constant (see for instance the value of the constant in [G, p. 33]) we obtain (1.8) with $C_p = (p-1)^{-\frac{5}{2}}$. We note that the standard proofs of (1.1) yield the constant $c_{p,n} = C(p-1)^{-1}n^c$ (for some absolute $C > 0$ and $c > 0$), which grows much faster in n , but blows up at a slightly slower rate when n is fixed and $p \rightarrow 1$.

In order to work with (4.1) as in the case $k = 3$, we need a suitable generalization of Lemma 3.1 for arbitrary k . First, we have

$$\begin{aligned} & \int_{\mathbb{R}^n} \prod_{j=1}^k P_{a_j}(\tau - x_j) d\tau \\ &= \frac{a_1 a_2 \dots a_k}{\pi^{(k-1)\frac{n}{2} + \frac{k}{2}}} \int_{[0, \infty)^k} \frac{(u_1 u_2 \dots u_k)^{\frac{n-1}{2}}}{(u_1 + u_2 + \dots + u_k)^{\frac{n}{2}}} \\ & \quad \times \text{Exp} \left\{ - \sum_{j=1}^k a_j^2 u_j - \sum_{\ell=1}^{k-1} \frac{(u_1 + \dots + u_\ell) u_{\ell+1}}{u_1 + \dots + u_{\ell+1}} \right. \\ & \quad \left. \times \left| x_{\ell+1} - \frac{x_1 u_1 + \dots + x_\ell u_\ell}{u_1 + \dots + u_\ell} \right| \right\} du_1 \dots du_k \end{aligned}$$

which can be obtained from the n -dimensional identity (3.10) working in a recursive fashion as in the proof of Lemma 3.1, until the $k(k-1)/2$ distances appearing in the original formula (2.1) are “squeezed” together into $k-1$ distances. Note that in the case $k = 3$ we applied (3.10) just once.

Next, if Δ_k denotes the $(k-1)$ -dimensional simplex

$$\Delta_k = \left\{ v = (v_1, \dots, v_k) : v_j \geq 0, \sum_{j=1}^k v_j = 1 \right\}$$

then the change of variables $u = \rho v$ with $v \in \Delta_k$ and $\rho > 0$ yields

$$\begin{aligned} & \int_{\mathbb{R}^n} \prod_{j=1}^k P_{a_j}(\tau - x_j) d\tau \\ &= \frac{a_1 a_2 \dots a_k \Gamma\left(\frac{k-1}{2}n + \frac{k}{2}\right)}{\pi^{\frac{k-1}{2}n + \frac{k}{2}}} \int_{\Delta_k} (v_1 \dots v_k)^{\frac{n-1}{2}} \\ & \quad \times \left(\sum_{j=1}^k a_j^2 v_j + \sum_{\ell=1}^{k-1} \frac{(v_1 + \dots + v_\ell) v_{\ell+1}}{v_1 + \dots + v_{\ell+1}} \right. \\ & \quad \left. \times |x_{\ell+1} - x_1 v_1 + \dots + x_\ell v_\ell v_1 + \dots + v_{\ell+1}| \right)^{-\frac{k-1}{2}n - \frac{k}{2}} dv_1 \dots dv_k. \end{aligned}$$

The further change of variables $v_j = (1-t)s_j$, $j = 1, 2, \dots, k-1$, $v_k = t$, with $s = (s_1, \dots, s_{k-1}) \in \Delta_{k-1}$ and $t \in [0, 1]$ gives

$$\begin{aligned} \int_{\mathbb{R}^n} \prod_{j=1}^k P_{a_j}(\tau - x_j) d\tau &= \frac{a_1 a_2 \dots a_k \Gamma\left(\frac{k-1}{2}n + \frac{k}{2}\right)}{\pi^{\frac{k-1}{2}n + \frac{k}{2}}} \\ &\times \int_{\Delta_{k-1}} \int_0^1 \frac{(1-t)^{\frac{k-1}{2}n + k - 2} t^{\frac{n-1}{2}} (s_1 \dots s_{k-1})^{\frac{n-1}{2}} dt ds}{\left(a_k^2 t + (1-t)B_k^2 + t(1-t)|x_k - q_k|^2\right)^{\frac{k-1}{2}n + \frac{k}{2}}} \end{aligned}$$

with

$$q_k = x_1 s_1 + \dots + x_{k-1} s_{k-1}$$

$$B_k^2 = \sum_{j=1}^{k-1} a_j^2 s_j + \sum_{\ell=1}^{k-2} \frac{(s_1 + \dots + s_\ell) s_{\ell+1}}{s_1 + \dots + s_{\ell+1}} \left| x_{\ell+1} - \frac{x_1 s_1 + \dots + x_\ell s_\ell}{s_1 + \dots + s_{\ell+1}} \right|,$$

and this is a suitable generalization of Lemma 3.1, with the property that B_k is independent of t , x_k , a_k .

We now apply Lemma 3.2 with $\alpha = \frac{n-1}{2}$, $\beta = \frac{k-1}{2}n + k - 2$, $\gamma = \frac{k-1}{2}n + \frac{k}{2}$ and proceed as we did in the previous case $k = 3$, assuming $a_k \leq a_{k-1} \leq \dots \leq a_1$. We obtain

$$\begin{aligned} \int_{\mathbb{R}^n \times (0, a_{k-1})} a_k \int_0^1 \frac{t^{\frac{n-1}{2}} (1-t)^{\frac{k-1}{2}n + k - 2}}{\left(a_k^2 t + (1-t)B_k^2 + t(1-t)|x_k - q_k|^2\right)^{\frac{k-1}{2}n + \frac{k}{2}}} dt d\mu(x_k, a_k) \\ \leq \frac{\kappa(\mu) \pi^{n/2+1} \left(\frac{k-1}{2}n + \frac{k}{2} - 1\right) \Gamma\left(\frac{(k-2)(n+1)}{2}\right)}{\left(B_k^2\right)^{\frac{k-2}{2}n + \frac{k+1}{2}} \Gamma\left(\frac{(k-1)n}{2} + \frac{k}{2}\right)} \\ = \frac{\kappa(\mu) \pi^{n/2+1} \Gamma\left(\frac{(k-2)(n+1)}{2}\right)}{\left(B_k^2\right)^{\frac{k-2}{2}n + \frac{k+1}{2}} \Gamma\left(\frac{(k-1)(n+1)}{2} - \frac{1}{2}\right)}. \end{aligned}$$

Iterating this estimate $k - 3$ more times, and doing one last estimate as in [V, equation (13)] we get, for any $k \geq 4$, that

$$\begin{aligned} & \frac{1}{k!} \int_{E^{k-1}} \int_{\mathbb{R}^n} \prod_{j=1}^k P_{a_j}(\tau - x_j) d\tau dx_2 \dots dx_k \\ & \leq \frac{C\kappa(\mu)^{k-1}}{\left(\frac{k-1}{2}n + \frac{k}{2} - 1\right)^{-1}} \frac{\Gamma\left(\frac{(k-2)(n+1)}{2}\right)}{\Gamma\left(\frac{(k-2)(n+1)}{2} - \frac{1}{2}\right)} \frac{\Gamma\left(\frac{(k-3)(n+1)}{2}\right)}{\Gamma\left(\frac{(k-3)(n+1)}{2} - \frac{1}{2}\right)} \dots \\ & \dots \frac{\Gamma(n+1)}{\Gamma\left(n + \frac{1}{2}\right)} \frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n}{2} + 1\right)} \end{aligned}$$

which is of the order of

$$\kappa(\mu)^{k-1} \sqrt{(k-1)!} (n+1)^{\frac{k-3}{2} + \frac{1}{2}} \sim C^k \kappa(\mu)^{k-1} \Gamma(k)^{\frac{1}{2}} n^{\frac{k-2}{2}}.$$

Since this estimate is uniform with respect to (x_1, a_1) we obtain (4.1).

To prove (1.9) we first note that applying the Marcinkiewicz interpolation theorem between (1.5) (i.e. $p = 2$) and the trivial $p = \infty$ estimate yields

$$(4.3) \quad \|Pf\|_{L^p(\mathbb{R}_+^{n+1}, d\mu)} \leq c' p^{\frac{1}{p}} (p-2)^{-\frac{1}{p}} \kappa(\mu)^{\frac{1}{p}} \|f\|_{L^p(\mathbb{R}^n)},$$

for all $p > 2$. Finally, using the Riesz-Thorin interpolation theorem, we interpolate between (1.8) with $p = \frac{3}{2}$ and (4.3) with $p = \left(\frac{1}{2} - \frac{1}{\log n}\right)^{-1}$ ($n > 10$) to obtain (1.9).

References

- [A] M. ANDERSSON, “*Topics in complex analysis*”, Universitext, Springer-Verlag, New York, 1997.
- [C1] L. CARLESON, An interpolation problem for bounded analytic functions, *Amer. J. Math.* **80** (1958), 921–930.
- [C2] L. CARLESON, Interpolations by bounded analytic functions and the corona problem, *Ann. of Math. (2)* **76** (1962), 547–559.
- [G] L. GRAFAKOS, “*Classical and Modern Fourier Analysis*”, Pearson Education, Upper Saddle River, New Jersey, 2004.
- [H] L. HÖRMANDER, L^p estimates for (pluri-) subharmonic functions, *Math. Scand.* **20** (1967), 65–78.
- [N] N. K. NIKOL'SKIĬ, “*Treatise on the shift operator*”, Spectral function theory, With an appendix by S. V. Hruščev and V. V. Peller, Translated from the Russian by Jaak Peetre, Grundlehren der

- Mathematischen Wissenschaften **273**, Springer-Verlag, Berlin, 1986.
- [S] E. M. STEIN, “*Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*”, With the assistance of Timothy S. Murphy, Princeton Mathematical Series **43**, Monographs in Harmonic Analysis III, Princeton University Press, Princeton, NJ, 1993.
- [SW] E. M. STEIN AND G. WEISS, “*Introduction to Fourier analysis on Euclidean spaces*”, Princeton Mathematical Series **32**, Princeton University Press, Princeton, N.J., 1971.
- [V] I. E. VERBITSKY, A dimension-free Carleson measure inequality, in: “*Complex analysis, operators, and related topics*”, Oper. Theory Adv. Appl. **113**, Birkhäuser, Basel, 2000, pp. 393–398.

Loukas Grafakos:
Department of Mathematics
University of Missouri
Columbia, Missouri 65211
U.S.A.
E-mail address: `loukas@math.missouri.edu`

Enrico Laeng:
Dipartimento di Matematica
Politecnico di Milano
Piazza L. da Vinci 32
20133 Milano
Italy
E-mail address: `enrlae@mate.polimi.it`

Carlo Morpurgo:
Department of Mathematics
University of Missouri
Columbia, Missouri 65211
U.S.A.
E-mail address: `morpurgo@math.missouri.edu`

Primera versió rebuda el 25 d'abril de 2006,
darrera versió rebuda el 3 d'octubre de 2006.