

**RESTITUCIÓ FUNCIONAL I
ESPECIFICITAT DE LA REINNERVACIÓ
DESPRÉS DE LESIONS I REPARACIONS
DEL NERVI PERIFÈRIC**

Antoni Valero Cabré

Tesi Doctoral

1999

VI. Conclusions

- 1. Després de lesions de compressió, secció o resecció del nervi ciàtic de la rata, els axons tenen capacitat per regenerar i reinnervar dianes musculars, sensorials i sudomotores i permetre el restabliment de respostes funcionals.**

A excepció de les lesions de compressió (grau II de Sunderland), la regeneració axonal en lesions de grau V de Sunderland assolix nivells de restitució funcional inferior a la dels valors preoperatoris o d'un grup control no lesionat. Els mètodes electrofisiològics i funcionals utilitzats permeten mesurar la progressió de la reinnervació i la recuperació funcionals per part de diferents tipus de fibres nervioses ($A\alpha$, $A\beta$, $A\delta$ i C) simultàniament.

- 2. L'inici i el nivell final de la regeneració de diferents tipus de fibres nervioses després de lesions depèn de la gravetat de la lesió i de l'eficàcia del mètode reparatiu aplicat.**

La compressió nerviosa mostra una bona recuperació fins a valors semblants als preoperatoris. Els gups de resecció mostren nivells de reinnervació distals inferiors als dels grups de secció. La reparació per sutura directa és la millor tècnica reparativa de troncs nerviosos plurifasciculars i mixtes, sempre i quan es pugui garantir un correcte alineament. La tubulització en distàncies interneurals curtes és una bona alternativa quan hi hagi risc de realitzar sutures a tensió o alterar l'orientació fascicular. En lesions de resecció i distàncies interneurals mitges/llargues la tubulització única o fascicular amb guies de silicona obté pitjors resultats que la reparació per un autoinjert ideal. La tubulització amb guies semipermeable de PLC és una bona alternativa a l'empelt.

- 3. Les fibres nervioses mielíniques i amielíniques manifesten diferent capacitat de regeneració i reinnervació distals.**

Les funcions mediades per fibres primes amielíniques nociceptives o sudomotores (C) mostren en tots els grups una reinnervació més ràpida i assolixen nivells finals de recuperació superiors a les mediades per fibres mielíniques gruixudes motores ($A\alpha$) i sensorials ($A\beta$). La recuperació de funcions mediades per fibres primes no permet discriminar entre els efectes de procediments reparatius diversos, que sí presenten valors diferents de reinnervació muscular i sensorial per fibres gruixudes.

- 4. No totes les dianes musculars i sensorials perifèriques tenen la mateixa capacitat de ser reinnervades després de lesions.**

La reinnervació de dianes distals té lloc de manera més tardana i en menor grau que la dels òrgans diana proximals. El múscul plantar i els nervis digitals mostren dies d'inici més tardans i valors finals inferiors de recuperació que la dels músculs gastrocnemi i tibial anterior i la del nervi tibial a nivell del taló. La recuperació de la funció motora i sensorial a nivells proximals no discrimina entre grups sotmesos a diferents tipus de reparacions que mostren per contra, nivells significativament diferents de reinnervació distal.

5. L'assoliment d'uns bons nivells de reinnervació motora i sensorial a l'extremitat posterior no garanteix la recuperació de la funció global de locomoció.

Només després de lesions compresions s'observa una recuperació significativa del patró de locomoció. Les lesions de secció o resecció alteren de manera no reversible la activitat locomotora, independentment de l'èxit d'alguns procediments reparatius com a promotors de la reinnervació muscular i sensorial.

6. Després de lesions de nervis plurifasciculars, el patró de reinnervació no reproduïx el mapa original de projeccions d'axons motors, sensorials i sudomoters.

Tots els grups experimentals mostren, en diferents graus, reinnervació aberrant de dianes per axons de fascicles no corresponents, neurones motores amb dobles o triples projeccions axonals en diversos músculs i poliinnervació de fibres musculars. Noranta dies després de la lesió, els nivells de reinnervació fascicular aberrant no superen, però, en cap grup els nivells de reinnervació funcional correcta. Els mètodes electrofisiològics i histològics utilitzats permeten quantificar les alteracions que la lesió produeix sobre l'especificitat de la reinnervació, el grau d'hiperinnervació de dianes i el nombre de motoneurons amb projeccions múltiples funcionals.

7. El grau de reinnervació funcional aberrant, innervació múltiple i poliinnervació de dianes depèn del tipus de lesió i del mètode reparatiu utilitzat

Les lesions de compressió presenten la reinnervació més específica de tots els grups estudiats, encara que no lliure d'alguns errors de la reinnervació. Les lesions de secció mostren percentatges de reinnervació funcional aberrant, dobles projeccions musculars i poliinnervació inferiors als dels grups de resecció. En lesions de secció, la sutura directa ben alinada és la reparació de referència, encara que la tubulització origina menys errors de reinnervació que la sutura equivocada amb desalineament fascicular. En distàncies interneurals mitges/llargues, la tubulització fascicular redueix de manera significativa el grau de reinnervació aberrant i de múltiples projeccions. La tubulització amb guies de PLC redueix significativament els nivells de reinnervació muscular aberrant i el nombre de motoneurons amb múltiples projeccions respecte a la reparació per empelt o la tubulització amb guies de silicona.

8. Els diferents tipus de fibres nervioses no mostren diferències en la seva capacitat per reinnervar específicament dianes perifèriques.

La major capacitat de regeneració i reinnervació de les fibres nervioses amielíniques no comporta un increment del percentatge de reinnervació funcional aberrant, hiperinnervació o poliinnervació respecte a les mielíniques. No hi ha diferències entre els nivells de reinnervació funcional aberrant de dianes proximals o distals, però les primeres mostren nivells de poliinnervació muscular superiors i les segones d'hiperinnervació polineuronal.

9. La regeneració i la reinnervació comporten una ràpida restitució de les respostes reflexes monosinàptiques i polisinàptiques després de lesions perifèriques.

Les lesions del nervi perifèric comporten l'abolició de les respostes reflexes espinals. La reinnervació sensorio-motora restitueix en tots els grups experimentals els circuits del reflex monosinàptic ($A\alpha$) i dels components inicials del reflex polisinàptic (C1 i C2, mediat per fibres sensorials $A\beta$ i $A\delta$). El component C3, mediat per fibres C, es recupera de manera més lenta en un proporció inferior d'animals.

10. El temps de conducció dels reflexos monosinàptics i polisinàptics s'incrementa amb la lesió i es recupera amb la reinnervació fins a valors finals propers als originals.

En el reflex monosinàptic, els grups de secció i els músculs proximals mostren una recuperació del temps de conducció més enlentida que la dels grups de resecció i els músculs distals, però sense diferències significatives al final del seguiment. En els reflexos polisinàptics, no hi ha diferències entre els nivells finals de recuperació del temps de conducció dels diferents components (C1, C2, C3) ni entre diferents tipus de lesió i reparacions.

11. Les lesions perifèriques indueixen una facilitació de respostes reflexes monosinàptiques i polisinàptiques que disminueix a mesura que té lloc la reinnervació distal

La relació d'amplituds entre l'ona M i l'ona H i l'amplitud màxima dels components C1, C2 i C3 dels reflex polisinàptic s'incrementen respecte dels valors controls en les primeres fases posteriors a la lesió i tenen tendència a normalitzar-se a mesura que progressa la reinnervació. Els components reflexos mediat per fibres gruixudes mielíniques i els circuits reflexos amb dianes musculars distals mostren nivells de facilitació superiors als mediat per fibres amielíniques o als de grups musculars proximals.

12. El tipus de lesió i de reparació condicionen la involució de la facilitació de reflexes monosinàptics i polisinàptics induïda per la lesió perifèrica

La facilitació de l'activitat reflexa i l'increment de la resposta és dependent del grau i la qualitat de la innervació muscular i sensorial en cada moment del seguiment. Al final, el grup de compressió mostra valors d'amplitud més propers als preoperatoris. Els grups de secció i de resecció no mostren diferències significatives en el grau de màxima facilitació de respostes reflexes en les primeres fases de la reinnervació. La reparació per sutura en distàncies interneurals curtes i l'empelt i el tub de PLC en lesions de resecció mostren al final del seguiment valors de resposta reflexa més ajustats als preoperatoris que la resta de grups homòlegs. La recuperació de la facilitació dels reflexos polisinàptics és inferior a la dels monosinàptics.

VII. Bibliografia

- Abernethy DA, Rud A, Thomas PK. Neurotrophic influence of the distal stump of transected peripheral nerve on axonal regeneration: absence of topographic specificity in adult nerve. *J Anat* 1992, 180:395-400.
- Abernethy DA, Thomas PK, Rud A, King RHM. Mutual attraction between emigrant cells from transected denervated nerve. *J Anat* 1994, 184:239
- Aguayo AJ, Epps J, Charron LC, Bray GN. Multipotentiality of Schwann cells in cross anastomosed and grafted myelinated and unmyelinated nerves. *Brain Res* 1976, 104: 1-20
- Aldskogius H, Arvidsson J, Grant G. Axotomy induced changes in primary sensory neurons. A Scott SA (ed) *Sensory Neurons Diversity: Development and Plasticity*. pp 363-383. Oxford Univ. Press 1992, New York, Oxford.
- Aldskogius H, Kozlova E. Central neuron-glial and glial-glial interactions following axon injury. *Progr Neurobiol* 1998, 55:1-26
- Aldskogius H, Molander C. Specificity in regenerative outgrowth and target reinnervation by mammalian peripheral axons. *Rest Neurol Neurosci* 1990, 1:275-280
- Amir R, Devor M. Axonal cross-excitation in nerve-end neuromas: comparison of A- and C-fibers. *J Neurophysiol* 1992 68(4):1160-1166
- Andrew D, Greenspan JD. Modality-specific hyper-responsivity of regenerated cat cutaneous nociceptors. *J Physiol* 1999, 516(3):89-906
- Angelov DN, Gunkel A, Stennert E, Neiss WF. Phagocytic microglia during delayed neuronal loss in the facial nucleus of the rat: time course of the neuronofugal migration of brain macrophages. *Glia* 1995, 13:113-129
- Angelov DN, Gunkel A, Stennert E, Neiss WF. Recovery of original nerve supply after hypoglossal-facial anastomosis causes permanent motor hyperinnervation of the whisker-pad muscles in the rat. *J Comp Neurol* 1993, 338:214-224.
- Angelov DN, Neiss WF, Streppel M, Andermahr J, Mader K, Stennert E. Nimodipine accelerates axonal sprouting after surgical repair of rat facial nerve. *J Neurosci* 1996, 16(3): 1041-1048
- Angelov DN, Skouras E, Guntinas-Lichius O, Streppel M, Popratiloff A, Walther M, Klein J, Stennert E, Neiss WF. Contralateral trigeminal nerve lesion reduces polyneuronal muscle innervation after facial nerve repair in rats. *Eur J Neurosci* 1999, 11:101-110
- Archibald SJ, Krarup C, Shefner J, Li ST, Madison RD. A collagen-based nerve guide conduit for peripheral nerve repair: an electrophysiological study of nerve regeneration in rodents and nonhuman primates. *J Comp Neurol* 1991, 306: 685-696.
- Archibald SJ, Shefner J, Krarup C, Madison RD. Monkey median nerve repaired by nerve graft or collagen nerve guide tube. *J Neurosci* 1995, 15(5):4109-4123
- Bain JR, Mackinnon S, Hunter R. Functional evaluation of complete sciatic, peroneal, and posterior tibial nerve lesions in the rat. *Plastic Reconstr Surg* 1989, 83:129-189
- Bamburg JR, Lockerbie RO, Bernstein BW, Morgan TE, Whitney M, Adams ME. Actin dynamics in neuronal growth cones. A Letourneau PC, Kater SB, Macagno ER (eds). *The nerve growth cone*. Raven Press, New York, 1992, pp 27-38.
- Baños JE, Verdú E, Butí M, Navarro X. Effects of dizocilpine on autotomy behavior after nerve section in mice. *Brain Res* 1994, 636:107-110
- Baranauskas G, Nistri A. Sensitization of pain pathway in the spinal cord: cellular mechanisms. *Progr Neurobiol* 1998, 54:349-365.
- Ben Johnston R, Zachary L, Dellon AL, Seiler WA, Teplica DM. Improved imaging of rat hindfoot prints for walking track analysis. *J Neurosci Meth* 1991, 38:111-114.
- Benett G, Kajander KC, Sahara Y, Iadarla MJ, Sugimoto T. Neurochemical and anatomical changes in the dorsal horn of rats with an experimental painful peripheral neuropathy. A Cerveró, Benett GJ, Headley PM (eds). *Processing of sensory information in the superficial dorsal horn of the spinal cord*. Plenum 1989, pp.463-472, New York
- Bento RF, Miniti A. Anastomosis of the intratemporal facial nerve using fibrin tissue adhesive. *Ear. Nose. Throat. J* 1993, 72:663

- Bernstein JJ, Guth L. Nonselectivity in establishment of neuromuscular connections following nerve regeneration in the rat. *Exp Neurol* 1961, 4:262-275
- Bharali LA, Burgess SA, Lisney SJ, Pearson D. Reinnervation of sweat glands in the rat hind paw following peripheral nerve injury. *J Auton Nerv Syst* 1988, 23:125-129
- Biela G, Riva L, Sotgiu ML. Interaction between neurons in different laminae of the dorsal horn of the spinal cord. A correlation study in normal and neuropathic rats. *Eur J Neurosci* 1997, 9:1017-1025
- Bisby JL. Regeneration of peripheral nerve system axons. A Waxman SG, Kocsis JD, Stys PK (eds). *The Axon: Structure, Function and Pathophysiology*. Oxford University Press, New York, Oxford. 1995, pp. 355-374
- Blinzinger K, Kreutzberg GW. Displacement of synaptic terminals from regenerating motoneurons by microglial cells. *Z Zellforsch* 1986, 85:145-157
- Bolin LM and Shooter EM. Neurons regulate Schwann cell genes by diffusible molecules. *J Cell Biol* 1993, 123: 237-243.
- Bowe MA, Fallon JR. The role of agrin in synapse formation. *Annu Rev Neurosci* 1995, 18: 443-462.
- Boyd IA, Smith S. The muscle spindle. A Dyck PJ, Thomas PK, Lambert EH i Bunge R (eds). *Peripheral neuropathy*. pp 171-202. Saunders 1984, Philadelphia.
- Brännstrom T, Havton L, Kellerth JO. Changes in size and dendritic arborization patterns of adult cat spinal α -motoneurons following permanent axotomy. *J Comp Neurol* 1992a, 318:439-451
- Brännstrom T, Havton L, Kellerth JO. Restorative effects of reinnervation on the size and dendritic arborization patterns of axotomized cat spinal a-motoneurons. *J Comp Neurol* 1992b. 318:452-461
- Bray D. Cytoskeletal basis of nerve axon growth. A Letorneau PC, Kater SB, Macagno ER (eds). *The nerve growth cone*. Raven Press, New York, 1992, pp 7-17.
- Bregman BS, Kunke-Bagden E, Schnell L, Dai HN, Gao D, Schwab ME. Recovery from spinal cord injury mediated by antibodies to neurite growth inhibitors. *Nature* 1995, 378:498-501.
- Bridgman PC. Functional anatomy of the growth cone in relation to its role in locomotion and neurite assembly. A Letorneau PC, Kater SB, Macagno ER (eds). *The nerve growth cone*. Raven Press, New York, 1992, pp 39-53.
- Brookes JP. Assays for cholinergic properties in cultured rat Schwann cells. *Proc R Soc London Ser B* 1984, 222: 121-134.
- Brown AG, Fyffe REW. The morphology of group Ia afferent fibre collaterals in the spinal cord of the cat. *J Physiol (London)* 1978, 111-127
- Brown CJ, Evans PJ, Mackinnon SE, Bain JR, Makino AP, Hunter RT, hare G. Inter and intraobserver reliability of walng-track analysis used to assess sciatic nerve function in rats. *Microsurg* 1991, 12:76-79.
- Brown CJ, Mackinnon, Evans PJ, Bain JR, Makino AP, Hunter RT, Hare GMT. Self evaluation of walking track measurement using a sciatic function index. *Microsurg* 1989, 10:226-235
- Brown MC, Hardman VJ. A reassessment of the accuracy of reinnervation by motoneurons following crushing or freezing of the sciatic nerve or lumbar spinal nerves of the rat
- Brown TG. The intrinsic factors in the act of progression in the mammal. *Proc R Soc London* 1911, 84:308-319
- Brushart TM. Trophic and tropic influences on peripheral nerve regeneration. A Omer GE, Spinner M, Van Beek AL(eds.). *Management of peripheral nerve problems*. 2nd edition, W.B Saunders Company 1998.
- Brushart TM, Seiler WA. Selective reinnervation of distal motor stumps by peripheral motor axons. *Exp Neurol* 1987, 97:290-300
- Brushart TME, Gerber J, Kessens P, Chen YG, Royall, RM. Contributions of pathway and neuron to preferential motor reinnervation. *J Neurosci* 1998, 18(21):8674-8681.
- Brushart TME, Henry EW, Mesulam M. Reorganization of muscle afferent projections accompanies peripheral nerve regeneration. *Neuroscience* 1996, 2053-2061

- Brushart TME, Mathur V, Koschorke GM, Sood R. Trajectory of regenerating axons across enclosed gaps. *J Hand Surg* 1995.
- Brushart TME, Seiler WA. Selective reinnervation of peripheral distal stumps by peripheral motor axons. *Exp Neurol* 1987, 97:289-300
- Brushart TME. Motor axons prefferentially reinnervate motor pathways. *J Neurosci* 1993, 13:2730-2738
- Brushart TME. Preferential motor reinnervation: a sequential double-labeling study. *Restor Neurol Neurosci* 1990, 1:281-287
- Brushart TME. Preferential reinnervation of motor nerves by by regenerating motor axons. *J Neurosci* 1988, 8:248-253
- Brushart TME. The mechanical and humoral control of specificity in nerve repair. A Gelberman RH (ed): *Operative nerve repair and reconstruction*. Philadelphia, J.B Lippincott Company 1991, p. 215
- Brushart TMH, Mesulam MM. Alterations in in connections between muscle and and anterior horn motoneurons after peripheral nerve repair. *Science* 1980, 208:603-605.
- Buchtahl F, Kuhl V. Nerve conduction, tactile sensibility, and the electromyogram after suture or compression of peripheral nerve: a longitudinal study in man. *J Neurol Neurosurg Psychiat* 1979, 42:436-451.
- Bunge MB, Johnson MI, Ard MD, Kleitman N. Factors influencing the growth of regenerating nerve fibers in culture. *Progr Brain Res* 1987, 71: 61-74.
- Butí M, Verdú E, Labrador RO, Vilches JJ, Forés J, Navarro X. Influence of Physical Parameters of Nerve Chambers on Peripheral Nerve Regeneration and Reinnervation. *Exp Neurol* 1996, 137:26-33.
- Butí M. Reparació de nervi perifèric per tubulització. Tesi doctoral. Universitat Autònoma de Barcelona. Barcelona 1994
- Cameron AA, Cliffer KD, Dougherty PM, Garrison CJ, Willis WD, Carlton SM. Time course of degenerative and regenerative changes in the dorsal horn in a rat model of peripheral neuropathy. *J Comp Neurol* 1997, 379:428-442
- Canedo A. Mèdula espinal, motoneuronas y unidades motoras; reflejos espinales. A Delgado JM, Ferrús A, Mora F, Rubia F (eds.). *Manual de Neurociencia*. Litoral 1998, Madrid
- Cannon A. Law of denervation. *Am J Med Sci* 1939, 198:739-750
- Caroni P. Driving the growth cone. *Science* 1998, 281:1465-1466.
- Carrier L, Brunstein E, Rossignol S. Locomotion of the hindlimb after neurectomy of the ankle flexors in intact and spinal cats: model for the study of locomotor plasticity. *J Neurophysiol* 1997. 77:1979-1993.
- Castro-Lopes JM, Malcangio M, Pan BH, Bowery NG. Complex changes of GABA_A and GABA_B receptor binding after in the spinal cord dorsal horn following peripheral inflammation or neurectomy. *Brain Res* 1995, 679:289-297
- Castro-Lopes JM, Távora I, Coimbra A. GABA decreases in the spinal cord dorsal horn after peripheral neurectomy. *Brain Res* 1993, 620:287-291
- Ceballos D, Navarro X, Dubai N, Wendelschafer-Crabb G, Kennedy WR, Tranquillo RT. Nerve regeneration using a magnetically aligned gel. *Exp Neurol* 1999, 158:290-300.
- Chen H, He Z, Tessier-Lavigne M. Axon guidance mechanisms: semaphorins as simultaneous repellents and antirepellents. *Nature Neurosci* 1998, 1(6):436-439.
- Chew LJ, Gallo V. Regulation of ion channel expression in neural cells by hormones and growth factors. *Mol Neurobiol* 1999, 18(3):175-255
- Chiu AY, Espinosa de los Monteros A, Cole RA, Loera S, de Vellis J. Laminin and s-laminin are produced and released by astrocytes, Schwann cells and schwannomas in culture. *Glia* 1991, 4: 11-24.
- Clare MH, Landau W. Distal hind-limb reflex responses in cats with largely intact spinal reflex circuits. *Exp Neurol* 1975, 46:470-495

- Collins WF, Honig G, Mendell LM. Heterogeneity of group Ia synapses on homonymous a-motoneurons as revealed by high frequency stimulation of Ia afferent fibers. *J Neurophysiol* 1984, 52(5): 980-993
- Cope TC, Bonasera J, Nichols TR. Reinnervated muscles fail to produce stretch reflexes. *J Neurophysiol* 1994, 71(2):817-820
- Cope TC, Clarck BD. Motor-Unit recruitment in self reinnervated muscle. *J Neurophysiol* 1993, 70(5):1787-1796
- Cossu G, Valls-Solé J, Valldeoriola F, Muñoz E, Benítez P, Aguilar F. Reflex excitability of facial motoneurons at onset of muscle reinnervation after facial nerve palsy. *Musc Nerv* 1999, 22:614-620
- Cova JL, Aldskogius H. Effect of axotomy on perineuronal glial cells in the hypoglossal and dorsal motor vagal nucleus of the cat. *Exp Neurol* 1986, 93: 662-667
- Covault J, Sanes JR. Neural adhesion molecule (N-CAM) accumulates in denervated and paralyzed skeletal muscles. *Proc Natl Acad Sci USA* 1985, 82:4544-4548
- Cragg BG, Thomas PK. Changes in conduction velocity and fibre size proximal to peripheral nerve lesions. *J Physiol (London)* 1961. 157:315-327
- Cragg BG, Thomas PK. The conduction velocity of regenerated peripheral nerve fibres. *J Physiol* 1964, 171:164-175
- Cragg BG. What is the signal for chromatolysis?. *Brain Res* 1970, 23: 1-21.
- Curtis R, Scherer SS, Somogyi R, Adryan KM, Ip NY, Zhu Y, Lindsay RM, DiStefano PS. Retrograde axonal transport of LIF is increased by peripheral nerve injury; correlation with increased LIF expression in the distal nerve. *Neuron* 1994, 12: 191-204.
- Dassesse D, Cuvelier L, Krebs C, Streppel M, Gunthinas Lichius O, Neiss W, Pochet R. Differential expression of calbindin and calmodulin in motoneurons after hypoglossal axotomy. *Brain res* 1998, 786:181-188.
- Davey DF, Brown CS, Ansselin AD. Pathway selection by regeneration peripheral nerve axons growing through a Y-shaped tube is related to distal stump size. *Soc Neurosci Abstr* 1992, 18:609
- Davis BM, Collins WF, Mendell LM. Potentiation of transmission at Ia-motoneuron connections induced by repeated short bursts of afferent activity. *J Neurophysiol* 1985, 54(6): 1541-1551
- Dawson DM, Krarup C. Perioperative nerve lesions. *Arch Neurol* 1989, 46: 1355-1360.
- De la Cruz R, Pastor AM, Delgado-Garcia JM. Effects of target depletion on adult mammalian central neurons functional correlates. *Neurosci* 1994, 58(1):81-97
- De la Cruz R, Pastor AM, Delgado-Garcia JM. Influence of the postsynaptic target on the functional properties of neurons in the adult mammalian central nervous system. *Rev Neurosci* 1996, 7:115-149
- De Medinaceli L, Freed WJ, Wyatt RJ. An index of the functional condition of rat sciatic nerve based on measurements made from walking tracks. *Exp Neurol* 1982, 77: 634-643.
- De Medinaceli L. Functional consequences of experimental nerve lesions: effects of reinnervation blend. *Exp Neurol* 1988, 100: 166-178.
- De Santis, Berger PK, Laskowski MB, Norton A. Regeneration by skeletomotor axons in neonatal rats is topographically selective at an early stage of reinnervation. *Exp Neurol* 1992, 116:229-239.
- De Santis, Norman W. Location and completeness of reinnervation by two types of neurons at a single target: the feline muscle spindle. *J Comp Neurol* 1993, 336:66-76
- Dellon AL, Mackinnon SE. An alternative to the classical nerve graft for the management of the short nerve gap. *Plastic Reconstruct Surg* 1988, 82: 849-856.
- Dellon AL. Evaluation of sensibility and reeducation of sensation in the hand. Baltimore, Williams and Wilkins 1981.
- Dellon ES, Dellon AL. Functional assessment of neurologic impairment: track analysis in diabetic and compression neuropathies. *Plast Reconstructiv Surg* 1991, 88(4):686-694

- Dellon ES, Mackinnon SE. Sciatic nerve regeneration in the rat. Validity of walking track assessment in the presence of cronic contractures. *Microsurg* 1989, 10:220-225.
- Denn Dunen WFA, Van der Lei B, Robinson PH, Holwerda A, Pennings AJ, Schakenraad JM. Biological performance of a degradable poly(lactic acid-e-caprolactone) nerve guide. Influence of tube dimensions. *J Biomed Mat Res* 1995, 29:757-766
- Devor M, Govrin-Lippmann, Angellides K. Na⁺ channel immunolocalization in prepipheral mamalian axons and changes following nerve injury and neuroma formation. *J Neurosci* 1993, 13(5):1976-1992
- Devor M, Govrin-Lippmann R. Selectie Regeneration of sensory fibers following nerve crush injury. *Exp neurol* 1979. 65:243-254.
- Devor M, Keller CH, Deerinck TJ, Ellisman MH. Na⁺ channels accumulation on axolemma of afferent endings in nerve end neuromas in *Apteronotus*. *Neurosci. Lett* 1989, 102:149-154
- Devor M, Schonfeld D, Seltzer Z, Wall PD. Two modes of cutaneous reinnervation following peripheral nerve injury. *J Comp Neurol* 1979, 185:211-220.
- Devor M, Wall PD. Type of sensory nerve fibre sprouting to form a neuroma. *Nature* 1976, 262:705-708
- Devor M. The patophysiolgy of damaged peripheral nerves. A Wall PD, Melzack R (eds.). *Textbook of pain* (3rd edition). Churchill Livingstone 1994, pp. 79-100, Edinburgh
- DeVrind HH, Van Dam WM, Wondergem J, Haveman J. Latent X-ray damage in the rat sciatic nerve results in delay in functional recovery after a heat treatment. *Int J Radiat Biol* 1993, 63: 83-89.
- DeVrind HH, Wondergem J, Haveman J. Hyperthermia-induced damage to rat sciatic nerve assessed in vivo with functional methods and with electrophysiology. *J Neurosci Meth* 1992, 45: 165-174.
- Di Giamberardino L, Coraud J, hassig R, Marini P, Gorio A. The use of Acetyl cholineestrase molecular forms to appraise post-traumatic recovery of nerve and muscle. A Gorio A, Millesi H, Migrino S (eds.). *Post-traumatic peripheral nerve regeneration*. Raven Press 1981, pp. 481-492, New York.
- Dib-Hajj S, Black JA, Cummins TR, Kenney AM, Kocsis JD, Waxman SG. rescue of a-SNS sodium chanel expression in small dorsal root ganglion neurons after axotomy by nerve growth factor in vivo. *J Neurophysiol* 1998, 79:2668-2675
- Dietz V, Colombo G, Jensen L. Locomotor activity in spinal man. *Lancet* 1994, 344:1260-1263
- Distefano PS, Friedman B, Raziejewski C, Alexander C, Boland P, Schick CN, Lindsay RM, Wiegand SJ. The neurotrophins BDNF, NT-3 and NGF display distincts patterns of retrograde axona transport in peripheral and central neurons. *Neuron* 1992, 8:983-993
- Doetsch G, Harrisom T, McDonald A, Litaker M. Short term plasticity in primary somatosensory cortex of the rat: rapid changes in magnitudes and latencies of neuronal responses following digit denervation. *Exp Brain Res* 1996, 112:505-512
- Donoghue JP, Suner S, Sanes JN. Dynamic organization of primary motor cortex output to target muscles in adult rats. II. Rapid reorganization following motor nerve lesions. *Exp Brain Res* 1990,70:492-503
- Doolabh V, Hertl MC, Mackinnon SE. The role of conduits in nerve repair. *Reviews in neurosciences* 1996, 7:47-84
- Dorner H, Otte P, Platt D. Trainning influence on age dependent changes in the gait of rats. *Gerontology* 1996, 42:7-13
- Dovstrovsky JO, Millar J, Wall PD. The immediate shift of afferent drive of dorsal column nucleus cells following deafferentation: a comparison of acute and chronic deafferentation in gracile nucleus and spinal cord. *Exp Neurol* 1976, 562:480-495.
- Dubovy P, Aldskogius H. Degeneration and regeneration of cutaneous sensory nerve formations. *Micr Res Tech* 1996, 34:362-375
- Dubovy P, Svizenska I. Denervated skeletal muscle stimulates migration of Schwann cells from the distal stump of transected peripheral nerve: an in vivo study. *Glia* 1994, 12: 99-107.

- Dyck PJ. The causes, classification, and treatment of peripheral neuropathy. *New Eng J Med* 1982, 307: 283-286.
- Eccles JC, Libet B, Young R. The behaviour of chromatolysed motoneurons studied by intracellular recording. *J Physiol (Lond)* 1958, 143:11-40
- Edshage S. Peripheral nerve suture. A technique for improved intraneural topography. *Acta Chirurg Scand.* 1964. Suppl 331: 1-104.
- Eriksson NP, Persson JKE, Arvidsson J, Molander C, Aldskogius H. A, Svensson M. Quantitative analysis of the microglial cell reaction in central primary sensory termination areas and treatment with nerve growth factor following peripheral nerve injury in the adult rat. *Exp Brain Res* 1997, 114:393-404.
- Evans PJ, Bain JR, Mackinnon SE, Makino AP, Hunter DA. selective reinnervation: a comparison of recovery following microsuture and conduit nerve repair. *Brain Res* 1991, 559:315-321
- Evans PJ, Midha R, Mackinnon SE. The peripheral nerve allograft: a comprehensive review of regeneration and neuroimmunology. *Prog Neurobiol* 1994, 43: 187-233.
- Farel PB. Selective synaptic changes following spinal motoneuron axotomy. *Brain Res* 1980, 189: 67-77.
- Fawcett JW, Keynes RJ. Peripheral nerve regeneration. *Annu Rev Neurosci* 1990, 13: 43-60.
- Feldberg W. Synthesis of acetylcholine in sympathetic ganglia and cholinergic nerves. *J Physiol (Lond)* 1943, 101: 432-445.
- Fields RD, Le Beau JM, Longo FM, Ellisman MH. Nerve regeneration through artificial tubular implants. *Progr Neurobiol* 1989, 33: 87-134.
- Fisher M. AAEM Minimonograph #13. H reflexes and F waves: physiology and clinical indications. *Muscl Nerv* 1992, 15:1223-1233.
- Floeter MK. Spinal motor control, reflexes and locomotion. A Zigmond MJ, Bloom FE, Landis SC, Roberts JL, Squire L (eds). *Fundamental Neuroscience*. Academic Press 1999, San Diego
- Florence SL, Garrathy PE, Wall TJ, Kaas JH. Sensory afferent projections and area 3b somatotopy following median nerve cut and repair in macaque monkeys. *Cereb Cortex* 1994, 4:391-407.
- Florence SL, Kaas JH. Large-scale reorganization at multiple levels of the somatosensory pathway follows therapeutic amputation of the hand in monkeys. *J Neurosci* 1995, 15:8083-8095
- Foehring RC, Sybert GW, Munson JP. Properties of self-reinnervated motor units of medial gastrocnemius of cat. II. Axotomized motoneurons and time-course of recovery. *J Neurophysiol* 1986, 55:947-965
- Forsman J. Über die Ursachen, welche die wachstumsrichtung der peripheren Nervenfasern bei der regeneration bestimmen. *Beitr Z Pathol Anat* 1898, 24:56
- Forssberg H. Stumbling corrective reaction: A phase dependent compensatory reaction during locomotion. *J Neurophysiol* 1979, 42:936-953
- Fraher JP, O'Leary DO, Moran MA, Cole M, King RH, Thomas PK. Relative growth and maturation of axon size and myelin thickness in the tibial nerve of the rat. *Acta Neuropathol* 1990, 79:364-374.
- Friedman B, Kleinfeld D, Ip N, Verge VMK, Moulton R, Boland P, Zlotchenko E, Lindsay RM, Liu L. Neurotrophic influence on injured adult spinal motor neurons. *J Neurosci* 1995, 15: 1044-1056.
- Fritzch B, Sonntag R. Sequential double labelling with different fluorescent dyes coupled to dextran amines as a tool to estimate the accuracy of tracer application and of regeneration. *J Neurosci Methods* 1991, 39:9-17
- Frizell M, Sjöstrand J. The axonal transport of [³H] fucose labelled glycoproteins in normal and regenerating peripheral nerves. *Brain Res* 1974a, 78: 109-123.
- Fu SY, Gordon T. The cellular and molecular basis of peripheral nerve regeneration. *Mol Neurobiol* 1997, 14: 67-116.
- Fujito Y, Kawasaki H, Aoki M. Motor recovery after cross-reinnervation of a forelimb nerve by the phrenic nerve in cats. *Brain res.* 1989, 492:36-44

- Galbraith JA, Mrosko BJ, Myers RR. A system to measure thermal nociception. *J Neurosci Meth* 1993, 49: 63-68.
- Gallego R, Kuno M, Nuñez R, Snider WD. Disuse enhances synaptic efficacy in spinal motoneurons. *J Physiol (Lond)* 1979, 291:191-205
- Gallego R, Kuno M, Nuñez R, Snider WD. Enhancement of synaptic function in motoneurons during peripheral sensory regeneration. *J Physiol (Lond)* 1980, 306:205-218
- Garrathy PE, Kaas JH. Functional reorganization in adult monkeys thalamus after peripheral nerve injury. *Neuroreport* 1991, 2:274-750.
- Garrathy, P.E; Clower, R.D; La Chica, E.A; Kaas, J.H. Thalamic and cortical reorganization after chronic transection of the median and ulnar nerves in adult monkeys. *Soc Neuroscience Abstract* 1990, 16:240.
- Gerding R, Robbins N, Antosiak J, Efficiency of reinnervation of neonatal rat muscle by original and foreign nerves. *Develop Biol* 1977, 61:177-183
- Getting PA. Emerging principles governing the operation of neural networks. *Annu Rev Neurosci* 1989, 12:185-204.
- Giannini C, Lais AC, Dyck PJ. Number, size, and class of peripheral nerve fibers regenerating after crush, multiple crush, and graft. *Brain Res* 1989, 500:131-138.
- Gibby WA, Koerber HR, Horch KW. A quantitative evaluation of suture and tubulization nerve repair techniques. *J Neurosurg* 1983, 58: 574-579.
- Gibson KL, Remson L, Smith A, Satterlee N, Strain G, Daniloff J. Comparison of nerve regeneration through different types of neural prostheses. *Microsurg* 1991, 12:80-85
- Glazner GW, Ishii DN. Insulin-like growth factor gene expression in rat muscle during reinnervation. *Muscle Nerve* 1995, 18: 1433-1442.
- Gomez T, Spitzer N. In vivo regulation of axon extension and pathfinding by growth-cone calcium transients. *Nature* 1999, 397:351-355.
- Gómez N, Cuadras J, Butí M, Navarro X. Histologic assessment of sciatic nerve regeneration following resection and graft or tube repair in the mouse. *Restor Neurol Neurosci* 1996, 10:187-196.
- González M, Collins WF. Modulation of motoneurons excitability by brain derived neurotrophic factor. *J Neuropysiol* 1997, 77:502-506
- Goodrum JF and Bouldin TW. The cell biology of myelin degeneration and regeneration in the peripheral nervous system. *J Neuropathol Exp Neurol* 1996, 55: 943-953.
- Gordon T, Stein R, Thomas CK. Innervation and function of hindlimb muscles in the cat after cross-union of the tibial and peroneal nerves. *J Physiol* 1986, 374:429-441
- Gordon T, Totossy de Zepetnek, Rafuse V, Erdebil S. Motoneuronal branching and motor unit size after complete and partial nerve injuries. A Wernig A(ed): *Plasticity of motoneuronal connections*. Elsevier Science Publishers 1991
- Gordon T. Dependence of peripheral nerves on their target organs. A Burnstock G, Vrbova G, O'Brien RA (eds.). *Somatic and Autonomic Nerve-Muscle Interactions*. Elsevier 1991, New York, pp. 289-323
- Gordon T. Dependence of peripheral nerves on their target organs. A Burnstock G, O'Brien R, Vrbova G (eds.). *Somatic and Autonomic Nerve-Muscle Interactions*, New York, Elsevier Science Publisher, 1983, pp 289-325.
- Gorio A, Carmignoto G, Finesso M, Polato P, Nunzi MG. Muscle Reinnervation- II. Sprouting, Synapse formation and repression. *Neurosci Vol.* 1983, 8(3): 403-416
- Gozariu M, Bragard D, Willer JC, Le Bars D. Temporal summation of C-fiber afferent inputs: competition between facilitatory and inhibitory effects on C-fiber reflex in the rat. *J Neurophysiol* 1997, 78:3165-3179
- Graeber MB, Kreutzberg GW. Astrocytes increase in glial fibrillary acidic protein during retrograde changes of facial motor neurons. *J neurocytol* 1986, 15:363-373
- Graeber MB, Kreutzberg GW. Delayed astrocyte reaction following facial nerve axotomy. *J. Neurocytol* 1988, 17:209-220

- Graeber MB, Tetzlaff W, Streit WJ, Kreutzberg GW. Microglial cells but not astrocytes undergo mitosis following rat facial nerve axotomy. *Neurosci Lett* 1988, 85:317-321
- Graham-Brown T, Sherrington CS. The rule of reflex response in the limbs reflexes of the mammal and its exceptions. *J Physiol* 1912, 44:125-130
- Grant MP, Landis SC, Siegel RE. The molecular and pharmacological properties of muscarinic cholinergic receptors expressed by rat sweat gland are unaltered by denervation. *J Neurosci* 1991, 11:3763-3771.
- Grant MP, Landis SC. Developmental expression of muscarinic cholinergic receptors and coupling to phospholipase C in rat sweat glands are independent of innervation. *J Neurosci* 1991, 11:3772-3782
- Greenberg SG, Lasek RJ. Neurofilament protein synthesis in DRG neurons decreases more after peripheral axotomy than after central axotomy. *J Neurosci* 1988, 8:1739-1746
- Greene EC. *Anatomy of the rat*. Hafner Publishing Co 1955, New York.
- Gregoric M. Suppression of flexor reflex by transcutaneous electrical nerve stimulation in spinal cord injured patients. *Musc Nerv* 1998, 21:166-172
- Groves MJ, Christopherson T, Giometto B, Scaravilli F. Axotomy-induced apoptosis in adult rat primary sensory neurons. *J Neurocyt* 1997, 26:615-624.
- Gruart A, Gunkel A, Neiss WF, Angelov DN, Stennert E, Delgado-Garcia JM. Changes in eye blink responses following hypoglossal-facial anastomosis in the cat: evidence of unadaptability to new motor tasks. *Neurosci* 1996, 73 (1): 233-247
- Gundersen HJ. Stereology of arbitrary particles. A review of unbiased number and size estimators and the presentation of some new ones in memory of William R Thomson. *J Microsc* 1986, 143: 3-45.
- Gunthinas-Lichius O, Neiss WF, Gunkel A, Stennert E. Differences in glial synaptic and motoneuron responses in the facial nucleus of the rat brainstem following facial nerve resection and nerve suture reanastomosis. *Eur Arch Oto-Rhino-Laryngol* 1994, 251:410-417
- Guntinas-Lichius, streppel M, Angeov DN, Stennert E, Neiss W. Effect of delayed facial-facial nerve suture on facial nerve regeneration. *Acta Otorinol* (Stockh) 1997, 1176:670-674
- Gustafsson B. Changes in motoneurons electrical properties following axotomy. *J Pysiol (Lond)* 1979, 293:197-215
- Guth L. Neuromuscular function after regeneration of interrupted nerve fibers into partially denervated muscle. *Exp Neurol* 1962, 6:129-141
- Guth L. regeneration in the mammalian peripheral nervous system. *Physiol. Rev.* 1980, 5:441-478.
- Gutmann E, Guttmann L, Medawar PB, Young JZ. The rate of regeneration of nerve. *J Exp Biol* 1942, 19: 14-44.
- Gutmann E, Guttmann L. Factors affecting recovery of sensory function after nerve lesions. *J Neurol Psychiat* 1942, 5:117-129.
- Gutmann, E. Factors affecting recovery of motor function after nerve lesions. *J Neurol Psychiat* 1942, 5:81-95.
- Haas CA, Donath C, Kreutzberg GW. Differential expression of immediate early genes after transection of the facial nerve. *Neuroscience* 1993, 53:91-99
- Haas CA, Streit WJ, Kreutzberg GW. Rat facial motoneurons express increased levels of calcitonin gene related peptide mRNA in response to axotomy. *J. Neurosci Res* 1990, 27:270-275
- Hare GM, Evans PJ, Mackinnon SE, Best TJ, Bain JR, Szalai JP, Hunter RT. Walking track analysis: a long term assessment of peripheral nerve recovery. *Plast Reconstruct Surg* 1992, 89(2):251-258.
- Hargreaves K, Dubner R, Brown F, Joris J. A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia. *Pain* 1988:77-88.
- Härkönen MHA, Kauffman FC. Metabolic alterations in the axotomized superior cervical ganglion of the rat. II. The pentose phosphate pathway. *Brain Res* 1974, 65: 141-157.
- Havton L, Kellerth JO. Elimination of intramedullary axons collaterals of cat spinal a-motoneurons following peripheral nerve injury. *Exp Brain Res* 1990. 79:65-74

- Hellgren J, Kellert JO. A physiological study of the monosynaptic reflex responses of cat spinal a-motoneurons after partial lumbosacral deafferentation. *Brain Res* 1989, 488:149-162
- Hendry LA, Hill CE, Watters DJ. Long-term retention of Fast Blue in symathetic neurones after axotomy and regeneration-deonstration of incorrect reconnections. *Brain Res* 1986, 376:292-298
- Henneman E, Olson CB. Relations between structure and function in the design of skeletal muscles. *J Neurophysiol* 1965, 28:581-598
- Henning R, Dietrichs E. Transient reinnervation of antagonistic muscles by the same neuron. *Exp Neurol* 1994, 130:331-336
- Henry EW, Chiu T-H, Nyilas E, Brushart TM, Dikkes P, Sidman RL. Nerve regeneration through biodegradable polyester tubes. *Exp Neurol* 1985, 90: 652-676
- Hermanson M, Olsson T, westmark B, Funa K. PDGF and its receptor following facial nerve axotomy in rats: expression in neurons and sorroundings glia. *Exp Brain Res* 1995, 102:415-422
- Himes BT, Tessler A. Death of some dorsal root ganglion neurons and plasticity of others following sciatic nerve section in adult and neonate rats. *J Comp Neurol* 1989, 284:215-213
- Hirakawa M, Kawata M. Changes of chemoarchitectural organization of the rat spinal cord following ventral and dorsal root transection. *J Comp Neurol* 1992, 320:339-352
- Hoffman P. Über die beziehung der Shenreflexe zur willkurlichen Bewegun zum Tonus. *Z Biol* 1918, 68:351-370.
- Hoffman P. Untersuchungen über die Eigenreflexe (Shenreflexe) Menschlicher Muskeln. Springer Verlag 1922, Berlin.
- Hoffman PL, Lasek RJ. Axonal transport of the cytoskeleton in regenerating motor neurons: constancy and change. *Brain Res* 1980, 202: 317-333.
- Holmberg H, Schouenborg J. Developmental adaptation of withdrawal reflexes to early alteration of peripheral innervation in the rat. *J Physiol* 1996a, 495(2):399-409
- Holmberg H, Schouenborg J. Postnatal development of the nociceptive withdrawal reflexes in the rat: a behavioural and electromyographic study. *J Physiol* 1996b, 493(2):239-252
- Horch K. The influence of mechanoreceptors structures in regenerating sensory axons after cutaneous nerve transaction in the cat. *Neurosci Lett* 1982, 32:281-284
- Horch KW, Lisney JW. Changes in primary afferent depolarisation of sensory neurones during peripheral nerve regeneration in the cat. *J Pysiol (Lond)* 1981, 313:287-299
- Hörkfelt T, Zhang X, Wiesenfeld-Hallin Z. Messenger plasticity in primary sensory neurons following axotomy and its functional implication. *Trends Neurosci* 1994, 17:22-30
- Hruska RE, Kennedy S, Silbergeld EK. Quantitative aspects of normal locomotion in rats. *Life Sci* 1979, 25:171-180
- Hsieh ST, Choi S, Lin WM, Chang YC, McArthur J, Griffin J. Epidermal denervation and its effects on keratinocytes and Langerhans cells. *J Neurocyt* 1996, 25:513-524
- Huerta M, Wall J. The functional territory of the primate median nerve in the hand skin and the cortical map. *Soc Neurosci Abstr.* 1987, 13:250
- Hultborn H, Meunier S, Pierrot-deseilligny E, Shindo M. Changes in the presynaptic inhibition of Ia fibers at the onset of voluntary contraction in man. *J physiol (London)* 1987, 389:757-772
- Hultborn H, Lipski J, Mackel R, Wigstrom H. Distribution of recurrent inhibition within a motor nucleus. I. Contribution from slow and fast units to the excitation of Renshaw cells. *Acta Physiol Scand* 1988, 134:347-361
- Imai H, Tajima T, Natsumi Y. Successful reeducation of functional sensibility after median nerve repair at the wrist. *J hand Surg* 1991, 16A:157-164
- Ishikawa K, Tanaka M, Black JA, Waxman SG. Changes in expression of voltage-gated potassium channels in dorsal root ganglion neurons following axotomy. *Musc and Nerve* 1999, 22:502-507
- Jahromi BS, Robitaille R, Charlton MP. Transmitter release increases intracellular calcium in perisynaptic Schwann cells in situ. *Neuron* 1992, 8: 1069-1077.
- Jain N, Florence SL, Kaas JH. Reorganization of somatosensory cortex after nerve and spinal cord injury. *New Physiol Sci* 1998, 13:143-149

- Jenkins WM, Merzenich NM, Ochs MT, Allard T, Guic Robles E. Functional reorganization of primary somatosensory cortex in adult owl monkeys after behaviourally controlled tactile stimulation. *J Neurophysiol* 1990a, 63:82-104
- Jenkins WM, Merzenich NM, Rencanzone G. Neocortical representational dynamics in adult primates: Implications for neurophysiology. *Neuropsychologia* 1990b, 28:573-584.
- Jenq CB, Coggeshall RE. Effects of sciatic nerve regeneration on axonal populations in tributary nerves. *Brain Res* 1984, 295:91-100
- Jenq CB, Coggeshall RE. Long-term patterns of axon regeneration in the sciatic nerve and its tributaries. *Brain Res* 1985, 345:34-44
- Jenq CB, Coggeshall RE. Regeneration of transected rat sciatic nerves after using isolated nerve fragments as distal inserts in silicone tubes. *Exp Neurol* 1986, 91:154-162
- Johnson EM, Taniuchi M, DiStefano PS. Expression and possible function of nerve growth factor receptors on Schwann cells. *Trends Neurosci* 1988, 11: 299-304.
- Karlstrom L, Dahlström A. The effect of different types of axonal trauma on the synthesis and transport of amine storage granules in rat sciatic nerves. *J Neurobiol* 1973, 4: 191-200.
- Kato S, Ide C. Axonal sprouting at the node of Ranvier of the peripheral nerve disconnected with the cell body. *Restor Neurol Neurosci* 1994, 6: 181-187
- Kelly RB, Grote E. Protein target in the neuron. *Annu Rev Neurosci* 1993, 16: 95-127.
- Kennedy WR, Navarro X, Kamei H. Reinnervation of sweat glands in the mouse: axonal regeneration versus collateral sprouting. *Musc Nerv* 1988, 11:603-609.
- Kennedy WR, Sakuta M, Quick DC. Rodent eccrine sweat glands. A case of multiple efferent innervation. *Neuroscience* 1984, 11: 741-749.
- Kennedy WR, Sakuta M. Collateral Reinnervation of sweat glands. *An Neurol* 1984, 15:73-78.
- Kim D, Connolly SE, Gillespie JT, Voorhies RM, Kline D. Electrophysiological studies of various graft length and lesion in repair of nerve gaps in primates. *J Neurosurg* 1991, 75:440-446
- Kimura J, Rodnitzky L, Owara SH. Electrophysiologic analysis of aberrant regeneration after facial nerve paralysis. *Neurology* 1975, 25:989-993.
- King AE, Thomson WN. Brief and prolonged changes in spinal excitability following peripheral injury. *Seminars Neurosci* 1995, 7:233-243
- Klapdor K, Dulfer B, Hammann A, Van der Staay FJ. A low-cost method to analyse footprint patterns. *J Neurosci Meth* 1997, 75:49-54
- Kline DG, Hudson AR. Selected recent advances in peripheral nerve injury research. *Surg Neurol* 1985, 24: 371-376.
- Kline DG. Surgical repair of peripheral nerve injury. *Muscle Nerve* 1990, 13:843-852.
- Koerber HR, Mirnics K, Brown PB, Mendell LM. Central sprouting and functional plasticity of regenerated primary afferents. *J Neurosci* 1994, 14(6):3655-3671
- Koerber HR, Mirnics K, Mendell LM. Properties of regenerated primary afferents and their functional connections. *J Neurophysiol* 1995, 73(2):693-702.
- Koerber HR, Mirnics K. Plasticity of dorsal horn cell receptive fields after peripheral nerve regeneration. *J Neurophysiol* 1996, 75(6):2255-2267
- Korshing S. The neurotrophic factor concept: a reexamination. *J Neurosci* 1993, 13: 2739-2748
- Korthals JK, Gieron MA, Wisniewski M. Nerve regeneration patterns after acute ischemic injury. *Neurology* 1989, 39: 932-937.
- Kreutzberg GW. reaction of the neuronal cell body to axonal damage. *The Axon: Structure, Function and Pathophysiology* (Waxman SG, Kocsis JD, Stys PK eds.) Oxford University Press, New York, Oxford. 1995, pp. 355-374
- Kuecherer-Ehret A, Graeber MB, Edgard D, Thoenen H. Immunoelectron microscopic localization of laminin in normal and regenerating mouse sciatic nerve. *J Neurocytol* 1990, 19: 101-109.
- Kuffler DP. Regeneration of muscle axons in the frog is directed by diffusible factors from denervated muscle and nerve tubes. *J Exp Neurol* 1989. 281:416
- Kuiken TA, Childress DS, Rymer WZ. The hyper-reinnervation of rat skeletal muscle. *Brain Res* 1995, 676:113-123

- Kuno M, Llinás R. Alterations of synaptic action in chromatolysed motoneurons of the cat. *J Physiol (Lond)* 1970, 210:823-838
- Kuno M, Miyata Y, Martínez-Muñoz EJ. Differential reaction of fast and slow α -motoneurons to axotomy. *J Physiol (Lond)* 1974, 240:725-739
- Labrador RO, Butí M, Navarro X. Influence of Collagen and Laminin gels concentration on nerve regeneration after resection and tube repair. *Exp Neurol* 1998, 149:243-252
- Labrador RO, Butí M, Navarro X. Peripheral nerve repair: role of agarose matrix density on functional recovery. *NeuroReport* 1995, 6:2022-2026
- Laird JM, Bennett GJ. An electrophysiological study of dorsal horn neurons in the spinal cord of rats with an experimental neuropathy. *J Neurophysiol* 1993, 69:2072-2085
- Laird JM, Hargreaves RG, Hill RG. Effect of RP 67580, a non-peptide neurokinin1 receptor antagonist, on facilitation of a nociceptive spinal flexion reflex in the rat. *Br J Pharmacol* 1993, 109:713-718.
- Lasek RJ. Protein transport in neurons. *Int Rev Neurobiol* 1970, 13: 289-324.
- Laskowski MB, Sanes J. Topographically selective reinnervation of adult mammalian skeletal muscles. *J Neurosci* 1988, 8(8):3094-3099.
- Lázaro JJ, Franquelo C, Navarro X, Castellano B, Verdú E, Cristòfol C, Arboix M. Prolongation of nerve and Epidural Anesthetic Blockade by bupivacaine in a lipid emulsion. *Anesth Anal* 1999, 89:121-127.
- Le Beau JM, Ellisman MH, Powell HC. Ultrastructural and morphometric analysis of long-term peripheral nerve regeneration through silicone tubes. *J Neurocytol* 1988, 17: 161-172.
- LeBlanc AC, Poduslo JF. Axonal modulation of myelin gene expression in the peripheral nerve. *J Neurosci Res* 1990, 26: 317-326.
- LeBlanc AC, Windebank AJ, Poduslo JF. P0 gene expression in Schwann cells is modulated by increase of cAMP which is dependent on the presence of axons. *Mol Brain Res* 1992, 12: 31-38.
- Lekan H, Carlton SM, Coggeshall. Sprouting of A β fibers into lamina II of the rat horn in peripheral neuropathy. *Neurosci Lett* 1996, 208:147-150
- Letinsky MS, Fischbeck KH, McMahan UJ. Precision of reinnervation of original postsynaptic sites in frog muscle after nerve crush. *J Neurocytol* 1976, 5:691-718
- Letourneau P, Shattuck TA. Distribution and possible interactions of actin-associated proteins and cell adhesion molecules of nerve growth cones. *Development* 1989, 195: 505-519.
- Levin M, Feldman AG. The role of stretch threshold regulation in normal and impaired motor control. *Brain Res* 1994. 657:23-30
- Lewin GR, McMahon S. Physiological properties of primary sensory neurons appropriately and inappropriately innervating skeletal muscle in adult rats. *J Neurophysiol* 1991a, 66(4):1218-1230
- Lewin GR, McMahon S. Physiological properties of primary sensory neurons appropriately and inappropriately innervating skin in the adult rat. *J Neurophysiol* 1991b, 66(4):1205-1217
- Li Y, Hsieh ST, Chien HF, Zhang X, McArthur J, Griffin J. Sensory and Motor denervation influence Epidermal Thickness in rat Foot Glabrous Skin. *Exp Neurol* 1997, 147:452-462
- Liddell EGT, Sherrington CS. Reflexes in response to stretch (myotatic reflexes). *Proc R Soc London, Ser. B* 1924, 96:212-242
- Lieberman AR. The axon reaction: a review of the principal features of perikaryal responses to axon injury. *Int Rev Neurobiol* 1971, 14: 49-124.
- Liu CN, Amir R, Devor M. Effect of age and nerve injury on cross-excitation among sensory neurons in rat dorsal root ganglia. *Neurosci Lett* 1999, 259:95-98
- Liu HM. Growth factors and extracellular matrix in peripheral nerve regeneration, studied with a nerve chamber. *J Periph Nerv Sys* 1996, 1: 97-110.
- Lloyd DPC. The interaction of antidromic and orthodromic volleys in a segmental spinal motor nucleus. *J Neurophysiol* 1943, 6:143-151.
- Lo A. C, Hoienu LJ, Oppenheimer RW. Apoptosis in the nervous system: morphological features, methods, pathology and prevention. *Arch. Histol. Cytol.* 1995, 58:139-149

- Longo FM, Hayman EG, Davis GE, Ruoslathi E, Engvall E, Manthorpe M, Varon S. Neurite-promoting factors and extracellular matrix components accumulating in vivo within nerve regeneration chambers. *Brain Res* 1984, 309: 105-117.
- Longo FM, Manthorpe M, Skaper SD, Lundborg G, Varon S. Neuronotrophic activities accumulate in vivo within silicone nerve regeneration chambers. *Brain Res* 1983, 261:109-117.
- Lubinska L. Early course of Wallerian degeneration in myelinated fibres of the rat phrenic nerve. *Brain Res*. 1977, 130: 47-63.
- Lundberg A, Malmgren K, Schomburg ED. Reflex pathways from group II muscle afferents. 3. Secondary spindle afferents and the FRA: A new hypothesis. *Exp Brain res* 1987, 65:294-306
- Lundberg A. Multisensory control of spinal reflex pathways. *Progr Brain Res* 1979, 50:11-28
- Lundborg G, Dahlin LB, Danielsen N, Geberman RH, Longo FM, Powell HC, Varon S. Nerve regeneration in silicone chambers: influence of gap length and of distal stumps components. *Exp Neurol* 1982b, 76:361-375
- Lundborg G, Dahlin LB, Danielsen N, Johannesson A, Hansson HA. A: Lee AJC, Albrektsson T, Branemark PI (eds). *Clinical Applications of Biomaterials*. Chichester, UK, John Wiley Sons Ltd. 1982a, p. 323
- Lundborg G, Dahlin LB, Danielsen N, Nachmson AK. Tissue specificity in nerve regeneration. *Scand J Plast Reconstruct Surg* 1986, 20:279
- Lundborg G, Longo FM, Varon S. Nerve regeneration model and trophic factors in vivo. *Brain Res* 1982c, 232: 157-161.
- Lundborg G, Rosen B, Abrahamsson SO, Dahlin LB, Danielsen N. Tubular repair of the median nerve in the human forearm. Preliminary findings. *J Hand Surg* 1994. 19B:273-276
- Lundborg G, Rosen B, Dahlin L, Danielsen N, Holmberg J. Tubular versus conventional repair of median nerve and ulnar nerves in the human forearm: early results from a prospective, randomized, clinical study. *J Hand Surg* 1997, 22: 99-106.
- Lundborg G. *Nerve Injury and Repair*. Churchill Livingstone, Edinburgh, 1988.
- Lundborg G. Nerve regeneration and repair. *Acta Orthop Scand* 1987, 58: 145-169.
- Lundborg G. Peripheral nerve injuries. Pathophysiology and strategies for treatment. *J Hand Therapy* 1993, 6:179-189
- Lunn ER, Brown MC, Perry VH. The pattern of axonal degeneration in the peripheral nervous system varies with different types of lesion. *Neuroscience* 1990, 35: 157-165.
- Luo ZJ, Lu SB. Selective reinnervation of regenerating mixed nerve fibers across a silicone tube gap. *J Hand Surg* 1996, 21B:660-663
- Mackinnon S, Dellon AL, Lundborg G, Hudsins AR, Hunter DA. A study of neurotrophism in a primate model. *J Hand Surg* 1985a, 11A:888
- Mackinnon S, Dellon AL. Reinnervation of distal sensory nerve environments by regenerating sensory axons. *Neurosci* 1992, 46: 595-603
- Mackinnon S, Hudson AR, Hunter DA. Histologic assessment of nerve regeneration in the rat. *Plast Reconstr Surg* 1985b, 75:384-388
- Mackinnon SE, Dellon AL, Hunter DA. Histological assessment of the effects of the distal nerve in determining regeneration across a nerve graft. *Microsurg* 1988, 9:46-51.
- Mackinnon SE, Dellon AL, O'Brien JP. Changes in the nerve fiber numbers distal to a nerve repair in the rat sciatic nerve model. *Musc Nerv* 1991, 14:1116-1122
- Mackinnon SE, Dellon AL. A study of nerve regeneration across synthetic (Maxon) and biologic (Collagen) nerve conduits for nerve gaps up to 5 cm in the primate. *J Reconstruct Microsurg* 1990, 6: 117-121.
- MacMahon S, Gibson S. Peptide expression is altered when afferent nerves reinnervate inappropriate tissues. *Neurosci Lett* 1987, 73:9-15
- Madison RD, Archibald SJ, Brushart TME. Reinnervation accuracy of the rat femoral nerve by motor and sensory neurons. *J Neurosci* 1996, 16: 5698-5703.

- Madison RD, Da Silva CF, Dikkes P, Sidman RL, Chiu T-H. Peripheral nerve regeneration with entubulation repair: Comparison of biodegradable nerve guides versus polyethylene tubes and the effects of a laminin-containing gel. *Exp Neurol* 1987, 95:378-390.
- Marinesco G. Sur les phenomenes de reparation dans les centres nerveux apres la section des nerfs periferiques. *Compt Rendu Soc Biol* 1896, 111:930
- Mathews MR, Nelson VH. Detachment of structurally intact nerve endings from chromatolytic neurones of rat superior cervical ganglion during the depression of synaptic transmission induced by post-ganglionic axotomy. *J Physiol (Lond)* 1975, 245: 91-135.
- Matre AD, Sinkjaer T, Svensson P, Arendt-Nielsen L. Experimental muscle pain increass the human stretch reflex. *Pain* 1998, 75:331-339
- Matzner O, Marshall D. Hyperexcitability at sites of nerve injury depends on voltage-sensitive Na⁺ channels. *J Neuropysiol* 1994, 72(1):349-359
- Mayer R, Denny-Brown D. Conduction velocity in peripheral nerve during experimental demyelination in the cat. *Neurology* 1964, 14:714726.
- McAllister A, Katz LC, Lo D. Neurotrophins ans synaptic plasticity. *Annu Rev Neurosci* 1999, 22:295-318
- McCrea D, Pratt CA, Jordon LM. rensshaw cell activity and recurrent effects on motor neurons during fictive locomotion. *J Neurophysiol* 1980, 44:475-488
- McGrath GJ. Matthews PBC. Evidence from the use of procaine nerve block that the spindle group II fibres contribute to the excitation in the tonic stretch reflex of the decerebrate cat. *J Physiol (London)* 1973, 235:371-408
- McMahon S, Gibson S. Peptide expression is altered when afferent nerves reinnervate inappropriate tissue. *Neurosci Lett* 1987, 73:9-15
- McMahon S, Lewin GR, Wall P. Central hyperexcitability trigered by noxious inputs. *Curr Op Biol* 1993, 3:602-610.
- Melville S, R Coggeshall. Sensory cell death caused by sciatic nerve transection prevented by a silicone tube. *Anat Rec* 1989.
- Melville S, Sherburn TE, Coggeshall RE. Preservation of sensory cells by placing stumps of transected nerve in an impermeable tube. *Exp Neurl* 1989, 105: 311-315.
- Melzer P, Zhang MZ, Mckanna JA. Infraorbital nerve transection and whisker follicle removal in adult rats affect microglia and astrocytes in trigeminal brainsteam. A study with lipocortin 1 and S100 immunohistochemistry. *Neurosci* 1997, 80:459-472
- Mendell L. Modifiability of spinal synapses. *Physiol Rev* 1984. 64: 260-324
- Mendell LM, Johnson R, Munson J. Neurotrophin modulation of the monosynaptic reflex after peripheral nerve transection. *J Neurosci* 1999, 19(8):3162-3170
- Mendell LM, Munson JB, Scott JG. Alterations of synapses on axotomized motoneurones. *J Physiol (Lond)* 1976, 255: 67-79.
- Mendell LM, Scott JG. The effects of peripheral nerve cross-union on connections of single Ia fibers to motoneurones. *Exp Brain res* 1975, 22:221-234
- Mendell LM, Wall PD. Responses of single dorsal cord cells to peripehral cutaneous unmyelinated fibres. *nature* 1965, 206:97-99
- Mendell LM. Physiological properties of unmyelinated fibre projections to the spina cord. *Exp Neurol* 1966, 16:316-332
- Merle M, Dellon L, Campbell J, Chang PS. Complications from silicon-polymer intubulation of nerves. *Microsurg* 1989, 10:130-133
- Merzenich MM, Kaas JH, Wall JT, Sur M, Nelson RJ, Felleman DJ. Progression of change following median nerve section in the cortical representation of the hand in areas 3b and 1 of adult owl and squirrel monkeys. *Neuroscience* 1983, 10:639-665.
- Mews M, Meyer M. Modulation of Schwann cell phenotype by TGF-b-1: Inhibition of P₀ mRNA expression and downregulation of the low affinity NGF receptor. *Glia* 1993, 8:208-217

- Meyer S, Abrams R, Botte J, Davey J, Bodine-Fowler S. Functional recovery following neurotomy of the sciatic nerve by epineurial repair compared with tubulization. *J Orthop Res* 1997, 5:664-669
- Meyerson BA, Ren B, Herregodts P, Linderöth B. Spinal cord stimulation in animal models of mononeuropathy: effects on the withdrawal response and the flexor reflex. *Pain* 1995, 61:229-243
- Millesi H, Terzis JK. Nomenclature in peripheral nerve surgery. *Clin Plast Surg* 1984, 11: 3-8.
- Millesi H. Reappraisal of nerve repair. *Surg Clin North Am* 1981, 61: 321-340.
- Milner-Brown HS, Stein RB, Lee RG. The pattern of recruiting human motor units in neuropathies and motor neurone disease. *J Neurol Neurosurg Psychiatry* 1974, 37:665-669
- Mira JC, Fardeau M. Nerve and muscle changes induced by repeated localized freezings of the sciatic nerve in the rat. In: *Peripheral Neuropathies*. N. Canal and G. Pozza eds. Elsevier/North-Holland Biomedical Press. 1978: 83-90.
- Mira JC. Étude Quantitatives sur la régénération des fibres nerveuses myélinisées. II. variation du nombre et du calibre des fibres régénérées après un écrasement ou une section du nerf. *Arch Anat Microscopique* 1976, 65:255-284
- Mogilner A, Grossman AI, Ribary U, Joliot M, Volkmann J, Rapaport D, Beasley RW, Llinás RR. Somatosensory cortical plasticity in adult humans revealed by magnetoencephalography. *Proc Natl Acad Sci USA* 1993, 90:3593-3597.
- Molander H, Engkvist O, Hägglund J, Olsson Y, Torebjök E. Nerve repair using a polyglactin tube and nerve graft: an experimental study in the rabbit. *Biomaterials* 1983, 4: 276-280.
- Molecular Probes. Lipophilic Tracers DiI, DiO, DiD and DiA. Product Information Sheet. MP. 0282, 1/1/1997
- Moneta ME, Gehrman J, Topper RB, Banati RB, Kreutzberg GW. Cell adhesion molecule expression in the regenerating rat facial nucleus. *J Neuroimmunol* 1993, 45:203-206
- Montserrat L, Benito M. Facial Synkinesis and aberrant regeneration of facial nerve. *Adv. Neurol* 1988, 49:211-224.
- Moreno-López B, De la Cruz R, Pastor AM, Delgado-García JM. Effects of botulinum neurotoxin type A on abducens motoneurons in the cat: alterations of the discharge pattern. *Neurosci* 1997, 81(2): 437-455
- Müller HW, Williams LR, Varon S. Nerve regeneration chamber: evaluation of exogenous agents applied by multiple injections. *Brain Res* 1987, 413:220-326
- Munger BL. The reinnervation of denervated skin. A Hamann W and Iggo A (eds.). *Progr Brain Res* 1988, 74:259-262
- Munro C, Szalai J, Mackinnon S, Midha R. Lack of association between outcome measures of nerve regeneration. *Musc Nerv* 1998, 21:1095-1097
- Munson JB, Johnson RD, Mendell LM. NT-3 increases amplitude of EPSP produced by axotomized group Ia afferents. *J Neurophysiol* 77:2209-2212
- Murray M, Goldberger ME. Restitution of functional and collateral sprouting on the cat spinal cord: the partially hemisectioned animal. *J Comp Neurol* 1974, 158:19-36
- Nakanishi T, Norris FH. Motor nerve fibers in rat sural nerve. *Exp Neurol* 1970, 26:433-435
- Navarrete R, Shahani U, Vrbová G. Long-Lasting modification of reflexes after neonatal nerve injury in the rat. *J Neurol Sci* 1990, 96:257-265
- Navarro X, Kennedy W. Effect of age on collateral reinnervation of sweat glands in the mouse. *Brain Res* 1988, 463:174-181.
- Navarro X, Kennedy W. The effects of autologous nerve transplants on motor and sudomotor reinnervation by regenerative axons. *Brain Res* 1991, 565:181-187
- Navarro X, Kennedy WR, Fries TJ. Small nerve fiber dysfunction in diabetic neuropathy. *Muscle Nerve* 1989, 12: 498-507.
- Navarro X, Kennedy WR. Changes in sudomotor nerve territories with aging in the mouse. *J Auton Nerv Syst* 1990a, 31: 101-108.
- Navarro X, Kennedy WR. The effect of a conditioning lesion on sudomotor axon regeneration. *Brain Res* 1990b, 509:232-236.

- Navarro X, Rodríguez FJ, Labrador RO, Butí M, Ceballos D, Gómez N, Cuadras J, Perego G. Peripheral nerve regeneration through bioresorbable and durable nerve guides. *J Periph Nerv Syst* 1996, 1: 53-64.
- Navarro X, Valero A, Gudiño G, Forés J, Rodríguez FJ, Verdú E, Pascual R, Cuadras J, Nieto-Sampedro M. Ensheathing glia transplants promote dorsal root regeneration and spinal reflex restitution after multiple lumbar rhizotomy. *Ann Neurol* 1999, 45(2):207-215
- Navarro X, Verdú E, Butí M. Autotomy prevention by amitriptyline after nerve section in different strains of mice. *Restor Neurol Neurosci* 1994a, 6:151-157
- Navarro X, Verdú E, Butí M. Comparison of regenerative and reinnervating capabilities of different functional types of nerve fibers. *Exp Neurol* 1994b, 129:217-224 .
- Navarro X, Verdú E, Wendelschafer-Crabb G, Kennedy WR. Innervation of cutaneous structures in the mouse hindpaw: A confocal microscopy immunocytochemical study. *J Neurosci Res* 1995, 41:111-120
- Navarro, X, Rodríguez FJ, Labrador RO, Butí M, Ceballos D, Gómez N, Cuadras J, Perego G. Peripheral nerve regeneration through bioresorbable and durable nerve guides. *J Periph Nerv Syst* 1996, 1:53-64
- Nguyen Q, Parsadanian AS, Snider W, Lichtman JW. Hyperinnervation of neuromuscular junctions caused by GDNF overexpression in muscle. *Science* 1998, 279:1725-1729
- Nieto A. Molecular biology of axon guidance. *Neuron* 1996, 17:1039-1048
- Nieto-Sampedro M. Plasticidad neural: una propiedad básica que subyace desde el aprendizaje a la reparación de lesiones. A: Mora F (ed.). *El cerebro íntimo*. Barcelona 1997, Ariel pp.66-96
- Nishimura H, Johnson RD, Munson JB. Rescue of motoneurons from the axotomized state by regeneration into a sensory nerve in cats. *J Neurophysiol* 1991, 66:1462-1470.
- Nothias F, Tessler A, Murray M. Restoration of substance p and calcitonin gene-related peptide in dorsal root ganglia and dorsal horn after neonatal sciatic nerve lesion. *J Comp Neurol* 1993, 334:370-384
- Novikova L, Novikov L, Kellerth JO. Persistent neuronal labeling by retrograde fluorescent tracers a comparison between Fast Blue, Fluoro-Gold and various dextran conjugates. *J Neurosci Meth* 1997, 74:9-15
- Nurse CA, Macintyre L, Diamond J. Reinnervation of the rat touch dome restores the merkel cell population reduced after denervation. *Neurosci* 1984, 13 (2): 563-571
- O'Brien RAD. Axonal transport of acetylcholine, choline acetyltransferase and cholinesterase in regenerating peripheral nerve. *J Physiol (Lond)* 1978, 282: 91-103.
- Ochi M. Experiments on orientation of regenerating fibers in the severed peripheral nerve. *Hiroshima J Med Sci* 1983, 31:386-406
- Ochs G, Schenk M, Struppler A. Painful dysaesthesias following peripheral nerve injury; a clinical and electrophysiological study. *Brain Res* 1989, 496: 228-240.
- Önne L. Recovery of sensibility and sudomotor activity in the hand after suture. *Acta Chirurg Scand* 1962, Suppl. 300:1-69.
- Ossowska K, Lorenc-Koci E, Schulze G, Wolfarth S. The influence of dizocilpine (MK-801) on the reserpine-enhanced electromyographic stretch reflex in rats. *Neurosci Lett* 1996, 203:73-76
- Ostberg AJ, Vrbová G, O'Brien AD. Reinnervation of fast and slow mammalian muscles by a superfluous number of motor axons. *Neuroscience* 1986, 18:205-213
- Papaloizos MY, Holmquist B, Lundborg G. An experimental study of nerve grafting combined with silicone tubes in the rat model: functional outcome and specificity of muscle reinnervation. *Restor Neurol Neurosci* 1997, 11:161-168.
- Pasterkamp RJ, Giger R, Verhaagen J. Regulation of semaphorin III/collapsin-1 gene expression during peripheral nerve regeneration. *Exp Neurol* 1998, 153:313-327.
- Pastor AM, Moreno-López B, de la Cruz R, Delgado-García JM. Effects of botulinum neurotoxin type A on abducens motoneurons in the cat: Ultrastructural and synaptic alterations. *Neurosci* 1997, 81(2): 457-478
- Pearson JC, Finkel LH, Edelman GM. Plasticity in the organization of adult cerebral cortical maps: a computer simulation based on neuronal group selection. *J Neurosci* 1987, 7:4209-4223.

- Perry VH, Brown MC. Macrophages and nerve regeneration. *Curr Op Neurobiol* 1992a, 2: 679-682.
- Persson JKE, Aldskogius H, Arvidsson J, Holmberg A. Ultrastructural changes in the gracile nucleus of the rat after sciatic nerve transection. *Anat Embryol* 1991, 184:591-604
- Persson JKE, Svensson M, Aldskogius H. Cell proliferation in the gracile nuclei and dorsal spinal cord following sciatic nerve transection in the adult rat. *Primary Sensory Neuron* 1995, 1:47-64.
- Peyronnard JM, Charron L, Lavoie J, Messier JP, Bergouignan FX. Comparative study of the effects of chronic axotomy, crush lesion and re-anastomosis of the rat sural nerve on horseradish peroxidase labelling of primary sensory neurons. *Brain Res* 1988, 443:295-309.
- Peyronnard JM, Charron L. Motor and sensory neurons of the rat sural nerve: A horseradish peroxidase study. *Musc Nerv* 1982, 5:654-660
- Pi R, Navarro X, Valero A. Lesions traumàtiques del sistema nerviós perifèric: de la clínica al laboratori experimental. A: Valero A (ed). *Neurociències clíniques: una visió interdisciplinària de les malalties del sistema nerviós*. Monografies de l'AENC-UAB. Universitat Autònoma de Barcelona 1999, Bellaterra
- Picard O, Boucher S, Regoli D, Gitter B, Howbert J, Couture R. Use of non peptide tachykinin receptor antagonist to substantiate the involvement of NK-1 and NK-2 receptors in a spinal cord nociceptive reflex in the rat. *Eur J Pharmacol* 1993, 232:255-261.
- Pinco M, Lev-Tov. Modulation of monosynaptic excitation in the neonatal rat spinal cord. *J Neurophysiol* 1993, 70:1151-1158
- Politis MJ, Ederle K, Spencer PS. Tropism in nerve regeneration in vivo. Attraction of regenerating axons by diffusible factors derived from cells in distal nerve stumps of transected peripheral nerves. *Brain Res* 1982, 253:1-12.
- Politis MJ, Spencer PS. An in vivo assay of neurotropic activity. *Brain Res* 1983, 278:229-231
- Politis MJ, Steiss JE. Electromyographic evaluation of a novel surgical preparation to enhance nerve-muscle specificity that follows mammalian peripheral nerve trunk transection. *Exp Neurol* 1985, 87:326-333.
- Politis MJ. Specificity in mammalian peripheral nerve regeneration at the level of nerve trunk. *Brain res* 1985, 328:271
- Pons TP, Garrathy PE, Ommaya AK, Kaas JH, Taul E, Mishkin M. Massive cortical reorganization after sensory deafferentation in adult macaques. *Science* 1991, 252:1857-1860.
- Popratiloff A, Kharazia VN, Weinberg RJ, Laonipon B, Rustioni A. Glutamate receptors in spinal motoneurons after sciatic nerve transection. *Neurosci* 1996, 74:953-958
- Popratiloff A, Weinberg RJ, Rustioni A. AMPA receptor subunits underlying terminals of fine-caliber primary afferent fibers. *J Neurosci* 1996, 16(10):3363-3372
- Popratiloff A, Weinberg RJ, Rustioni A. AMPA receptors at primary afferent synapses in substantia gelatinosa after sciatic nerve section. *Eur J Neurosci* 1998, 10:3220-3230
- Povlsen B, Hansson T. Functional evaluation of regenerated and misrouted axons to glabrous and hairy skin of the rat hind foot after sciatic neurectomy and suture. *Exp Neurol* 1995, 132:99-194
- Povlsen B, Hildebrand C, Stankovic N. Functional projections of sensory lateral plantar and superficial peroneal nerve axons to glabrous and hairy skin of the rat hindfoot after sciatic nerve lesions. *Exp neurol* 1994, 129-135.
- Povlsen B, Hildebrand C, Wisenfeld-Hallin Z, Stankovic N. Functional projection of regenerated rat sural nerve axons to hindpaw skin after sciatic nerve lesions. *Exp Neurol* 1993, 119:99-106
- Prats-Galino A, Puigdemívol-Sánchez A, Ruano-Gil D, Molander C. Representations of hindlimb digits in rat dorsal root ganglia. *J Comp Neurol* 1999, 408:137-145
- Pratt CA, Jordan LM. Ia inhibitory interneurons and Renshaw cells as contributors to the spinal mechanisms of fictive locomotion. *J Neurophysiol* 1987, 57:56-71
- Proske U. The mammalian muscle spindle. *News Physiol Sci* 1997, 12:37-42
- Purves D. Functional and structural changes in mammalian sympathetic neurones following interruption of their axons. *J Physiol (Lond)* 1975, 252: 429-463.
- Rafuse V, Gordon T. Incomplete rematching of nerve and muscle properties in motor units after extensive nerve injuries in cat hindlimb muscle. *J Physiol* 1998, 509.3:909-926

- Raivich G, Gehrmann J, Kreutzberg GW. Increase of macrophage colony stimulating factor and granulocyte macrophage in the regenerating rat facial nucleus. *J Neurosci Res* 1991, 30:682-686
- Raivich G, Kreutzberg GW. Pathophysiology of glial growth factor receptors. *Glia* 1994, 11:129-146.
- Raivich G, Kreutzberg GW. Peripheral nerve regeneration: role of growth factors and their receptors. *J Devl Neuroscience* 1993, 3: 311-324.
- Rall W. Distinguishing theoretical synapting potentials computed for different soma-dendritic distributions of synaptic input. *J Neurophysiol* 1967, 30:1138-1168
- Ramón y Cajal S. Estudios sobre la degeneración y regeneración del sistema nervioso. Tomo I. Degeneración y Regeneración de los nervios. Madrid 1913.
- Reichardt LF, Tomaselli KJ. Extracellular matrix molecules and their receptors: functions in neural development. *Ann Rev Neurosci* 1991, 14: 531-570.
- Ren B, Linderorth B, Meyerson B. Effects of spinal cord stimulation on the flexor reflex and involvement of supraspinal mechanisms: an experimental study in mononeuropathic rats. *J Neurosurg* 1996. 84:244-249
- Rende M, Granato A, Manaco ML, Zelagno G, Toesca. Accuracy of reinnervation by peripheral nerve axons regenerating across 10 mm gap within an impermeable chamber. *Exp neurol* 1991; 111:332-339.
- Reynolds ML, Woolf CJ. Reciprocal schwann cell-axon interactions. *Curr Opinion Neurobiol* 1993, 3: 683-693.
- Robitaille R. Purinergic receptors and their activation by endogenous purines at perisynaptic glial cells of the frog neuromuscular junction. *J Neurosci* 1995, 15: 7121-7131.
- Rodríguez FJ, Gómez N, Labrador RO, Butí M, Ceballos D, Cuadras J, Verdú E, Navarro X. Improvement of regeneration with predegenerated nerve transplants in silicone chambers. *Rest Neurol Neurosci* 1999a, 12:65-79
- Rodríguez FJ, Gómez N, Perego G, Navarro X. Highly permeable polylactide-caprolactone nerve guides enhance peripheral nerve regeneration through long gaps. *Biomaterials* 1999b, 20:1489-1500
- Rosén B, Lundborg G, Dahlin LB, Holmberg J, Karlson B. Nerve repair. Correlation of restitution of functional sensibility with specific cognitive capacities. *J hand Surg (Br)* 1994, 19B: 452-458
- Rosenberg ME, Silkensen J. Clustering: physiologic and neurophysiologic considerations. *Int J Biochem Cell Biol* 1995, 27:633-645
- Rost FWD. Fluorescence Microscopy. Volumes I/II. Cambridge University Press 1995, Cambridge.
- Rushton R, Steinberg H, Tinson C. Effects of a single experience on subsequent reactions to drugs. *Brit J Pharmacol* 1962, 20:99-105
- Sadjadpour K. Postfacial palsy phenomena: faulty nerve regeneration or ephaptic transmission? *Brain Res* 1975, 95:403-406.
- Sakai J, Honmou O, Kocsis J, Hashi K. The delayed depolarization in rat cuaneous aferent axons is reduced followng nerve transection and ligation, but nit crush: implications for injury-induced axinal Na⁺ channel reorganization. *Muscl Nerv* 1998, 21:1040-1047
- Sandkühler J, Xianguo L. Induction of long-term potentiation at spinal synapses by noxious stimulation or nerve injury. *Eu J Neurosci* 1998, 10:2476-2480
- Sanes JN, Suner S, Donoghue JP. Dynamic organization of primary motor cortex output to target muscles in adult rats. I. Long-term patens of reorganization following motor or mixed peripheral nerve lesions. *Exp Brain Res* 1990, 79:479-491.
- Sanna PP, Celio M, Bloom F, Rende M. Presumptive Renshaw cells contain decreased calbindin during recovery from sciatic nerve lesions. *Proc Natl Sci* 1993, 90:3048-3042
- Sato A, Sato Y, Suzuki H. Aging effects on conduction velocities of myelinated and unmyelinated fibers of peripheral nerves. *Neurosci Lett* 1985. 53:15-20
- Scaravilli F. The influence of distal environment on peripheral nerve regeneration across a gap. *J Neurocytol* 1984, 13:1027-1041

- Schlaepfer WW, Bunge RP. Effects of calcium ion concentration on the degradation of amputated axons in tissue culture. *J Cell Biol* 1973, 59: 456-470.
- Schlaepfer WW. Calcium-induced degeneration of axoplasm in insulated segments of rat peripheral nerve. *Brain Res* 1974, 69: 203-215.
- Schmalbruch H. Fiber composition of the rat sciatic nerve. *Anat Record* 1986, 215:71-81.
- Schmelzer JD, Zochodne DW, Low PA. Ischemic and reperfusion injury of rat peripheral nerve. *Proc Natl Acad Sci USA* 1989, 86: 1639-1642.
- Schouenborg J, Holmberg H, Yu Y, Weng HR. The organization of spinal nociceptive withdrawal reflexes depends on ontogenetic learning in the rat. *Soc Neurosci Abst* 1994, 20:547 (232.2)
- Schouenborg J, Weng HR, Kalliomäki, Holmberg H. A survey of spinal dorsal horn neurones encoding the spatial organization of withdrawal reflexes in the rat. *Exp Brain Res* 1995, 106:19-27
- Schuurman T, Klein H, Beneke M, Traber J. Nimodipine and motor deficits in the aged rat. *Neurosci Res Comm* 1987, 1(1):9-15
- Scott JJ, Panesar G. Small-signal sensitivity of muscle spindles reinnervated after long-term denervation. *Brain Res* 1995, 671:325-328
- Scott JJ. Regeneration of gamma-fusimotor axons after nerve-freeze injury in the cat. *Brain Res* 1985, 348:159-162.
- Seburn KL, Cope TC. Short Term afferent axotomy increases both strength and depression at Ia-motoneuron synapses in rat. *J Neurosci* 1998, 18(3):1142-1147
- Seckel BR, Chu TH, Niyilas E, Sidman R. Nerve regeneration through synthetic biodegradable nerve guides. Regulation by the target organ. *Plast Reconstr Surg* 1984, 74:171-181.
- Seckel BR, Ryan SE, Gagne RG, Ho-Chiu T, Watkins E. Target specific nerve regeneration through a nerve guide in the rat. *Plast Reconstr Surg* 1986, 78:793
- Seddon HJ. Three types of nerve injury. *Brain* 1943, 66: 237-288.
- Selzer ME. Regeneration of peripheral nerve. En Sumner AJ (ed): *The physiology of peripheral nerve disease*. Saunders, Philadelphia, 1980, pp.358-431.
- Sendtner M, Holtmann B, Kolbeck R, Thoenen H, Barde YA. Brain-derived neurotrophic factor prevents the death of motoneurons in newborn rats after nerve section. *Nature* 1992, 360: 755-759.
- Shenag JM, Shenag SM, Spira M. Reliability of sciatic function index in assessing nerve regeneration across a 1 cm gap. *Microsurg* 1989, 10:214-219
- Sherrington CS. Flexion-reflex of the limb crossed extension-reflex, and reflex stepping and standing. *J Physiol (London)* 1910, 40:28-121
- Sherrington CS. *Integrative action of the nervous system*. Scribner's 1906, New York
- Shindo M, Yanagawa S, Morita H, Hashimoto T. Conditioning effect in single human motoneurons: a new method using the unitary H reflex. *J Physiol* 1994, 481(2):469-477
- Shortland P, Fitzgerald M. Functional connections formed by saphenous nerve terminal sprouts in the dorsal horn following neonatal sciatic nerve section. *European J Neurosci* 1991, 3:383-396
- Shortland P, Fitzgerald M. Neonatal sciatic nerve section results in a rearrangement of the central terminals of saphenous and axotomized sciatic nerve afferents in the dorsal horn of the spinal cord of the adult rat. *Eur J Neurosci* 1994, 6:75-86
- Shortland P, Woolf CJ. Chronic peripheral nerve section results in a rearrangement of the central axonal arborizations of axotomized Aβ primary afferents in the rat spinal cord. *J Comp Neurol* 1993, 330:65-82
- Simonsen EB, Dhyre-Poulsen P. Amplitude of the human soleus H reflex during walking and running. *J Physiol* 1999, 515(3):929-939
- Siuciak JA, Altar CA, Wiegand SJ, Lindsay RM. Antinociceptive effect brain-derived neurotrophic factor and neurotrophin-3. *Brain Res* 1994, 633:326-330
- Skene JHP. Axonal growth-associated proteins. *Annu Rev Neurosci* 1989, 12: 127-156.
- Smith KJ, Hall M. Nerve conduction during peripheral demyelination and remyelination. *J Neurol Sci* 1980, 48: 201-219

- Snider WD, Elliot JL, Yan Q. Axotomy-induced neuronal death during nervous system development. *J. Neurosci Res* 1992, 37:278-286
- Son Y-J, Trachtenberg T, Thompson WJ. Schwann cells induce and guide sprouting and reinnervation of neuromuscular junctions. *TINS* 1996, 19: 280-285.
- Spengler F, Godde B, Dinse HR. Effects of ageing on topographic organization of somatosensory cortex. *NeuroReport* 1995, 6:469-473.
- Stevens LM, Landis SC. Development and properties of the secretory response in rat sweat glands: relationship to the induction of cholinergic function in sweat gland innervation. *Dev Biol* 1987, 123: 179-190.
- Stoll G, Griffin JW, Li CY, Trapp BD. Wallerian degeneration in the peripheral nervous system: participation of both Schwann cells and macrophages in myelin degradation. *J Neurocytol* 1989, 18: 671-683.
- Streppel M, Angelov DN, Guntinas-Lichius O, Hilgers RD, Rosenblatt JD, Stennert E, Neiss WF. Slow axonal regrowth but extreme hyperinnervation of target muscle after suture of the facial nerve in aged rats. *Neurobiol Ageing* 1998, 19(1):83-88
- Strittmatter SM, Valenzuela D, Vartanian T, Sudo Y, Zuber MX, Fishman MC. Growth cone transduction: G_o and GAP-43. *J Cell Sci* 1991, suppl 15: 27-33.
- Sugimoto T, Bennett GJ, Kajander KC. Transsynaptic degeneration in the superficial dorsal horn after sciatic nerve injury: effects of a chronic constriction injury, transection, and strychnine. *Pain* 1990, 42:205-213.
- Sumner AJ. Aberrant peripheral nerve regeneration. In *AAEE Symposium*, 1989, pp 35-36.
- Sumner AJ. Aberrant reinnervation. *Muscle Nerve* 1990, 13: 801-803.
- Sumner BEH, Sunderland S. Quantitative electron microscopy on the injured hypoglossal nucleus in the rat. *J Neurocytol* 1973, 2: 315-328.
- Sumner BEH, Watson WE. Retraction and expansion of the dendritic tree of motor neurones of adult rats induced in vivo. *Nature* 1971, 233: 273-275.
- Sumner BEH. A quantitative analysis of the response of presynaptic boutons to postsynaptic motor neurin axotomy. *Exp Neurol* 1975, 46:605-615.
- Sumner BEH. Responses in the hypoglossal nucleus to delayed regeneration of the transected hypoglossal nerve: a quantitative ultrastructural study. *Exp Brain Res* 1977, 29:219-231.
- Sumner BEH. Ultrastructural data with special reference to bouton glial relationships from the hypoglossal nucleus after a second axotomy of the hypoglossal nerve. *Exp Brain Res* 1979, 36:107-118
- Sunderland S. A classification of peripheral nerve injuries producing loss of function. *Brain* 1951, 74: 491-516.
- Sunderland S. Factors influencing the course of regeneration and the quality of the recovery after nerve suture. In: *Nerve and nerve injuries*. Sunderland S (ed). Churchill Livingstone, Edinburgh 1978, pp. 569-592.
- Sunderland S. *Nerve injuries and their repair. A critical appraisal*. Churchill Livingstone, Edinburgh y London 1991.
- Svendsen F, Tjolsen A, Hole K. LTP of spinal Ab-fibre evoked responses after electrical sciatic nerve stimulation. *NeuroReport* 1997, 8:3427-3430
- Svensson M, Aldskogius H. Evidence for activation of the complement cascade in the hypoglossal nucleus following peripheral nerve injury. *J Neuroimmunol* 1992, 40:99-110
- Svensson M, Aldskogius H. Regeneration of hypoglossal nerve axons following blockade of the brain inhibits the microglial cell proliferation after hypoglossal nerve injury. *Glia* 1993, 7:286-298.
- Svensson M, Ericksson NP, Liu L, Persson J, Aldskogius H. The response of microglia to axon injury. A Ling EA, Tan K, Tan CBC (eds.). *Topical issues in microglial research*. Singapore Neuroscience Association 1996, Singapore.
- Svensson M, Eriksson NP, Aldskogius H. Evidence for activation of astrocytes via reactive microglial cells following hypoglossal nerve transection. *J Neurosci Res* 1993, 35:373-381

- Swett J, Hong CZ, Miller P. All peroneal motoneurons of the rat survive crush injury but some fail to reinnervate their original targets. *J Comp neurol* 1991, 304:234-252.
- Swett J, Hong CZ, Miller PG. Most dorsal root ganglion neurons of the adult rat survive nerve crush injury. *Somatosens Motor Res* 1995, 12(3-4):177-189.
- Swett JE, Wikholm RP, Blanks R, Swett A, Conley L. Motoneurons of the sciatic nerve. *Exp neurol* 1986, 93:227-252
- Swett JE, Woolf CJ. Somatotopic organization of primary afferents terminals in the superficial laminae of the dorsal horn of the rat spinal cord. *J Comp Neurol* 1985, 231:66-77
- Takahashi Y, Nakajama Y, Sakamoto T. Dermatome mapping in the rat hindlimb by electrical stimulation of the spinal nerves. *Neurosci Lett* 1994, 168:85-88.
- Tal M, Wall P, devor M. Myelinated afferent fiber types that become spontaneously active and mechanosensitive following nerve transection in the rat. *Brain Res* 1999, 824:218-223.
- Tanaka K, Zhang QL, Webster HF. Myelinated fiber regeneration after sciatic nerve crush: morphometric observations in young adult and aging mice and effects of macrophage suppression and conditioning lesions. *Exp Neurol* 1992, 118: 53-61.
- Tanelian DL, Barry MA, Johnston SA, Le T, Smith GM. Semaphorin III can repulse and inhibit adult sensory afferents in vivo. *Nat Med* 1997, 3:1398-1401.
- Tessier-Lavigne M, Goodman CS. The molecular biology of guidance. *Science* 1996, 274:1123-1133
- Tetzlaff W, Graeber MB, Bisby MA, Kreutzberg GW. Increased glial fibrillary acidic protein synthesis in astrocytes during retrograde reaction of the rat facial nucleus. *Glia* 1988, 1:90-96
- Thomas PK, King RHM. The degeneration of unmyelinated axons following nerve section: an ultrastructural study. *J Neurocytol* 1974, 3: 497-512.
- Thomson FJ, Reier PJ, Lucas CC, Parmer R. Altered patterns of reflex excitability subsequent to contusion injury of the rat spinal cord. *J Neuropysiol* 1992, 68(5):1473-1486
- Tötösy de Zepetnek J, Gordon T, Stein RB, Ven Zung Hor. Comparison of force and EMG measures in normal and reinnervated tibialis anterior muscles of the rat. *Can J Physiol Pharmacol* 1991, 69:1774-1783
- Tresch M, Saltiel P, Bizzi E. The construction of movement by the spinal cord. *Nature Neurosci* 1999, 2(2):162-167.
- Valls-Solé J, Tolosa E, Pujol M. Myokymic discharges and enhanced facial nerve reflex responses after recovery from idiopathic facial palsy. *Musc Nerv* 1992, 15:37-42
- Valls-Solé J, Tolosa E. Blink reflex excitability cycle in hemifacial spasm. *Neurology* 1989, 39:1061-1066
- Van Meeteren N, Brakee JH, Helders PJ, Gispen WH. The effect of exercise training on functional recovery after sciatic nerve crush in the rat. *J Periph Nerv Syst* 1998, 3(4):277-82
- Verdú E, Butí M, Navarro X. Functional Changes of the peripheral nervous system with aging in the mouse. *Neurobiol Aging* 1996, 17:73-77
- Verdú E, Navarro X. Comparison of immunohistochemical and functional reinnervation injury of skin and muscle after peripheral nerve injury. *Exp Neurol* 1997, 146:187-198.
- Verdú E, Navarro X. The role of Schwann cell in nerve regeneration. A Nieto-Sampedro M, Castellano B, Gonzalez B (eds.). *Understanding Glial Cells* Kluwer Academic Publishers 1998, pp. 319-359, Boston.
- Verdú E. Variaciones de la capacidad regenerativa del sistema nervioso periférico con el envejecimiento. Tesis doctoral. Universitat Autònoma de Barcelona 1994, Bellaterra
- Verge VMK, Gratto KA, Karchewski LA, Richardson PM. Neurotrophins and nerve injury in the adult. *Phil Trans Roy Soc Lond* 1996, 351: 423-430.
- Vilches J, Rodríguez FJ, Verdú E, Valero A, Navarro X. Changes in cholinergic responses of sweat glands during denervation and reinnervation. *J Auton Nerv Syst* 1999, 74:134-142.
- Vilches J. Evaluación de la función sudomotora normal y en procesos de denervación y reinnervación. Tesis Doctoral, Universitat Autònoma de Barcelona, Bellaterra, 1997.

- Von Bartheld C, Cunningham D, Rubel E. Neuronal tracing with DiI: Decalcification, cryosectioning and photoconversion for light and electron microscopic analysis. *Journal of Histochemistry and Cytochemistry*, 1990 38(5): 727-733.
- Walker JL, Resig P, Guarnieri S, Siskin B, Evans J. Improved footprint analysis using video recordings to assess functional recovery following injury to the rat sciatic nerve. *Rest Neurol Neurosci* 1994, 6:189-193
- Wall J, Felleman DJ, Kaas J. Recovery of normal topography in the somatosensory cortex of monkeys after nerve crush and regeneration. *Science* 1983, 221:771-773.
- Wall J, Kaas J. Long-term cortical consequences of reinnervation errors after nerve regeneration in monkeys. *Brain Res* 1986, 372:400-404.
- Wall J, Kaas J. Long-term cortical consequences of reinnervation errors after nerve regeneration in monkeys. *Brain Res* 1986, 372:400-404.
- Wall JT, Kaas JH, Sur M, Nelson RJ, Felleman DJ, Merzenich MM. Functional reorganization in somatosensory cortical areas 3b and 1 of adult monkeys after median nerve repair: Possible relationship to sensory recovery in humans. *J Neurosci* 1986, 1986:218-233.
- Wall P, Woolf C. Muscle but not cutaneous C-afferent input produces prolonged increases in the excitability of the flexion reflexes in the rat. *J Physiol* 1984, 356:443-458
- Wall PD, Woolf CJ. The brief and prolonged facilitatory effects of unmyelinated afferent input on the rat spinal cord are independently influenced by peripheral nerve section. *Neurosci* 1986, 17(4):1199-1205
- Wall TJ, Felleman DJ, Kaas JH. Recovery of normal topography in the somatosensory cortex of monkeys after nerve crush and regeneration. *Science* 1983, 221:771-773.
- Walport M. Complement. A Roitt M, Brostoff J, Male K, (eds.). *Immunology*. Mosby 1997, St Louis, Baltimore, Boston, Chicago, London, Philadelphia, Sydney, Toronto
- Wasserschaff M. Coordination of reinnervated muscle and reorganization of spinal cord motoneurons after nerve transection in mice. *Brain Res* 1990, 515:241-246.
- Waxman SG. Biophysical mechanisms of impulse conduction in demyelinated axons. *Adv Neurol* 1988, 47:185-213.
- Waxmann SG, The molecular basis for electrogenic computation in the brain: you can't step in the same river twice. *Mol Psychiatry* 1999, 4:222-228
- Weiss P, Hoag A. Competitive reinnervation of rat muscles by their own and foreign nerves. *J Neurophysiol* 1946, 9:413-418
- Weiss P, Taylor AC. Further experimental evidence against neurotropism in nerve regeneration. *J Exp Zool* 1944, 95:233
- Weiss P, Hoag A. Competitive reinnervation of rat muscles by their own and foreign nerves. *J Neurophysiol* 1946, 9:413
- Wendell A, Gibby BS, Koerber, Horch K. A quantitative evaluation of suture and tubulization nerve repair techniques. *J Neurosurg* 1983, 58:574-579
- Wessendorf M. Fluoro-Gold: composition, and mechanisms of uptake. *Brain Res* 199, 553:135-148.
- White FV, Toews AD, Goodrum JF, Novicki DL, Bouldin TW, Morell P. Lipid metabolism during early stages of Wallerian degeneration in the rat sciatic nerve. *J Neurochemistry* 1989, 52: 1085-1092.
- Wiesenfeld-Hallin Z, Kinmann, Aldskogius H. Expansion of innervation territory by afferents involved in plasma extravasation after nerve regeneration in adult and neonatal rats. *Exp Brain Res* 1989, 76:88-96.
- Wiesenfeld-Hallin Z, Kinmann, Aldskogius H. Studies of normal and expansive cutaneous innervation territories of intact and regenerating C-fibres in the hindlimb of the rat. *Agents and Actions* 1988, 25(3/4):260-262
- Wiethölter H, Eckert S, Stevens A. Measurement of atactic and paretic gait in neuropathies of rats based on analysis of walking tracks. *J Neurosci meth* 1990, 32:199-205
- Wilgis EFS, Maxwell GP. Distal digital nerve grafts: clinical and anatomical studies. *J Hand Surg* 1979, 4: 439-441.

- Williams LR, Longo FM, Powell HC, Lundborg G, Varon S. Spatial-temporal progress of peripheral nerve regeneration within a silicone chamber. Parameter for a bioassay. *J Comp Neurol* 1983, 218: 160-170.
- Willis WD. Is central sensitization of nociceptive transmission in the spinal cord a variety of long-term potentiation?. *NeuroReport* 1997, 8:16
- Woolf C, Swett J. The cutaneous contribution to the hanstring flexor reflex in the rat: an electrophysiological and anatomical study. *Brain Res* 1984, 303:299-312
- Woolf CJ, King AE. Dynamic alterations in the cutaneous mechanoreceptive fields of dorsal horn neurons in the rat spinal cord. *J Neurosci* 1990, 10(8):2717-2726
- Woolf CJ, Shortland P, Coggeshall E. Peripheral nerve injury triggers central sprouting of myelinated afferents. *Nature* 1992, 355:75-78
- Woolf CJ, Shortland P, Reynolds M, Ridings J, Doubell T, Coggeshall RE. Reorganization of central terminals of myelinated primary afferents in the rat dorsal horn following peripheral axotomy. *J Comp Neurol* 1995, 360:121-131
- Yakhnitsa V, Linderöth B, Meyerson B. Spinal cord stimulation attenuates dorsal horn neuronal hyperexcitability in a rat model of mononeuropathy. *Pain* 1999, 79:223-233
- Young JB, Rowe JW, Pallotta AP, Sparrow D, Landsberg L. Enhanced plasma norepinephrine response to upright posture and oral glucose administration in elderly human subjects. *Metabolism* 1980, 29: 532-539.
- Zalewski AA. Effects of reinnervation on denervated skeletal muscle by axons of motor, sensory, and sympathetic neurons. *Am J Physiol* 1970, 219:1675-1679
- Zehr EP, Komiyama T, Stein RB. Cutaneous reflexes during human gait: electromyographic and kinematic to electrical stimulation. *J Neurophysiol* 1997, 77:3311-3325.
- Zelená J. Multiple axon terminals in reinnervated pacinian corpuscles of adult rat. *J Neurocytol* 1984, 13:665-684
- Zhang J, Ming J, Donnelly F, Song D, Xue-Jun, Lamotte RH. Axotomy increases the excitability of dorsal root ganglion cells with unmyelinated axons. *J Neurophysiol* 1997, 78: 2790-2794
- Zhang X, Bean J, Wiesenfeld-Hallin Z, Hökfelt T. Ultrastructural studies on peptides in the dorsal horn of the rat spinal cord- IV. Effects of peripheral axotomy with special reference to neuropeptide Y and vasoactive intestinal polypeptide/peptide histidine isoleucine. *Neurosci* 1995, 64(4):917-941
- Zhao Q, Dahlin LB, Kanje M, Lundborg G, Lu SB. Axonal projections and functional recovery following fascicular repair of the rat sciatic nerve with Y-tunnelled silicone chambers. *Rest Neurol Neurosci* 1992a, 4:13-19.
- Zhao Q, Dahlin LB, Kanje M, Lundborg G. Specificity of muscle reinnervation following repair of the transected sciatic nerves: A comparative study of different repair techniques in the rat. *J Hand Surg (Br)* 1992b, 17(B):257-261.
- Zhao Q, Dahlin LB, Kanje M. Reinnervation of muscles in rats after repair of transected sciatic nerves with Y-shaped and X-shaped silicone tubes. *Scand J Plast Reconstr Hand Surg* 1992c, 26(3):101-106.
- Zhu PH, Vrbová G. The role of Ca^{2+} in the elimination of polyneuronal innervation of rat soleus muscle fibers. *Eur J Neurosci* 1992, 4:433-437.
- Zigmond RE, Hyatt-Sachs H, Mohny RP, Schreiber RC, Shadiak AM, Sun Y, Vaccariello SA. Changes in Neuropeptide Phenotype after axotomy and the role of Leukaemia inhibitory factor. *Perspectives Dev. Neurobiol*, 1997.
- Ziskind-Conheim . NMDA receptors mediate poly and monosynaptic potentials in motor neurons of the rat embryos. *J Neurosci* 1990, 10:125-135.

0



Servei de Biblioteques

Reg. 1500459751

Sig. T0AB/4914

Ref. 12500

