

Patricia García Sans

Final Degree Project – June 2019



Figure 1. Location of the pancreas. 1. Duodenal lobe of the pancreas; 2. Descending part (duodenum); 3. Caudal duodenal flexure; 4. Ascending part (duodenum); 5. Muscular stomach; 6. Liver; 7. Gall bladder.

## INTRODUCTION

**Objectives:** to contribute to the morphological characterization of the pancreatic islets of the Japanese quail. To estimate the size of the pancreas and its  $\beta$  cell mass, to study the distribution pattern of the  $\beta$  cells inside the islets and to determine the relative number and size of the pancreatic islets.

The pancreas is a mixed gland that is located on the right side of the abdominal cavity, on the mesenteric border of the duodenum. In most birds, it consists of four lobes: ventral, dorsal, third and splenic, together with the main pancreatic ducts leading to the ascending duodenal loop.

The exocrine tissue is formed by glandular acini that drain into a branched duct system. The endocrine component is organized into islets of different shapes and sizes. Within the islets,  $\beta$  cells produce insulin,  $\alpha$  cells produce glucagon,  $\delta$  cells produce somatostatin, PP cells synthesize pancreatic polypeptide and  $\epsilon$  cells that produce ghrelin. Insulin is an hypoglycaemic hormone because it stimulates the uptake of glucose especially by hepatocytes, adipocytes and skeletal muscle cells, thus reducing its concentration in the blood.




Figure 2. Structure of the quail pancreas. Histological section of pancreatic tissue. Hematoxylin-Eosin stain (200X). 1. Pancreatic acini; 2. Pancreatic islets; 3. Intralobular conduit.

## MATERIALS & METHODS

### Measurements

- 12 quails (*Coturnix coturnix japonica*) of 5 weeks of age (6 males, 6 females).
- Blood glucose levels and body weight were determined at 5, 6 and 7 weeks of age.
- In 7 of the quails (3 males, 4 females) systematic morphometric study of the pancreas was performed.

- Blood glucose measurement (glucometer Accu-Chek Aviva, Roche Diabetes Care).
- Hematoxylin-eosin staining in paraffin sections.
- Immunohistochemistry for the detection of insulin and  $\beta$ -cells using HRP and DAB system.
- Microscope (Nikon Eclipse E800) and ACT-1 software for image capture.

- Point counting morphometry

$$\text{Tissue mass A} = \frac{\text{Dots on tissue A}}{\text{Dots on all tissues}} \times \text{Pancreas weight}$$

- Results were expressed a mean  $\pm$  SEM, and comparisons validated with Student's t test.

## RESULTS & DISCUSSION




Figure 3. Immunodetection of insulin in pancreatic quail  $\beta$  cells by development with DAB. A) 200X. 1.  $\beta$  cells; 2. Pancreatic acinus.

Table 2. Estimated mass of the  $\beta$  cells, exocrine tissue, ductal cells and other tissues (shown in percentages).

	$\beta$ -Cell Mass (mg)	Exocrine Tissue Mass (mg)	Ductal Cell Mass (mg)	Other Tissues Mass (mg)
$\textcircled{M}$ N=3	6.39 $\pm$ 0.64 (1.30%)	441.21 $\pm$ 52.0 1 (89.17%)	11.97 $\pm$ 0.99 (2.47%)	34.44 $\pm$ 6.36 (7.06%)
$\textcircled{F}$ N=4	6.71 $\pm$ 0.68 (0.64%)	997.67 $\pm$ 62.82 (95.86%)	12.24 $\pm$ 1.57 (1.20%)	23.63 $\pm$ 1.63 (2.30%)
Total N=7	6.57 $\pm$ 0.44 (0.92%)	759.19 $\pm$ 118. 96 (92.99%)	12.12 $\pm$ 0.92 (1.75%)	28.26 $\pm$ 3.36 (4.34%)

The percentage of  $\beta$  cells in male quail is significantly higher than females, being approximately 1.3%, This percentage is similar to that observed in male mice (0.5-1%)

Table 3. Distribution of the number of pancreatic islets according to their size (shown in percentages).

	Small islets (<50 $\mu\text{m}$ )	Medium islets (50-149 $\mu\text{m}$ )	Large islets (>150 $\mu\text{m}$ )	Total
$\textcircled{M}$ N=3	492.67 $\pm$ 141.1 3 (81.06%)	97.33 $\pm$ 2.91 (18.32%)	3.67 $\pm$ 0.88 (0.62%)	593.67 $\pm$ 139.6 4 (100%)
$\textcircled{F}$ N=4	157.75 $\pm$ 10.23 (68.09%)	73.75 $\pm$ 12.40 (31.17%)	1.75 $\pm$ 0.25 (0.74%)	233.25 $\pm$ 17.23 (100%)
Total N=7	301.29 $\pm$ 86.33 (73.65%)	83.86 $\pm$ 8.24 (25.67%)	2.57 $\pm$ 0.53 (0.69%)	387.71 $\pm$ 90.40 (100%)

## CONCLUSIONS

- Males and females presented similar and stable blood glucose levels throughout the follow-up.
- Body weight was 30% higher in females than in males.
- The pancreas weight in females was twice than in males, and represents 0.29% and 0.19% of the total body weight, respectively.
- The general structure of the quail pancreas is similar to that observed and reported in mammals.

- The percentage of  $\beta$ -cells relative to the total pancreas weight in males was twice than in females. Thus, a similar  $\beta$ -cell mass was observed in males and females.
- The morphology of the islets is irregular and their  $\beta$  cells are also irregularly distributed within the islet.
- Then number of islets observed is higher in males than in females, and the size distribution seems to be different in both.