

A hopeful future

Degree in Biology June 2020

Foliar epiphytes as bioindicators of *Posidonia oceanica* state of preservation


Júlia Ribalta Alet

Out of the 5 marine phanerogams found in the Mediterranean Sea, Posidonia oceanica (L.) Delile, 1813 is the only one which is endemic.

It forms extensive seagrass meadows which are very important at:

- Ecological level \rightarrow to act as a substrate, refuge or place of settlement for many species.
- Economical level \rightarrow to provide a wide variety of ecosystem services.

Unfortunately, they are subjected to a list of threats which put at risk the diversity associated with them.




AIMS OF THE STUDY

CHARACTERISATION

- To see if a Caulerpa spp. community has less biodiversity associated than a healthy P. oceanica community.
- To analyse the foliar epiphytes of *P. oceanica* in a good state of preservation vs. a degraded one.
- To understand the trend of *P. oceanica* meadows taking into account the current situation.

Bibliographical information was searched and classified into 2 groups: those which described a typical or healthy meadow and those which referred to a degraded one.




In= Indeterminate Ge= Generalist

Figure 1. Methodology diagram used for the habitat characterisation and comparison.

Only foliar epiphytes species were selected and they were classified according to 2 established criteria. Finally, the two communities were characterised and the foliar epiphytes species of the two habitats compared.

≈ 50 % ← 6 Pre ≈ 22 % 27 spp. ≈ 22 %



Phaeophyceae; 5 spp. Bryozoa; 5 spp. (4) Anthozoa; 2 spp.

Annelida; 3 spp. 7 Ascidiacea; 2 spp. (8) Clorophyta; 1 sp.

≈ 30 % 13 spp. ≈ 15 % ≈ 30 % P. oceanica meadow in a degraded state of preservation Rhodophyta; 5 spp. Phaeophyceae; 3 spp. 3 Bryozoa; 5 spp.

Figure 2. Results diagram with the taxonomic groups which characterise a healthy (left) and a degraded (right) P. oceanica meadow.


CURRENT SITUATION

Least concern at IUCN Red List

36 % Declining 20 % Increasing 44 % Showing no change Climate change (个 temperature & 个 extreme events).

Water quality degraded, anchoring, trawling & coastal modification. Geological dynamics change. Non-native macroalgae invasions.

COMPARISON



In Catalonia, P. oceanica meadows have different preservation states depending on the region.

classified by taxonomic groups in a healthy habitat and

a degraded habitat.

However, in general, they are showing no change in trajectory due to the management measures taken.

MANAGEMENT RECOMMENDATIONS

classified by taxonomic groups.

THREATS

- More scientific research should be done and it would be convenient to standardise sampling methods.
 - In order to preserve these habitats is extremely important to:
 - create networks of nature protection areas
 - keep local monitoring of the *P. oceanica* meadows
 - carry out environmental initiatives

CONCLUSIONS

- It can not be affirmed that Caulerpa spp. "meadows" have less biodiversity.
- As an habitat dominated by P. oceanica degrades, the composition of species changes: characteristic species decrease and generalist species increase.
- There is not a generalized trend of *P. oceanica* meadows to regression and the main degradation causes are the anthropogenic ones.

SELECTED REFERENCES:

- Ballesteros, E., García, A., Lobo, A., & Romero, J. (1984). L'alguer de *Posidonia oceanica* de les Illes Medes. En I. Olivella (Ed.), Els Sistemes naturals de les Illes Medes (p. 739-759).
- Romero, J. (2004). *Posidònia: els prats del fons del mar* (Escola del Mar, Àmbit de Medi Ambient Sostenibilitat i Habitatge, & Ajuntament de Badalona (ed.); 1a ed.).
- De los Santos, et al. (2019). Recent trend reversal for declining European seagrass meadows. Nat. Commun., 10(1), 1-8.