

TABLE 1. Percentages of main biosecurity measures applied by transporters (n=23)
(VC = Very common, C = Common, O = Occasional, R = Rare, N = Never)

Farm	VC	C	O	R	N
Enters the perimeter of the farm (%)	69	13	-	9	9
Use of loading and unloading bay (%)	22	39	22	13	4
Enters the pen where the animals are kept (%)	69	13	-	9	9
Maintains contact with the animals that stay on the farm (%)	61	17	4	9	9
Loading and unloading					
The footwear is replaced with own boots (%)	96	-	4	-	-
The footwear is replaced with boots provided by the farm (%)	17	-	-	-	83
Uses footwear covers (%)	4	-	-	-	96
Changes clothes for work (%)	82	-	9	-	9
Climbs into the cabin with overall and boots on (%)	19	10	10	-	61
Cleaning and disinfection					
Truck cleaning between two transports to different customer's farms (%)	74	9	9	4	4
Truck cleaning between two transports to farms of the same customer (%)	69	9	9	4	9
Truck disinfection between two transports to different customer's farms (%)	61	9	22	-	9
Truck disinfection between two transports to farms of the same customer (%)	56	9	22	-	13
Cleaning the cabin between two farm transports (%)	-	-	-	5	95




FIG. 1. Map of the location of surveyed transporters (n=23)

TABLE 2. Percentages of the completion of the load on one or more farms for the different types of cattle transported

	Calves for rearing (n=13)	Restocking (n=4)	Calves for fattening (n=11)	Cattle to slaughter (n=22)	Culling animals (n=2)
Complete the load on 1 farm (%)	76,92	50	72,73	45,45	0
Complete the load on >1 farm (%)	23,08	50	27,27	54,55	100




FIG. 2. Frequencies of time taken by transporters to clean and disinfect their transport vehicle (n=23)

